



## ANTHELMINTIC ACTIVITY OF THIRIKADUGU CHOORANAM AND ITS INGREDIENTS

M.S.SHREE DEVI<sup>1\*</sup>, B.SAMPATH KUMAR<sup>2</sup>

1. Ph.D., Scholar Vel's University, Chennai. 2. Professor, Dept of Pharmacology, Tirunelveli., India  
E mail: shreemd@gmail.com

### ABSTRACT

**Aim:** The alcoholic extract of Thirikadugu Chooranam and its ingredients were evaluated for anthelmintic activity. **Introduction:** The dried fruits of *Piper nigrum* L. (Piperaceae), *Piper longum* L. (Piperaceae) and rhizome of *Zingiber officinale* Roscoe. (Zingiberaceae) were powdered and mixed together in equiproportions to get a polyherbal formulation, thirikadugu chooranam. All these three ingredients are spicy, commonly used in our daily diet, also well known for their tremendous therapeutic potential, since from the Vedic period. **Materials and Methods:** The alcoholic extract of Thirikadugu chooranam and its ingredients were screened for preliminary phytochemical studies and also tested for anthelmintic activity against *Pheritima posthuma* and recorded the time taken for induction of paralysis and death. Piperazine citrate (10 mg/ml) was included as standard reference and distilled water as control. **Result and Discussion:** The results demonstrated that, the extracts of Thirikadugu chooranam and its plant ingredients showed the presence of alkaloids, flavonoids, tannins, lignins, saponins and steroids, but *P. nigrum* has shown negative for steroid tests, *P. longum* for saponins and tannins, *Z. officinale* for Labat test (for Lignans). Where as, the Thirikadugu chooranam showed positive for all the above said tests. All these test samples were also exhibited potent anthelmintic activity, but the highest activity was noticed in Thirikadugu chooranam, this might be due to the multifunctional effect of all the three plant ingredients of Thirikadugu chooranam. **Conclusion:** Based on the above results, it is confirmed that, combination of *Piper nigrum*, *Piper longum* and *Zingiber officinale* in Thirikadugu chooranam offered promising anthelmintic effect than using the ingredients alone.

**Key words:** Thirikadugu chooranam, Piperazine citrate, *Pheritima posthuma*.

### INTRODUCTION

Thirikadugu chooranam is one of the traditional polyherbal preparations, made up of combination of three important spicy materials, such as *Piper nigrum* L (Piperaceae), *Piper longum* L. (Piperaceae) and *Zingiber officinale* Roscoe. (Zingiberaceae). All these plant materials are used world wide as spices. They are also used as important ingredients in folklore medicine in many Asian countries. However, the consumption of these spices would exert several health beneficial effects by the virtue of their innumerable therapeutic potentials, such as fever, asthma, cold, cough and other general health disorders<sup>1-6</sup>. There are voluminous research carried out in this particular stream, a brief review of the notable work is highlighted and the combination of two or more these formulations were tested for anthelmintic activity against *Pheritima posthuma*<sup>7</sup>, aqueous extract of fruits of *Terminalia chebula* Retz.<sup>8</sup>. But so far no clinical trials are made on this compound poly herbal formulation, Thirikadugu chooranam. Hence, the present study was undertaken to explore the anthelmintic activity of Thirikadugu chooranam and compared its effects to its individual ingredients. The present study is also aimed to establish its clinical validity.

### MATERIALS AND METHODS

The anthelmintic activity was tested on earth worms (*Pheritima posthuma*) using ethanolic(95%) extracts of *Piper longum*, *Piper nigrum*, *Zingiber officinale* and Thirikadugu chooranam. The crude extracts of Thirikadugu chooranam and its ingredients were also screened for preliminary phytochemical studies to find out the occurrence of possible major chemical groups in the given extracts.

#### Collection of Plant materials:

The plant materials of *Piper nigrum* L, *Piper longum* L. along with the fruits were collected from the Agricultural University, Coimbatore, and plant along with rhizomes of *Zingiber officinale* Roscoe. Were collected from the farmers of Coimbatore in the month of October-November. All these plants were authenticated in Department of Botany, Agricultural University, Coimbatore (Tamilnadu, India). The fruits and the rhizomes of respective plants were surface sterilized by using 50% alcohol, then shade dried and powdered for the preparation of Thirikadugu chooranam.

#### Preparation of the Thirikadugu chooranam:

The Thirikadugu chooranam is a fine powder of drugs. It is prepared by mixing equal quantities of the powder of the dried fruits of *Piper*

*nigrum*, *Piper longum* and rhizomes of *Zingiber officinale* and then sieved through muslin cloth. This chooranam is stored in airtight container for further processing<sup>4</sup>.

#### Preparation of the extract:

The 100g of Thirikadugu chooranam and its ingredients were extracted in 95% ethanol at 50 - 60°C in a soxhlet apparatus separately. The extract was concentrated to dryness in a flash evaporator (Buchi type) under reduced pressure and controlled temperature (50 -60°C). The dried 95% of the ethanolic extracts weighed in a required dose and dissolved in known volume of distilled water, separately for further treatment.

#### Phytochemical Evaluation of the Crude Extracts:

Phytochemical screening of the extracts for the presence of secondary metabolites were performed using the following reagents and chemicals: for alkaloids with Mayer's, Wagner's and Dragendroff's reagents, for flavonoids with the use of Mg and HCl, tannins with 1% gelatin and 10%NaCl solutions, for saponins with distilled water<sup>9-13</sup>.

#### Preparation of Standard Solution and Control:

Piperazine citrate powder standard drug [Adani Pharma chem Pvt. Ltd, Gujarat, India] was dissolved in 100ml of normal saline solution to get 1, 2, and 4ml of solution. Normal saline alone was used as control.

#### Experiment Design:

Adult earth worms (*Pheritima posthuma*) were collected (due to their anatomical and physiological resemblance with the intestinal round worm parasites of human beings) Earth worms were thoroughly washed with normal saline to remove the adhering material. Petri dishes of equal size were collected and 20ml of normal saline alone was poured in the first petri dish, 20ml of Piperazine citrate solution of concentration 1, 2 and 4mg/ml were poured in second, third and fourth petri dishes, respectively. Then 20ml (4mg/ml) of the test solutions that is, the ethanolic extracts of *piper nigrum*, *Piper longum*, *Zingiber officinale* and Thirikadugu chooranam were taken in fifth, sixth, seventh and eighth Petri dishes, respectively. Placed six earth worms of nearly equal size in each Petri dish and time taken for the induction of paralysis (motion less) and complete death of earth worms was noted. The experiment was repeated thrice and confirmed the readings<sup>8</sup>.

**Table 1: Distribution of primary and secondary metabolites in Thirikadugu chooranam and its ingredients**

Tests		Piper nigrum	Piper longum	Zingiber officinale	Thirikadugu chooranam
Alkaloids	Mayer's test	+	+	+	+
	Wagner's test	+	+	+	+
	Dragendroff's test	+	+	+	+
Steroids	Salkowski' test	-	+	+	+
	Liebermann and Burchard test	-	+	-	+
Flavonoids	Extract + Mg turnings	+	+	+	+
	Extract + Aqueous	+	+	+	+
	NaOH + Conc H2SO4	+	+	+	+
Saponins	Foam test	+	-	+	+
Tannins	Gelatin test	+	-	+	+
Lignans	Labat test	-	+	-	+
	Lignan test	+	+	+	+

Note : '+' indicates the compound is present in the extract, '-' indicates absent.

**Table 2: Anthelmintic activity of ethanolic (95%) extracts of Piper longum, P.nigrum, Zingiber officinalis, Thirikadugu chooranam and Piperazine citrate**

Sl No	Treatment	Concentration (mg/ml)	Paralysis Time (min)	Death time (min)
01	Normal Saline	0.9% NaCl	No paralysis	No death
02	Piperazine Citrate	01	43.66 ±1.071	63.33 ±0.838
03	Piperazine Citrate	02	30.00 ±0.881	66.66 ±1.071
04	Piperazine Citrate	04	21.33 ±0.509	36.33 ±1.895
05	<i>Piper longum</i> L.	04	113.66 ±1.34*	207.00 ±1.20*
06	<i>Piper nigrum</i> L.	04	67.66 ±0.769	238.66 ±4.71*
07	<i>Zingiber officinale</i> Roscoe.	04	46.00 ±0.881	235.33 ±1.50*
08	Thirikadugu chooranam	04	29.66 ±0.693	90.66 ±1.34*

Note:\*=P<0.05 (significant difference) compared 5th, 6th, 7th and 8th groups with 2nd, 3rd, and 4th groups.

#### Statistical Analysis:

All the data are expressed as mean ± S.E.M. (standard error of the mean). The significance level was determined using the Student's 't' test. A p value of <0.05 was considered statistically significant.

#### RESULTS AND DISCUSSION

The results of the above studies demonstrated that, the alcoholic extract of Thirikadugu chooranam and its individual ingredients possess potent anthelmintic activity with varying magnitudes. But the extract of Thirikadugu chooranam showed highest activity, which is almost equal in effectiveness to the standard Piperazine citrate. The difference in the time taken for induction of paralysis in both Piperazine citrate and Thirikadugu chooranam was insignificant or almost same. However, significant difference was observed when compared the induction of paralysis time of Piperazine with ingredients of chooranam alone. The mode of action for the piperazine is generally by paralyzing parasites, which allows the host body to easily remove or expel the invading organism (Table-2).

The preliminary phytochemical observations of the alcoholic extracts of four different test samples such as Thirikadugu chooranam and its plant ingredients have shown the occurrence of alkaloids, flavonoids, tannins, saponins, lignins and steroids. But *P. nigrum* has shown negative for steroid tests, *P. longum* for saponins and tannins, *Z. officinale* for Labat test (for Lignans). Where as, the Thirikadugu chooranam alone showed positive for all the above said tests. (Table-1). It indicates that, the Thirikadugu chooranam is a mixture all these phyto constituents and interaction all these chemicals might be resulted in synergistically enhanced therapeutic efficacy of anthelmintic activity.

The main finding from this investigation that, the alcoholic extracts all the four test samples produced paralysis of earth worms after 30-120 minutes when treated 4mg/ml concentration of the Thirikadugu chooranam and individual extracts. Moreover earthworms did not get recovery from paralysis even after 2-3 hours of post treatment period. Chemotherapeutic drugs against helminthes infection act mainly through three different mechanisms, such as, disruption of the neuromuscular physiology, blocking the energy metabolism, disrupting the highly efficient reproductive system of the parasites [14]. Several important anthelmintics cause paralysis by disrupting one or the other aspect of neuromuscular system<sup>15</sup>. The paralytic effect of alcoholic extract of *Piper longum* was observed on *Gigantocotyle explanatum* by progressive reduction in the spontaneous muscular activity.

This may be associated with their inhibitory effect on the neuromuscular system of amphistome<sup>16</sup>. *In vivo* anthelmintic activity of ginger was confirmed against gastrointestinal nematodes of sheep, thus justifying the age-old traditional use of this plant in helminth infestation<sup>17</sup>. The oil of *Piper longum* elicited the pronounced effect on the rhythmic movements of *Ascaris* as evidenced from resulting paralysis that occurred between 12 to 15 minutes of exposure with 1:1000 v/v concentration of oil. The essential oil of *Piper longum* and its non-phenolic fraction did not significantly differ in their ability to inhibit the rhythmic motions, even though; minimal motions persisted in case of non-phenolic fraction<sup>18</sup>. Thus, it has happened obviously that, Thirikadugu chooranam is a mixed preparation of all these useful phyto constituents, perhaps the synergistic interaction of alkaloids, flavonoids, tannins, lignins, steroids and other constituents in the extract may impart strong anthelmintic activity to the poly herbal preparation. But the mechanism involved in the interaction between

the different plant extracts remain unclear and should be further evaluated.

#### CONCLUSION

Thirikadugu chooranam was found to possess higher the rate of phyto constituents and promising anthelmintic activity. It is also confirmed that, these spicy products triggers natural immune system to fight against various parasites as well as helminthes. This study would provide the preliminary scientific evidence for the folkloric, ethno-botanical and traditional use of this chooranam for destruction of helminthes / parasites and eliminates from host body and other health benefits.

#### REFERENCES

1. Anonymous. Indian Pharmacopoeia, Government of India, III-Edition, New Delhi, Appendix IV, 1985:90
2. Chopra R.N, Nayar SL, Chopra, IC. In Glossary of Indian Medicinal Plants, Publications and Information Directorate, CSIR, New Delhi, 1992:132, 210
3. George Watt. Dictionary of the economic products of India, Periodical experts, 42-D, Vivek Vihar, Delhi, 1972:217, 247.
4. Rakesh Lodha and Sushil K. Kabra. Bronchial asthma In Scientific Basis for Ayurvedic Therapies, Edited by Lakshmi Chandra Mishra, Published by CRC Press. 2003: 211-215
5. Namjoshi AM. Ayurvedic formulary of India, Controller of Publications, Delhi, 1976:85, 95.
6. Sivarajan VV, Indira Balachandran. Ayurvedic drugs and their plant sources. Published by Mohan Primalini for Oxford and IBH Publishing Company, PVT.LTD, New Delhi. 1996.
7. Dwivedi S, Dwivedi A, Kapadia R, Kaul S. Anthelmintic activity of alcoholic and aqueous extract of fruits of *Terminalia chebula* Retz. Ethanobotanical Leaflets, 2008;12:741-43.
8. Harborne JB. Phytochemical Methods. A guide to modern techniques of plant analysis, 3rd Ed. Chapman and Hall, London, 1998: 235.
9. Sadasivam S, Manickam A. Biochemical methods for Agricultural Sciences, Wiley Eastern Limited., Ansari Road, Daryaganj, New Delhi, 1992:1-20
10. Ogbonnia S, Adekunle AA, Bosa MK, Envuru VN, Evaluation of acute and sub acute toxicity of *Alstonia congensis* Engler (Apocynaceae) bark and *Xylopiiaethiopica* (Dunal)A. Rich (Annonaceae) fruits mixtures used in the treatment of diabetes., African Journal of Biotechnology 2008;7(6):701-705.
11. Nooman A. Khalaf, Ashok K. Shakya, Atif Al-Othman, Zaha El-Agbar, Husni Farah. Antioxidant activity of some common plants. Turk. J. Biol 2008; 32:51-55.
12. Mohd. Nawagish, Ansar SH, Shoaib Ahmad, Preliminary Pharmacognostical Standardisation of *Lawsonia inermis* Linn. Seeds. Research Journal of Botany 2007; 2(3):161-164
13. Geary TG, Klein RD, Vanover L, Bowman JW, Tompson DP.. The nervous system of helminthes as target for drugs. J. Parasitol 1992;78:215 -280.
14. Loukas A, Hotez PJ. Chemotherapy of helminth infections. In Brunton LL, Lazo JS, Parker KL., Goodman and Gilman's, The Pharmacological Basis of Therapeutics, 11th ed, U.S.A: McGraw-Hill Company, 2005:1073 - 1093.
15. Singh TU, Kumar D, Tandan SK, Paralytic effect of alcoholic extract of *Allium sativum* and *Piper longum* on liver amphistome, *Gigantocotyle explanatum*. Indian J. Pharmacol 2008; 40(2): 64-68
16. Iqbal Z, Lateef M, Akhtar MS, Ghayur MN, Gilani AH. In vivo anthelmintic activity of ginger against gastrointestinal nematodes of sheep. J. Ethnopharmacol 2006; 106(2):285-287.
17. Kokate CK, Chaudhari GN, Nimbkar AY. Search for anthelmintics of plant origin: activities of volatile principles of *Acorus calamus* and *Piper longum* against *Ascaris lumbricoides*, Asian Symposium on Medicinal Plants and Spices, Conference, Bangkok (Thailand), 1980:15-19.