

## TO EVALUATE HEPATOPROTECTIVE ACTIVITY OF ROOTS OF CYNODON DACTYLON -AN EXPERIMENTAL STUDY

ANURAG JAIN \*<sup>1</sup>, I.P. JAIN<sup>2</sup>, S.P. SINGH<sup>3</sup>, ASHA AGRAWAL<sup>4</sup>

<sup>1</sup>M.D. Lecturer, Department of Pharmacology & Therapeutics, S.N. Medical College, Agra, <sup>2</sup> & <sup>3</sup> M.D. Professor, Department of Pharmacology & Therapeutics, G.S.V.M. Medical college, Kanpur, <sup>4</sup> M.D. Professor, Department of Pathology, G.S.V.M. Medical college, Kanpur.

Email: dranurag2008@rediffmail.com

Received: 4 January 2013, Revised and Accepted: 22 February 2013

### ABSTRACT

**Objective :** To Evaluate Hepatoprotective Activity of Roots of Cynodon Dactylon in CCl<sub>4</sub> induced hepatotoxicity in albino Rabbits.  
**Materials and Methods:** The study was conducted on 12 healthy albino rabbits of either sex weighing 1.5-2.0 kg, divided into 2 groups. Hepatotoxicity was induced in rabbits by carbon tetra chloride (CCl<sub>4</sub>) 0.05 mg/kg, intraperitoneally. Alcoholic extracts of roots of Cynodon dactylon was administered orally for 20 days from 1 day to day 20 in the doses of 100mg/kg/day with the help of syringe.  
**Results:** Group I: The rise of serum transaminase (p<0.001), serum alkaline phosphatase (p<0.001), serum bilirubin (p<0.001) and decrease in serum albumin (p<0.001) due to hepatotoxic effect of CCl<sub>4</sub> when compared to zero day of same group (p<0.001).  
Group II: Cynodon dactylon extract was able to bring down the level of serum transaminase, serum alkaline phosphatase, serum bilirubin and increased in serum albumin in a statistical highly significant amount (p<0.001), when compared with group I.  
**Conclusion:** Cynodon dactylon extract found to be effective in reducing SGOT, SGPT, ALP, S.bilirubin. Cynodon dactylon extract helps in normalize S.Albumin level. Cynodon dactylon extract had shown protection in restoration of liver function and regeneration of liver cells as observed on histopathology.

**Key words:** hepatoprotective activity, Cynodon dactylon, liver injury

### INTRODUCTION

The liver is the largest internal organ in the body contributing about 2 percent of total body weight<sup>1</sup>, which plays an essential role in the metabolism of foreign substances entering the body. They are known as **Xenobiotics**. The liver has considerable reserve capacity, can often maintain function in spite of significant disease and is one of the few human organs capable of regeneration<sup>1</sup>.

More than 1000 xenobiotics substances are potentially hepatotoxic<sup>2</sup>. The ability of the chemical to produce liver damage in vivo often results from interaction of a series of complex process involved in the uptake, biotransformation and elimination of these potentially toxic compounds.

Conventional drugs used in the treatment of liver disease are often inadequate. It is therefore necessary to search for **alternative drugs** for the treatment of liver diseases to supplement the currently used drugs of limited efficacy and safety.

Hence, the present study is designed to evaluate hepatoprotective effect of Cynodon dactylon in CCl<sub>4</sub> induced hepatotoxicity in experimental animals, supported by histopathological evidences.

### MATERIALS AND METHODS

The present study was conducted in the department of Pharmacology and therapeutics in collaboration with department of Pathology, G.S.V.M. Medical college, Kanpur, after the clearance from Institutional Animal Ethical Committee for Prevention of cruelty and supervision of experiments on animals.

#### Animals

The study was conducted on 12 healthy albino rabbits of either sex weighing 1.5-2.0 kg, divided into 2 groups. The animals were made available in the animal house of Department of Pharmacology & Therapeutics.

Rabbits, also have metabolism similar to human beings. Hepatotoxicity induced in rabbit by carbon tetra chloride (CCl<sub>4</sub>) simulate the symptoms of drug induced hepatitis in human being

without the development of concurrent infections. So, experiment on rabbits correlate well with human subjects.

All the animals were fed normal stock diet for 7 days. During this time the animals got acclimatized to the new environment. All the animals were housed individually in clean cage and maintained under standard conditions (12 hr light and dark cycle, at room temperature 25± 3° C and 35-60% humidity).

#### Drugs Used

Alcoholic extracts of roots of **Cynodon dactylon**

This extracts was administered orally for 20 days from 1 day to day 20 with the help of syringe. Carbon tetra chloride (CCl<sub>4</sub>) was administered intraperitoneally for 10 days from 11<sup>th</sup> day to day 20.

#### Preparation of extract

Roots of Cynodon dactylon were obtained from Herbal garden of G.S.V.M. Medical College, Kanpur. Roots of **Cynodon dactylon** were dried in shade then turn into coarse powder. About 500 gms of coarse powder of each was thus obtained which was then subjected to cold percolation method for 7 days using 70% alcohol as solvent. After 7 days extract of each drug was collected. The alcohol free extract was weighed and preserved in a refrigerator at 4°C.

Carbon tetra chloride was obtained from market. Since CCl<sub>4</sub> is a hepatotoxic agent, it induces hepatitis in the animals. Hepatitis produces anorexia and decreases in the body weight, Therefore assessment of weight loss was done in all the groups.

1. 60 grams of diet was given to each rabbit. Diet was provided between 11 A.M and 1.00 P.M daily. Weighed diet was given and the amount consumed was calculated from difference between the left over amount of diet 24 hours later. Water was given ad libitum.
2. Weight of the animals; Weight was recorded daily from first day to 20<sup>th</sup> day. Any increase or decrease in the weight of rabbit during drug administration was recorded

3. SGOT , SGPT,Alkaline phosphatase ,Serum bilirubin, Serum albumin estimation was done, blood samples were collected on zero day ,11<sup>th</sup> day,21<sup>st</sup> day.
4. Liver weight :At the end of the study rabbit were sacrificed and liver was taken out. It was weighed and preserved in 10% buffered formalin for histopathological study.

## PROCEDURE

Rabbits were divided into 2 groups with 6 rabbits in each group.

**GROUP I:** Animals of this group were treated with hepatotoxic agent ie carbon tetrachloride (CCl<sub>4</sub>) for 10 days in the dose of 0.05ml /Kg/day intraperitoneally from 11<sup>th</sup> day to day 20 along with normal feed. On 21<sup>th</sup> day blood samples were collected and rabbits were sacrificed.

**GROUP II:** Animals of this group were given extract of roots of Cynodon dactylon 100 mg/Kg/day orally for 20 days along with normal feed ,from 11<sup>th</sup> day onward carbon tetrachloride (CCl<sub>4</sub>) 0.05mg/kg, i.p. was also given followed by herbal drug .

Blood samples were collected on zero day before giving any drug to see the control value of liver function tests (L.F.T.),on 11<sup>th</sup> day to see the per se effect of herbal drug on L.F. T. and on 21<sup>th</sup> day to see the protective effect of herbal drug on L.F.T. The value obtained were compared.Blood samples were drawn from the marginal vein of pinna using 22 gauge needle, after the ear hairs were shaved off. 3 ml blood was collected in the vial, for the liver function test.

Body weight was measured daily . The animals of group I were sacrificed on 11<sup>th</sup>day and the animals of group II were sacrificed on 21<sup>th</sup>day. They were made unconscious, by giving ketamine.. The abdomen was exposed and liver was excised, weighed and was preserved in 10% buffered formalin for histopathological study.

## Assesment of Liver Injury

Assessment of liver injury was done by biochemical estimation and histopathological study of liver under light microscope.

## Biochemical Estimation

Serum bilirubin ,SGOT,SGPT, Alkaline phosphatase, serum albumin levels were estimated by Olympus autoanalyser in the department of Pathology.

## Histopathology

Histopathological study of the rabbit's liver was done to assess the extent of toxicity. Liver was taken out after sacrificing the rabbit. It

was weighed and preserved in 10% buffered formalin. Tissue sectioned to prepare slides. Staining was done with Hematoxylin and Eosin. Then slides were examined under light microscope and these slides were photographed<sup>3</sup>.

## Statistical Calculations

Mean, standard deviation and standard error of mean was calculated and results were analyzed by using paired t test and Student 't' test. P Value of < 0.05 was considered significant.

## OBSERVATIONS AND RESULTS

### Effect on diet intake, body weight, liver weight (Table no:1)

In Rabbits of **group I**, who were administered CCl<sub>4</sub> (0.05 mg/kg/day, intraperitoneally) along with normal feed, the diet intake was found to be **39.13 ±0.77 gm/day**. The decrease in food intake has lead to a decrease in body weight. The mean decrease in body weight in group I was considerably more than when compared to the group II, group III. The mean weight of liver was **28.38±0.18 gms**.

In Rabbits of **group II** (received Cynodon dactylon extract), average diet intake was decreased by 9.5%when compared to average diet intake during first 10 days of same group and increased by 26% when compared with group I ( CCl<sub>4</sub> administered group).Mean weight of the liver was measured to be **45.91±0.27**.

In **group I**,There was a highly significant (p<0.001), increase in the levels of serum transaminases , serum alkaline phosphatase ,serum Bilirubin and significant decrease in serum albumin with p<0.001 compared to zero day of same group (self control).

Administration of Cynodon dactylon extract and Silymarin to Rabbits ,feed on normal, diet did not alter the level of serum transaminase (p>0.10),serum alkaline phosphatase (p>0.10) serum bilirubin and serum albumin (p>0.10),when compared to zero day of same group.

In **group II**, The rise of serum transaminase(p<0.001,table 2,3), serum alkaline phosphatase(p<0.001,table 4), serum bilirubin (p<0.01,table 5)and decrease in serum albumin(p<0.001,table 6) due to hepatotoxic effect of CCl<sub>4</sub> when compared to zero day of same group .

Cynodon dactylon extract was able to bring down the level of serum transaminase( Table No.2,3),serum alkaline phosphatase (Table No.4),serum bilirubin (Table No.5) and increased in serum albumin (Table No.6) in a statistical highly significant amount (p<0.001), when compared with 11<sup>th</sup> day of rabbits receiving CCl<sub>4</sub> alone.

**Table no. 1: Showing average diet intake per day, Mean Liver weight (gm) of Rabbits, Mean body weight (Kg) of different groups.**

Group	Average Diet Intake (gm/kg) 1 <sup>st</sup> to 10 <sup>th</sup> day	Average Diet Intake (gm/kg) 11 <sup>th</sup> to 20 <sup>th</sup> day	Mean Liver Weight(gm)	Average Body weight (in kg )1 <sup>st</sup> to 10 <sup>th</sup> day	Average Body weight( in kg) 11 <sup>th</sup> to 20 <sup>th</sup> day
I (CCl <sub>4</sub> )	39.13 ± 0.77	-\$	28.38±0.18	1.04± 0.11	-\$
II (Cynodon dactylon extract)	60.11 ± 0.45	54.67 ± 0.47	45.91± 0.27	1.58±0.21	1.50±0.22

All values are in MEAN ± SE, \$ group I rabbits were sacrificed on 11<sup>th</sup> day.

**Table No 2: Showing Mean Aspartate transaminase(AST,SGOT) in IU/L of Rabbits in different groups**

GROUP	Mean SGOT ± SE (IU/L) At zero day	Mean SGOT ± SE (IU/L) At 11th day	Mean SGOT ± SE (IU/L) At 21th day
I (CCl <sub>4</sub> )	30.67 ± 0.88	87.17±1.58*	-\$
II (Cynodon dactylon extract)	29.17±1.35	29.83±1.51**	52.67±1.15*,#

✚ \*P-Value<0.001, \*\*P-Value>0.10, \$ group I rabbits were sacrificed on 11<sup>th</sup> day ; \*,\*\* values are compared with zero day of same group (self control) ; # P-Value<0.0001,# compared with 11th day of group I (CCl<sub>4</sub>)

Table No 3: Showing Mean Alanine transaminase(ALT,SGPT) in IU/L of Rabbits in different groups

Group	Mean SGPT $\pm$ SE (IU/L) At zero day	Mean SGPT $\pm$ SE (IU/L) At 11th day	Mean SGPT $\pm$ SE (IU/L) At 21th day
I(CCl <sub>4</sub> )	30.33 $\pm$ 1.49	132.67 $\pm$ 3.77*	-\$
II(Cynodon dactylon extract)	29.17 $\pm$ 1.14	30.67 $\pm$ 0.92**	64.50 $\pm$ 1.43*,#

\*P-Value<0.001, \*\*P-Value>0.10,\$ group I rabbits were sacrificed on 11<sup>th</sup> day ;\*,\*\* values are compared with zero day of same group (self control) ;# P-Value<0.0001,# compared with 11th day of group I (CCl<sub>4</sub>).

Table No 4: Showing Mean Alkaline phosphatase(ALP) in IU/L of Rabbits in different groups

Group	Mean ALP $\pm$ SE (IU/L) At zero day	Mean ALP $\pm$ SE (IU/L) At 11th day.,	Mean ALP $\pm$ SE (IU/L) At 21th day
I (CCl <sub>4</sub> )	36.50 $\pm$ 2.53	121.83 $\pm$ 3.07*	-\$
II (Cynodon dactylon extract)	40.17 $\pm$ 1.91	40.33 $\pm$ 1.73**	58.00 $\pm$ 0.89*,#

\*PValue<0.001,\*\*P-Value>0.10\$ group I rabbits were sacrificed on 11<sup>th</sup> day ;\*,\*\*values are compared with zero day of same group(self control); # P-Value<0.0001,# compared with 11th day of group I (CCl<sub>4</sub>)

Table No 5: Showing Mean Serum Bilirubin (mg/dl) of Rabbits in different groups.

Group	Mean Serum Bilirubin (mg/dl) $\pm$ SE At zero day	Mean Serum Bilirubin (mg/dl) $\pm$ SE At 11th day	Mean Serum Bilirubin (mg/dl) $\pm$ SE At 21th day
I (CCl <sub>4</sub> )	0.33 $\pm$ 0.04	1.07 $\pm$ 0.07*	- \$
II (Cynodon dactylon extract)	0.32 $\pm$ 0.04	0.37 $\pm$ 0.04**	0.50 $\pm$ 0.04***,#

\*P-Value<0.001, \*\*P-Value>0.10,\*\*\* P-Value <0.01 ; \$ group I rabbits were sacrificed on 11<sup>th</sup> day ;\*,\*\*,\*\* values are compared with zero day of same group(self control) ; # P-Value<0.0001,# compared with 11th day of group I (CCl<sub>4</sub>)

Table No 6: Showing Mean Serum Albumin (gm/dl) of Rabbits in different groups

Group	Mean Serum Albumin (gm/dl) $\pm$ SE At zero day	Mean Serum Albumin (gm/dl) $\pm$ SE At 11th day	Mean Serum Albumin (gm/dl) $\pm$ SE At 21th day
I (CCl <sub>4</sub> )	4.00 $\pm$ 0.10	2.33 $\pm$ 0.07*	-\$
II (Cynodon dactylon extract)	4.10 $\pm$ 0.14	4.15 $\pm$ 0.16**	3.25 $\pm$ 0.07***,#

\*P-Value<0.001, \*\*P-Value>0.10,\*\*\* P-Value <0.02 ; \$ group I rabbits were sacrificed on 11<sup>th</sup> day ;\*,\*\*,\*\* values are compared with zero day of same group(self control) ;# P-Value<0.0001,# compared with 11th day of group I (CCl<sub>4</sub>)

### Histopathological Assessment

#### Rabbits administered carbon tetrachloride

Grade III fatty changes and hydropic degeneration was present in 75% of rabbits and Grade II fatty changes was present in 25% of rabbits. Centrilobular (perivenular) and periportal inflammation was found in 75% and 25% of rabbits respectively, chiefly infiltrated with monocytes .Grade II inflammation was present in all rabbits.Grade II necrosis and loss of cord pattern was found in all rabbits (Figure 1).




Figure 1: A section of Rabbit Liver treated with CCl<sub>4</sub> alone showing marked fatty.

#### Rabbits administered Cynodon dactylon extract and CCl<sub>4</sub>

When rabbits on CCl<sub>4</sub> were compared with rabbits receiving Cynodon dactylon extract and CCl<sub>4</sub>, there was good protection of hepatic lobules from the damage induced by CCl<sub>4</sub>. Grade I fatty changes was present in 50% rabbits. Portal inflammation of grade I, infiltrated by monocytes were present in 30% of rabbits and necrosis of grade I was found in 20% of rabbits in centrilobular zone. Cord pattern was maintained (Figure 2).




Figure 2: A section of Rabbit Liver treated with Cynodon dactylon extract and CCl<sub>4</sub> showing sub maximum protection from damage induced by CCl<sub>4</sub>. (Grade I fatty changes in 50% and Grade I Portal inflammation in 30% area).

The result in this study suggest that administration of Cynodon dactylon extract to the rabbits received CCl<sub>4</sub> from 11<sup>th</sup> day to 20<sup>th</sup> day caused a decline in hepatotoxicity induced by CCl<sub>4</sub>. This is evidenced in the marked decrease in serum SGPT and SGOT level relative to the group treated with CCl<sub>4</sub> alone. Cynodon dactylon extract appears to exhibit protection against liver injury.

## DISCUSSION

Today, most xenobiotics to which humans are exposed come from sources that include environmental pollution, food additives, cosmetics products, agro-chemicals, processed food and drugs. In general, these chemicals in the absence of metabolism would not be eliminated from the body efficiently, and thus would accumulate in the body resulting in toxicity. Hepatic injury is a common sequel of exposure to toxic agents.

The CCl<sub>4</sub> is one of the most commonly used hepatotoxins in the experimental study of liver diseases<sup>4</sup>. Plant derived natural products such as flavonoids, terpenoids and steroids etc have received considerable attention in recent years due to their diverse pharmacological properties including hepatoprotective and antioxidant activity<sup>5,6</sup>. Realizing the fact this study was carried out to evaluate the hepatoprotective of Cynodon dactylon extract in this direction.

Our findings regarding weight of liver, however, are different from earlier reports. **Simmon's et al**, 1995 in their study have reported that increased organ weight (whether absolute or relative) is a sensitive indicator of organ toxicity<sup>7</sup>. Also, **Sodhi et al** in their study have reported an increase in the specific liver weight in experimental animals given protein restricted diet administered INH+R (causing hepatotoxicity)<sup>8</sup>.

In this study, CCl<sub>4</sub> was able to produce hepatic damage which is manifested by increase in serological marker and abnormal histopathology. These changes are similar to previous studies.

The serum level of marker enzymes: SGOT, SGPT and ALP reflect the physiological state of the liver. The levels of these enzymes change accordingly to the distortion of liver resulting from cellular injury of the organ caused by toxic metabolites and diseases. Serum and plasma enzymes levels have been used as a marker for monitoring chemically induced tissue damages<sup>9-10</sup>.

The toxicity of CCl<sub>4</sub> to the liver of mammal is largely as a result of the active metabolite, trichloromethyl radical<sup>11</sup>. The above radical bind to tissue macro- molecule and thus induce peroxidative degradation of membrane lipids of Endoplasmic Reticulum (ER), which are rich in polyunsaturated fatty acids. **Shenoy et al 2001**<sup>12</sup>, postulated, that such development would ultimately lead to the formation of lipid peroxides. The increase enzyme level in the plasma of CCl<sub>4</sub>- treated rabbits suggests that the toxicant was able to reach the liver and induce a detectable damage.

The increase in levels of serum bilirubin reflects the depth of jaundice and increase in transaminases and alkaline phosphatase indicates the cellular leakage and loss of functional integrity of cell membrane<sup>13</sup>. Liver enzymes are usually raised in acute hepatotoxicity, but tend to decrease with prolonged intoxication due to damage to liver cells<sup>14</sup>.

Rabbits treated with Cynodon dactylon extract have shown significant reduction of SGOT, SGPT, Alkaline phosphatase & serum bilirubin and histopathology provide direct evidence of functional protection of hepatocyte. Various pathological changes such as steatosis, centrilobular necrosis, loss of cord pattern observed in group I rabbits, were prevented in group II.

## CONCLUSION

From the discussion, it is clear that the carbon tetrachloride administration produces hepatic injury as is evident both by the changes in the biochemical parameters and histopathological changes in the present study. There is evidence of varying degree of oxidative stress leading to hepatocellular damage. We observed that roots of Cynodon dactylon protect the liver against CCl<sub>4</sub> induced hepatotoxicity.

Cynodon dactylon extract found to be effective in reducing SGOT, SGPT, ALP, S.bilirubin & cause increase in S.albumin level. Cynodon dactylon extract has shown protection in restoration of liver function and regeneration of liver cells as observed on histopathology.

This study was done on small scale and for short duration so further research needs to be done to confirm the above results and to find the active principle and mechanism of action responsible for their hepatoprotective activity.

## Acknowledgement

The authors sincerely thank to Department of Pharmacology and Pathology, G.S.V.M. Medical college, Kanpur for providing the necessary facilities for carrying out this work.

I am highly thankful to Dr. Priyamvada Sharma, M.D. Professor, Department of Pharmacology & Therapeutics S.N. Medical college, Agra, whose wealth of knowledge and vast experience has been instrumental at all stages of this study.

## REFERENCES

1. Ward F.M., Daly M.J. Liver Disease. In Clinical Pharmacy and Therapeutics, 3<sup>rd</sup> ed: Elsevier Science Limited; 2003:209.
2. Guillouzo A. Liver cell models in in-vitro toxicology. Environ Health Perspect 1998; 108(Suppl 2): 511-32.
3. Willianson EM, Okpako DT, Evans FJ. Selection, preparation and pharmacological evaluation of plant material. John Wiley, England; 1996.
4. Jhonson DE, Kroeming C. Mechanism of early CCl<sub>4</sub> toxicity in cultured rat hepatocytes. Pharmacol Toxicol 1998; 83: 231- 9.
5. De Feudis FV, Papadopoulos V, Driev K. Ginkgo biloba extract and cancer, a research area in its infancy. Fundam Clin Pharmacol 2003; 17: 405- 17.
6. Takeoka GR, Dao LT. Antioxidant Constituent of Almond. J Agric Food Chem 2003; 51: 496- 501.
7. Raghavendran BRH, Sathivel A, Devaki T. Hepatoprotective Nature of Seaweed alcoholic extract on Acetaminophen induced oxidative stress. J Health Sci 2003; 50: 42- 46.
8. Anandan R, Deepa Rekha R, Devaki T. Protective effect of Picrorrhiza kurroa on mitochondrial glutathione antioxidant system in D- galactosamine- induced hepatitis in rats. Curr Sci 1999; 76: 1543-5.
9. Lin JK, Wang CJ. Protection of crocin dyes in the acute hepatic damage induced by aflatoxin B and dimethylnitrosamine in rats. Carcinogenesis 1986; 7: 595- 99.
10. Ngaha EO, Akanji MA, Madusolumuo MA. Studies on correlation between chloroquine- induced tissue damage and serum enzyme changes in the rats. Experimentia 1989; 45: 143- 46.
11. Johnson DE, Kroening C. Ethanol feeding stimulates trichloromethyl radical formation from CCl<sub>4</sub> in cultured rat hepatocytes. Pharmacol Toxicol 1998; 83: 231- 39.
12. Shenoy KA, Somayaji SN, Bairy KL. Hepatoprotective effects of Ginkgo biloba against CCl<sub>4</sub> induced hepatic injury in rats. Indian J Pharmacol 2001; 33: 260-66.
13. Saraswat B, Visen PK, Patnaik GK, Dhawan BN. Anti-cholestatic effect of picroliv, active hepatoprotective principle of Picrorrhiza kurroa, against CCl<sub>4</sub> induced cholestasis. Indian J Exp Biol 1993; 31: 316- 18.
14. Cornelius CE. Biochemical evaluation of hepatic function in dogs. J Am Hosp Ass 1979; 15:25- 9.