

IN VITRO EVALUATION OF ANTI MYCOTIC ACTIVITY OF ETHANOLIC EXTRACT OF
*GLYCYRRHIZA GLABRA*R.V GEETHA^{1*}, ROY ANITHA²¹Faculty of Microbiology, ²Faculty of Pharmacology, Saveetha Dental College, Velappanchavady, Chennai 77 India.

Email: rgeetha2010@yahoo.in

Received: 9 May 2013, Revised and Accepted: 29 May 20123

ABSTRACT

Objective: The aim of the present study was to assess the anti fungal activity of ethanolic extract of *Glycyrrhiza glabra*. *Glycyrrhiza glabra* commonly called as liquorice or sweet root is one of the oldest known herbal remedies famous for its vast variety of therapeutic properties. Ethanolic extract of *Glycyrrhiza glabra* was tested for antifungal activity against *Candida albicans*, *Aspergillus niger*, *Aspergillus fumigates*, *Mucor spp* and *Penicillium marneffe*.

Method: Disc diffusion technique was followed for screening anti fungal activity. The discs were loaded with 100 μ l of ethanolic extracts at different concentrations [500 μ g/disc, 1000 μ g/disc and 2000 μ g/disc]. Positive controls used were fluconazole (10 mcg/disc) and amphotericin B (100 units/disc).

Result : After incubation at 28 $^{\circ}$ C for 48 hours, the zone of Inhibition was measured.

Conclusion: The extract at different concentrations showed varying degree of antifungal activity against the micro organisms tested compared to standard.

Keywords: *Glycyrrhiza glabra*, disc diffusion, zone of inhibition, anti mycotic activity.

INTRODUCTION

The very old folk medicine is based on the use of plant and plant extracts. There are innumerable types of indigenous plants that have been used by people for centuries in the treatment of many ailments. The use of herbs to treat disease is almost universal and is often more affordable than expensive modern pharmaceuticals.[1] Many of the phytochemicals found in the herbs have beneficial effects and can be used to treat human diseases. It has become increasingly more in recent years as scientific evidence about the effectiveness of herbal medicine has become more widely available.[2] There are herbal remedies that have been tested for their use as a treatment for many fungal diseases.

Glycyrrhiza glabra, also known as liquorice and sweet wood, is native to the Mediterranean and certain areas of Asia. It is a perennial herb which possesses sweet taste. The main taproot, which is harvested for medicinal use, is soft, fibrous, and has a bright yellow interior. It was one of the most widely known medicines in ancient history, and records of its use include Assyrian tablets of around 2000 BC and cortisone has been found useful for arthritis and allergies.[3] In addition licorice has been used for mild Addison's disease and other adrenal insufficiencies, such as hypoglycemia. [4] Licorice also acts like the hormone, ACTH, causing sodium retention, potassium depletion, and water retention. The herb contains glycyrrhizin, glycyrrhetic acid, flavonoids, asparagine, iso-flavonoids, and chalcones. The glycoside, glycyrrhizin has a similar structure and activity as the adrenal steroids.[5,6] It also possess good anti bacterial, [7] anti fungal,[8] anti oxidant,[9] antitussive,[10] hepatoprotective [11] and anti inflammatory activity.[12] Historically, the dried rhizome and root of this plant were employed medicinally as an expectorant and carminative. It is used for treating upper respiratory ailments including coughs, hoarseness, sore throat and bronchitis.

TEST MICROORGANISMS

Fungal strains used were *Candida albicans*, *Aspergillus fumigates*, *Aspergillus niger*, *Mucor sps*, and *Penicillium marneffe*. The organisms were obtained from Department of Microbiology, Saveetha Medical College and maintained in SDA slope at 4 $^{\circ}$ C.

METHODOLOGY

The extracts were prepared in the following concentrations in sterile water. 5mg/ml and 10mg/ml and 20mg/ml. 100 μ l of extract of different concentrations were loaded on sterile filter paper discs measuring 6mm in diameter, so that the concentration of the extract on each disc was 500 μ g, 1000 μ g and 2000 μ g respectively. The discs were dried and kept aseptically.

SCREENING OF ANTIFUNGAL ACTIVITY [DIS DIFFUSION TECHNIQUE]

The ethanolic extract of *Glycyrrhiza glabra* was screened for antifungal activity by disc diffusion method. Activated cultures of fungal strains in Sabouraud's broth were adjusted to 0.5 McFarland standards [108 cfu/ml].[13-17] 100 μ l of the inoculum was introduced to molten Sabouraud's dextrose agar and poured in the sterile petri plates and allowed to set. Sterile filter paper discs (6.0 mm diameter) impregnated with 2000 μ g/disc, 1000 μ g /disc and 500 μ g /disc of the plant extract dissolved in sterile water were placed on fungal seeded plates and incubated at 28 $^{\circ}$ C for 48 hrs. As a positive control, Fluconazole (10 mcg /disc) and Amphotericin B (100 units /disc) were used. Following an incubation period of 48 hrs, plates were removed from the

incubator and antifungal activity was evaluated by measuring zones of inhibition of fungal growth. Clear zones within which fungal growth was absent were measured and recorded as the diameter (mm) of complete growth inhibition. The whole experiment was performed three times to minimize error.

RESULTS AND DISCUSSION

Effect of three different concentrations (2000, 1000, 500, μ g /disc) of the ethanolic extract of *Glycyrrhiza glabra* was tested against the fungal strains using disc diffusion technique. All the concentrations of the test solution inhibited the fungal species with varying degree of sensitivity. The antifungal activity of the extract against the fungal strains is shown in Table 1. The extract showed good antifungal activity against the two *Aspergillus Spp*, *Aspergillus niger* and

Aspergillus fumigates at all concentrations with a maximum zone of inhibition of 21mm and 20mm diameter respectively at 2000 µg

/disc concentration. The lowest concentration of the extract showed least activity against *Mucor sps* and *Penicillium morneffeji*, while the higher concentrations showed antifungal activity with a maximum zone diameter of 18mm and 17mm respectively at 2000 µg/disc concentration. With *Candida albicans*, the lowest concentration of the extract showed a zone of inhibition of 9mm while the highest concentration showed an inhibitory zone of 20mm. From the results, it was evident that the lower concentration showed very weak while the higher concentration of the extract showed good antifungal activity against all the fungal strains tested.

Table 1: Antifungal activity of ethanolic extract of the *Glycyrrhiza glabra*

Extract	Conc [µg]	Zone of inhibition [in mm diameter]				
		B1	B2	B3	B4	B5
Ethanolic	500	09	10	12	07	08
	1000	15	20	21	15	13
	2000	20	21	20	18	17
Fluconazole	10mcg /disc	24	21	22	22	23
Amphotericin B	100mcg /disc	25	23	20	24	25

B1-*Candida albicans*, B2-*Aspergillus fumigates*, B3-*Aspergillus niger*, B4-*Mucor sps*, B5-*Penicillium marneffeji*.

Over the past two decades fungal infections have evolved into important cause of morbidity and mortality in modern medicine. The prevalence of resistance to antifungal agents has significantly increased. So it makes necessary to discover new classes of antifungal compounds to treat fungal infections. The research on natural products derived compounds has accelerated in recent years due to their importance in drug discovery.[18] Plants are rich source of bioactive secondary metabolites of wide variety such as tannins, terpenoids, alkaloids, and flavonoids, which are reported to have in vitro antifungal properties. A series of molecules with antifungal activity against different strains of fungus have been found in plants, which are of great importance. The results obtained from our study shows that ethanolic extract of *Glycyrrhiza glabra* has got a very good anti mycotic activity against the selected fungal species

CONCLUSION

The results of present investigation clearly indicate the antifungal activity of ethanolic extract of the *Glycyrrhiza glabra*. The anti-fungal activities could be enhanced if active components are purified and adequate dosage determined for proper administration. Further studies on their cytotoxicity or toxicity will be beneficial in providing data on the possible harmful effects of this extract commonly used by the numerous local communities. Thus, the study ascertains the value of plants used in Ayurveda, which could be of considerable interest to the development of new drugs.

ACKNOWLEDGEMENTS

The authors are thankful to Mr.R.Rajendran, GreenChem. Herbal Extract & Formulations. Bangalore,for providing us with the Ethanolic extract of the *Glycyrrhiza glabra* as a gift sample for our research work.

REFERENCES

1. Sheetal verma and S.P.Singh., Current and future status of herbal medicines. Veterinary world, Vol.1 (11): 347 – 350.

2. Si-Yuan Pan, Si-Bao Chen, Hong-Guang Dong, ZhiLing Yu, Ji-Cui Dong, Zhi-Xian Long, Wang-Fun Fong, et al., New perspectives on Chinese herbal medicine research & development. Evidence Based Complementary and alternative medicine 2011.vol 2011,11 pages.
3. Olukoga A, Donaldson D. The history of liquorice: the plant, its extract, cultivation, and commercialisation and etymology. J R Soc Health 1998; vol 118:300-304
4. A.AAl-Qarawi, H.A.Abdel Rahman, B.H.Ali, Liquorice (*Glycyrrhiza glabra* and the adrenal kidney -pituitary axis in rats. Food and chemical toxicology, vol 40 [10] : 1525-1527
5. Fan Y. G.; Shi Z. Q.; He B. L.; Extraction separation and application for glycyrrhizic and glycyrrhetic acid. Natural product Research and Development. 1996, vol 8 : 93-99
6. Yang L.; Liu Y. L.; Lin S. Q.; The determination of flavonoid in 6 kinds of licorice root by HPLC. Acta Pharmaceutica Sinica. 1990; 25 : 840-848
7. Manoj M. Nitalikar, Kailas C. Munde, Balaji V. Dhore, Sajid N. Shikalgar Antibacterial Activities of *Glycyrrhiza glabra* Root Extract ,International Journal of PharmTech Research Vol.2, [1] :899-901
8. Tharkar P.R.,Tatiya A.U., Shinde P.R. , Surana S.J., Patil U.K. ,Antifungal Activity of *Glycyrrhiza glabra* Linn. And *Embllica Officinalis* Gaertn. by Direct Bioautography Method International Journal of PharmTech Research. ,2010; 2 [2] :1547-1549,
9. Morteza-Semnani K, Saeedi M, Shahnavaz B,Comparison of antioxidant activity of extract from roots of licorice (*Glycyrrhiza glabra* L.) to commercial antioxidants in 2% hydroquinone cream. J Cosmet Sci. 2003 Nov-Dec; 54 [6] :551-558.
10. Anderson DM, Smith WG. The antitussive activity of glycyrrhetic acid and its derivatives. J.Pharm.Pharmacol. Jul 1961;vol 13: 396-404.
11. Kang HE, Lee MG, Hwang SJ, Kim SC, Lee CH, et al. Liquiritigenin, a flavonoid aglycone from licorice, has a choleric effect and the ability to induce hepatic transporters and phase-II enzymes. Am J Physiol Gastrointest Liver Physiol. Feb 2009; 296[2]:372-81
12. Okimasu E, Moromizato Y, Watanabe S, et al. Inhibition of phospholipase A2 and platelet aggregation by glycyrrhizin,an anti inflammation drug. Acta Med Okayama1983;37[5] :385-391.
13. A.Manja malai, Sneha susan Varghese, Aishwarya Haridas and V.M.Belin grace., Antifungal,anti inflammatory and GC-MS analysis for Bioactive molecules of *Tridax procumbens* leaf, Asain J of pharmaceutical and clinical research , 2012; 5 [suppl 1]:139 -145
14. Collins, CH and Lyne, P.M .Microbiological methods, London, Butterworths and co; 1976
15. Betty A.Forbes., Daniel F.Sahm., Alice S.Weissfeld. Bailey & Scott's Diagnostic Microbiology 11th edition Mosby;1998
16. Connie R.Mahon., George Manuselis., Saunder's Diagnostic Microbiology 2nd Edition. W.B.Saunder's company;2000
17. J. G. Colle., A. G. Faster., B. P. Marmion. A. Simmons. Practical Microbiology [Mackie & Mc Cartney] 14th edition. Elsevier ;2006
18. 18.N.Jayashree, N.Narayanan, Lata sriram., Antibacterial antifungal and anti mycobacterial studies on some synthetic Dimethoxy flavones, Asain J of pharmaceutical and clinical research , 2012; 5 [suppl 1]:101 -103