



ANTI-NOCICEPTIVE ACTIVITY OF METHANOLIC EXTRACT OF *OCIMUM GRATISSIMUM* (LABIATE) ON EXPERIMENTAL ANIMALS

VIKRAM KUMAR^{1*}, MANJEET SINHA², DR.ANGSHU BANERJEE³, DR.JYOTI MOHANTY⁴

¹ Lecturer, Sri Balaji College of Pharmacy- Macheda, Jaipur, Rajasthan, India, 302013, ²Principal, Mahatma Gandhi College of Pharmacy- Jaipur, ³ Director, Sri Balaji College of Pharmacy- Jaipur, ⁴ Professor, Kanak Manjari Institute of Pharmaceutical Sciences-Rourkela, Orissa, India. Email: vikramyadav05@rediffmail.com

Received: 25 Jan 2011, Revised and Accepted: 27 Feb 2011

ABSTRACT

The methanolic leaves extract of *Ocimum gratissimum* was investigated for anti-nociceptive effects in mice. The models used to study the effect on nociception are the acetic acid induced abdominal constriction test, hot- plate method in mice. The extract demonstrated a significant ($P < 0.05$) anti-nociceptive activity at all the doses (50,100 and 200 mg/kg body weight i.p.) tested group and standard group (piroxicam 20mg/kg) compared to control normal saline. The activity resides more at the highest dose 200mg/kg body weight i.p. that was found to have the highest percentage (92.5 %) of inhibition of the abdominal constriction induced by acetic acid in mice. Also the extract was found to have a significant ($p < 0.05$) inhibitory effect on hot-plate method at all the three doses. The extract caused a significant ($p < 0.05$), dose dependent inhibition of acetic acid-induced writhing and hot-plate method. The intraperitoneal LD50 value of the extract was 1285.5mg/kg body weight in mice. Preliminary phytochemical screening revealed the presence of alkaloids, saponins, tannins and flavonoids. The results suggest the extract contained pharmacologically active principles, and supports the local application of the plant in painful conditions. Further studies may reveal the exact mechanisms of action responsible for the analgesic activities of *O. gratissimum* leaves extract.

Keywords: *Ocimum gratissimum*, Labiatae, Anti-nociceptive, Piroxicam, Acetic acid-induced writhing, hot-plate.

INTRODUCTION

Ocimum gratissimum is commonly known as fever leaf in general but is has different native names in different part of the country. In Yoruba language, it is known as Ewfirm ajase, Nchu-nwu in Ibo, Bunsuru daji in Hausa, Ireru in Ebira, Ebaubokho in Benin, ufuo-yibo in Urhobo and ntion in Efik¹. There are about 60 or more species of *Ocimum* and numerous varieties, belonging to the Family Labiatae. This different types of species are represented by the five most important representatives of the more than 60 *Ocimum* species and these include (i) *Ocimum gratissimum*, (ii) *Ocimum basilicum*, (iii) *Ocimum americanum*, (iv) *Ocimum sanctum* and (v) *Ocimum americanum*^{2,3}.

Ocimum gratissimum Linn. (Labiaceae) is a herbaceous plant commonly found in tropical Asia especially India. It is used in the treatment of epilepsy in the coastal area of Nigeria⁴, High fever⁵, and Diarrhea^{5,6}. The plant is also used to treat typhoid fever and diabetes⁷⁻⁹. Today, basil is used mainly as a culinary herb. Its medicinal value is not as widely appreciated in Western World. In France it is used in perfumes and cosmetics¹⁰.

This research was aimed at investigating the possible anti-nociceptivities of methanolic leaves extract of the plant in order to support or refute the claims by traditional herbalists.

MATERIALS AND METHODS

Preparation of extract

The fresh leaves of *Ocimum gratissimum* was chopped, cleaned and air dried for minimum one week. After that the size was reduced with a mortar and pestle into a fine powder. 100 g of the powder was extracted with 90% methanol (2.5 litres) using Soxhlet apparatus process for 50-55 h. The liquid extract was then concentrated on a water bath to give a brownish solid extract with a mean yield of 10% w/w.

Methanolic extract obtained was washed with hexane to obtain purified methanolic extract. Methanolic fraction was successively extracted with ethanol (90 % w/w) in Soxhlet apparatus for 50-55 h.¹¹

Acute toxicity study

This was conducted by using the method described by Lorke. In the initial phase, mice were divided into 3 groups of three and treated with the methanolic leaves extract of the plant at doses of 10, 100

and 1000mg extract/ kg body weight intra peritoneal (i.p.) and were then observed for 24 hrs for signs of toxicity including death. In the final phase, mice were divided into 4 groups of one mouse each and treated with the ethanol extract at doses of 600, 1000, 1600 and 2900 mg / kg body weight i.p. The median lethal dose (LD50) was calculated from the second phase¹².

Acetic acid induced abdominal constriction

Swiss mice (20-25gm) were divided into 5 groups of 6 mice each. The first group was given 10 ml/kg of Normal saline i.p. and served as control or placebo, groups 2 received Piroxicam 20mg/kg as a positive control or standard, 3, 4 and 5 received 50, 100 and 200 mg of extract per kg of body weight i.p. respectively. Thirty mins later, mice in all the groups were treated with Acetic acid (1%v/v, 10ml/kg body weight i.p.). Five minutes after Acetic acid injection, mice were placed in individual cages and the number of abdominal contractions was counted for each mouse for a period of 10 mins¹³.

Hot-plate assay

Swiss mice (20-25gm) were divided into 5 groups of 6 mice each. The first group was given 10 ml/kg of Normal saline i.p. and served as control or placebo, groups 2 received Piroxicam 20mg/kg as a positive control or standard, 3, 4 and 5 received 50, 100 and 200 mg of extract per kg of body weight i.p. respectively. A 500 ml of glass beaker was placed on hot-plate. The temperature of the hot-plate was then regulated to 50 ± 20 C in order to obtain the animal's response to electric heat-induced nociceptive pain stimulus (licking of the forepaws and eventually jumping out of the glass beaker). Jumping out of the beaker was taken as an indicator of the animal's response to heat-induced nociceptive pain stimulus. The time taken for each mouse to jump out of the beaker (i.e. reaction time) recorded in seconds. Readings were taken at intervals of 30, 60 and 90 mins after extract administration. Cut off time in the absence of response was 60 s to avoid tissue damage to the mice paws¹⁴.

Statistical analysis

Results were expressed as mean \pm Standard Error of Mean (SEM). The data was statistically analyzed using the one-way ANOVA to determine whether results in a particular group were significantly different from those in the corresponding control groups. Results were statistically significant when P values are less than 0.05 ($P < 0.05$) as described by Duncan¹⁵.

RESULTS

The freshly prepared extracts were subjected to preliminary phytochemical screening test for various constituents. This revealed the presence of alkaloids, tannins, saponins, flavonoids, terpenoids and steroids. The sign of toxicity was first noticed after 4-6 hrs of extract administration.

There was decreased loco-motor activity and decreased sensitivity to touch and jerking. Also there was decreased feed intake, and prostration after 10 hrs of extract administration. The median lethal

dose (LD50) in mice was calculated to be 1285.5 mg/kg body weight.

The extract demonstrated a significant ($P < 0.05$) anti-nociceptive activity at all the doses (50,100 and 200 mg/kg body weight i.p.) tested compared to control normal saline. The activity resides more at the highest dose 200mg/kg body weight i.p. that was found to have the highest percentage (92.5 %) of inhibition of the abdominal constriction induced by acetic acid in mice (Table 1). Also the extract was found to have a significant ($p < 0.05$) inhibitory effect on hot-plate method at all the three doses as shown in Table 2.

Table 1: Effect of methanolic extract of *O. gratissimum* on writhing assay

S.No	Treatment	Mean wriths
1	Vehicle (Placebo/control)	22.5
2	Piroxicam (Standard)	5.65*
3	Ocimum gratissimum 50ml/kg	7.54*
4	Ocimum gratissimum 100ml/kg	7.15*
5	Ocimum gratissimum 200ml/kg	2.64*

* $P < 0.05$ compared to vehicle (control).

Table 2: Effect of methanolic extract of *O. gratissimum* on Hot-plate assay

S.No	Treatment	Reaction latency (30 mins)	Reaction latency (60 mins)	Reaction latency (120 mins)
1	Vehicle (Placebo/control)	8.2	11.4	9.5
2	Piroxicam (Standard)	20.4	49.8*	47.4*
3	Ocimum gratissimum 50ml/kg	27.6*	33.8*	37.5*
4	Ocimum gratissimum 100ml/kg	23.3*	36.4*	37.2*
5	Ocimum gratissimum 200ml/kg	44.2*	44.4*	46.8*

* $P < 0.05$ compared to vehicle(control).




Fig. 1: Effect of methanolic extract of *O. gratissimum* on writhing assay, * $P < 0.05$ compared to vehicle.




Fig. 2: Effect of methanolic extract of *O. gratissimum* on hot-plate assay, * $P < 0.05$ compared to vehicle.

DISCUSSION AND CONCLUSION

The extract produce significant (* P<0.05) analgesic (anti-nociceptive) activity at all the doses for tested group and standard group compared to control group of animal.

Acute pain as writhing assay is chemical assay that may produce some inflammation and used as model of visceral pain. Possible mediators involved in the acetic acid induced inflammatory pain are not well known. It has been reported that bradykinin, neurokinins and prostanoids involves in the sensory C-fibres activation after intra-peritoneal injection of acetic acid. The activity demonstrated by the extract might be due to the presence of flavonoids and phenolics that were present in the extract¹⁶⁻¹⁸.

The Hot-plate assay is somatic pain model that doesn't produce inflammation. The repetitive afferent input enhances the response of the animal to a given noxious stimulus (radiant heat on Hot-plate). Activation of prostanoid receptors increases the opening of voltage sensitive Ca²⁺ channels and enhances primary afferent peptide release. It may be possible that extract exert its therapeutic action by antagonizing prostanoid receptor or inhibiting the synthesis of prostaglandins^{18,19}.

In conclusion, these preliminary investigations and data obtained from this study demonstrated that effect of herbal plant *Ocimum gratissimum* aqueous leaves extract have good analgesic or anti-nociceptive activity.

ACKNOWLEDGEMENT

I take this privilege and pleasure to acknowledge the contributions of many individuals including my wife Mrs Priyanka Yadav, our college director Dr. Angshu Banerjee who have been inspirational and supportive throughout my work undertaken and endowed with the most precious knowledge to see success in my endeavor.

REFERENCES

- Sofowora, L.A (1993): Medicinal plants and traditional Medicinal Medicine in African. Spectrum books Ltd, Ibadan. Pp. 55-71.
- Mandal, J and Pattnaik E.D. (2000) .Alginate encapsulation of auxiliary buds of *Ocimum Americanum*, *Basilicum*, *Gratissimum* and *Sanctum*. J. cellular developl. Biol., **36 (4)**:287-292.
- Martin, A.P, and Salguerio ,G.O .(1999). Composition of essential oil of *Ocimum Canum Ocimum Gratissimum*, *Ocimum Basilicum* and *Ocimum Minimum*. *Planta medica*, **65 (2)**:187-189.
- Osito, N.G. (1992): A system of traditional health care. Vol .2. pp. 56-58.
- Oliver, B. (1980): Medicinal plants in Nigeria. Published by Nigerian college of Arts, Science and Technology. Ibadan, pp.90-94.
- Sofowora, L.A (1993): Medicinal plants and traditional Medicinal Medicine in African. Spectrum books Ltd, Ibadan. Pp. 55-71.
- Adjanahoun, E., Aliyu, M.R.A., Ake-Assi, L., Elewude, J.A., Dramane, K., Fadoju, S.O., Gbile, Z.O., Goudole, E., Johnson, C.L.A., Keita, A., Morakinyo, O., Ojewole, J.A.O., Olatunji, A.O and Sofowora, E.A (1991), Traditional medicine pharmacopoeia contribution to ethanobotanical floristic studies in Western Nigeria, Pub. Organisation of African Unity, Scientific Technical and Research Commission Lagos, Nigeria. 420 p.
- Igoli, J.O., Tor- Anyim, T.A, Usman, S.S., Oluma, H.O.A and Igoli, N.P. (2002) Folk medicine of the lower Benue Valley of Nigeria .In: Recent progress in medicinal plants vol 7. Ethnomedicine and pharmacognosy. 11 (Eds. V.K Singh, J.N Govil, S, Hashmi and G.Singh), Sci. Tech. Pub. U.S.A. 327-338.
- Tor-Anyin, T.A, Shaatu, R, and Oluma, H.O.A. (2003) Ethnobotanical Survey of Antimalarial medicinal plant among the Tiv people of Nigeria. *J. Herbs, Spices and medicinal plants* **10 (3)**: 61-74.
- Ross, Ivan A. (2003): Medicinal plants of the World: Chemical constituents, traditional and modern medicinal uses. 2nd ed. Totowa, NJ, Humana Press, pp 205.
- Trease, G.E., Evans, M.C. (1989): Text book of Pharmacognosy 13th Edition Bailliere Tindall, London, Toronto, Tokyo. Pgs. 200-201, 340-348, 419-423, 626-630, 765-775.
- Lorke, D. (1983): A New Approach to Practical Acute Toxicity Testing. *Arch. Toxicol*, **53**: 275 - 287.
- Koster, R., Anderson, M. and Debeer, E.J. (1959). Acetic acid analgesic screening. *Fed. Proc.* **18**:412-417.
- Sharma, K.K., Khanna, T Sen, P. (1982). Current status of centrally acting peptides. In: (Dhawan, BN, ed.) *Advances in Biosciences* Vol 38, Oxford: Pergamon Press, p.147.
- Duncan, R. C; Knapp, R. G and Miller, M. C. (1977): Test of Hypotheses in population Means, In: *Introductory Biostatistics for Health Sciences*, John Wiley and Son, Inc. N. Y. pp. 71 - 76.
- Bentley, G.A., Newtons, S.H. and Starr, J. (1983). Studies on the antinociceptive action of agonist drugs and their interaction with opioid mechanisms. *Bri. J. Pharmacol.* **79**:125-134.
- Besra SE, Sharma RM, Gomes A (1996). Anti-inflammatory effect of petroleum ether extract of leaves of *Litchi Chinensis* Gaertn (Sapinadaceae). *J. Ethnopharmacol.* **54**: 1-6.
- Derardt, R., Jougney , S., Delevalce, F., Falhourt, M. (1980). Release of prostaglandins E and F in an allogenic reaction and its inhibition. *Eur. J. Pharmacol.* **51**: 17 -24.
- Ahmadiani, A., Fereidoni, M., Semnianian, S., Kamalinejad, M., Saremi, S. (1998). Anti-nociceptive and anti-inflammatory effects of *Sambucus ebulus rhizome* extract in rats. *J. Ethnopharmacol.* **61**:229-235.