

ANTIOXIDANT ACTIVITY, TOTAL PHENOLIC AND FLAVONOID CONTENTS OF *ARTOCARPUS HETEROPHYLLUS* AND *MANILKARA ZAPOTA* SEEDS AND ITS REDUCTION POTENTIAL

K.SHANMUGAPRIYA^{1*}, P.S.SARAVANA², HARSHA PAYAL², S. PEER MOHAMMED², WILLIAMS BINNIE²

¹Assistant Professor, ²School of Biotechnology, Dr. G.R. Damodaran College of Science, Coimbatore 641014, Tamil Nadu, India.
Email: shanmugapriya8@gmail.com

Received: 11 July 2011, Revised and Accepted: 24 Oct 2011

ABSTRACT

The four different extracts of *Artocarpus heterophyllus* and *Manilkara zapota* seeds were undertaken and studied for their total phenolics and flavonoids contents, reducing power and antioxidant activity. Total phenolic contents were determined spectrometrically by using Folin-Ciocalteu method and were calculated as Gallic acid equivalents. Flavonoids and reducing power were also determined spectrometrically using standards methods. The antioxidant capacity was studied using DPPH radical scavenging assay and ABTS radical scavenging assay methods. Plants containing phenols and flavonoids have been reported to possess strong antioxidant properties. The results of antioxidant activity showed high effect in both extracts. From the present study, it can be concluded that *Artocarpus heterophyllus* and *Manilkara zapota* seed extract has effective antioxidant activity due to the presence of high amounts of phenolic compounds. The results revealed that the extracts showed high flavonoids content and reducing potential. Both extracts may therefore be a good functional medicine as well as pharmaceuticals plant-based products.

Keywords: Antioxidant, Total Phenolic, Flavonoids, Reducing potential, DPPH, ABTS

INTRODUCTION

Nature has been a source of medicinal agents for thousands of years and an impressive number of modern drugs have been isolated from natural sources, many based on their use in traditional medicine. Free radicals are the main culprit in lipid peroxidation, highly reactive free radicals and oxygen species are present in biological systems from a wide variety of sources¹. Free radicals are fundamental to many biochemical processes and represent an essential part of aerobic life and metabolism^{2,4}. Free radical oxidative stress caused a wide variety of clinical disorders³. A serious imbalance between the production of free radicals and the antioxidant defense system is responsible for oxidative stress⁵. Antioxidants exert their mode of action by suppressing the formation of reactive oxygen species either by inhibition of enzymes or by chelating trace elements⁶.

Current research is now directed towards finding naturally occurring antioxidants of plant origin. Antioxidants have been reported to prevent oxidative damage by free radical and ROS; any may prevent the occurrence of disease, cancer and aging. It can interfere with the oxidation process by reacting with free radicals, chelating, catalytic metals, and also by acting as oxygen scavengers⁷.⁸. Natural antioxidants, particularly in fruits, vegetables and beverages have gained interest among consumers⁹. Antioxidants are dietary substance that protects body cells from the oxidative damage to a target molecules caused by oxidation from free radicals by reactive oxygen species (ROS)¹⁰.

During the last two decades there has been in search for new plant derived drugs containing the medically useful alkaloids, glycosides, polyphenolics, steroids and terpenoids derivatives, which contributes to the antioxidant property¹¹. Dietary phenolic compounds and flavonoids have generally been considered, as non-nutrients and their possible beneficial effect on human health have only recently been recognized. Flavonoids are known to possess antioxidant and anticarcinogenic activities¹². Therefore search for natural antioxidants of plant origin gained momentum in recent years.

The *Artocarpus heterophyllus* (Jackfruit) is a species of tree of the mulberry family Moraceae. It is native to Western Ghats of India, Malaysia and also found in central and eastern Africa, south-eastern Asia, Florida, Brazil, Australia and many Pacific Islands¹³. *Artocarpus heterophyllus* possesses known anti-bacterial, anti-fungal, anti-diabetic, anti-inflammatory, antioxidant and anti-helminthic activities¹⁴.

Manilkara zapota L. (Sapodilla) belongs to the family Sapotaceae. It is an evergreen, glabrous tree, 8-15 m in height. It is cultivated

throughout India, though it is native to Mexico and Central America. The seeds are aperients, diuretic tonic and febrifuge. Bark is antibiotic, astringent and febrifuge.

The main objectives of the study are to analysis the total phenolic and flavonoid content, reducing power, antioxidant activity.

METHODS AND MATERIALS

Chemicals used

All chemicals were purchased from Sigma Chemical, Merck Chemical Supplies, SD fine and Himedia. All other chemicals used were obtained commercially and were of analytical grade.

Collection of plant material

Artocarpus heterophyllus (Jackfruit) and *Manilkara zapota* (Sapodilla) were collected from Pallamudir, Coimbatore, Tamil Nadu, India.

Preparation of Extracts

The seeds were shade dried for one month and coarsely powdered. Extraction of the active ingredient from the seed powder was carried out using specific method¹⁵. 25g of the powdered seeds were extracted by soxhlet apparatus using 250ml of four different solvent (ethanol, acetone, ethylacetate and water) in a separate flask. The extraction lasted for six hours. The extract obtained was concentrated by evaporation using water bath at 100°C and stored at 4°C in cold room.

Determination of Total Phenolic content

Total phenolic content was determined by Folin-Ciocalteu reagent¹⁶. 10ml of the samples were taken in test tubes and made up to the volume of 1ml with distilled water. Then 0.5ml of Folin-Ciocalteu phenol reagent (1:1 with water) and 2.5ml of sodium carbonate solution (20%) were added sequentially in each tube in triplicate manner. Soon after vortexing the reaction mixture, the test tubes were placed in dark for 40 minutes and the absorbance was recorded at 725nm against a reagent blank. All tests were carried out in triplicate. Total content of phenolic compounds in extracts was expressed as Gallic Acid Equivalent (GAE) in milligram per gram extract.

Determination of Total Flavonoid content

Flavonoid content was determined according to the aluminum chloride colorimetric method with some modifications¹⁷. The sample solution (0.5ml) was mixed with 1.5ml of 95% ethanol, 0.1ml of 10% aluminum chloride hexahydrate, 0.1ml of 1M potassium acetate, and

2.8ml of distilled water. After incubation at room temperature for 40 minutes the absorbance of the reaction mixture was measured at 415nm. The same amount (0.1ml) of distilled water substituted for the amount of 10% aluminum chloride as the blank. All tests were performed in independent triplicates. The total flavonoid content was expressed in milligram per gram extract of Quercetin Equivalent (QE).

Determination of Reduction potential

The reducing antioxidant power was determined by specific method¹⁸. 1.0ml of the different concentration of various extracts of the sample was mixed with 2.5ml potassium ferric cyanide and 2.5ml phosphate buffer (pH 6.6). The mixture was incubated at 50°C for 20 minutes. After the incubation, 2.5ml of Trichloroacetic acid (10%) was added to it and centrifuged at 3000rpm for 10 minutes. 2.5ml of the supernatant was taken and mixed with 2.5ml distilled water and 0.5ml of ferric chloride (0.1%) were added to it. The absorbance of the colour was measured spectrophotometrically at 700nm. Increased absorbance of the reaction mixture indicates increase in reducing power. All tests were carried out in triplicate.

Determination of antioxidant Activity

DPPH (2, 2-diphenyl, 1-picryl hydrazyl) radical scavenging assay

The antioxidant effect of extract on DPPH radical was assayed using specific method¹⁹. 3.0ml of 1.0mM DPPH in ethanolic solution was added to 2.5µl and 5µl of the plant extract. DPPH solution with ethanol was used as positive control and ethanol alone acted as blank. After 30 minutes, the discoloration from deep violet to yellow colour was measured at 518nm in a spectrophotometer.

The percentage inhibition was calculated by the following formula:

$$\text{Scavenging activity (\% of inhibition)} = \frac{\text{Ac} - \text{Ae}}{\text{Ac}} \times 100$$

Where,

Ac – Absorbance of DPPH radical as control.

Ae – Absorbance of DPPH radical in the presence of plant extract.

ABTS (2, 2'-azinobis (3-ethylbenzthiazoline)-6-sulphonic acid) radical scavenging assay

The effect of extract on ABTS radical was assayed using specific method²⁰. ABTS⁺ radical was prepared by adding 2.45mM ammonium per sulphate with 7mM ABTS solution and kept in dark for 12-16 hours at room temperature. 0.5ml of various plant extracts were added to 0.3ml of ABTS solution and the final volume was made up to 1.0ml with ethanol. After 6 minutes, the absorbance was read at 745nm and the percentage inhibition scavenging activity was calculated using the scavenging activity formula as presented earlier.

RESULTS AND DISCUSSION

Plants containing phenols and flavonoids have been reported to possess strong antioxidant properties.

Determination of Total Phenolic content

Phenolic compounds may contribute directly to antioxidative action. The four different solvents at 100 mg/g of seed extract of *Artocarpus heterophyllus* contains total phenolic content was found to be 4.16±0.01mg/g (ethanolic), 2.30±0.02mg/g (acetone), 2.77±0.02mg/g (ethyl acetate) and 1.18±0.01mg/g (aqueous) of Gallic acid equivalent (GAE). The total phenolic content was 4.00±0.01mg/g for ethanolic extract, 2.26±0.02mg/g for acetone extract, 2.54±0.02mg/g for ethyl acetate extract and 1.21±0.01mg/g for aqueous extract of Gallic acid equivalent per 100mg seed extract of *Manilkara zapota* respectively (Table 1). The ethanolic extract of *Artocarpus heterophyllus* and *Manilkara zapota* seeds showed high phenol content than other different solvents.

Phenolic compounds are known as powerful chain breaking antioxidants and act as free radical terminators²¹, may contribute directly to antioxidative action²². These compounds are very important constituents of plants and their radical scavenging ability is due to their hydroxyl groups present in it²³. The mechanisms of action of flavonoids are through scavenging or chelating process^{24, 25}. In general, extracts with high antioxidant activity show high phenolics content. Polyphenols are the major plant compounds with antioxidants activity due to their redox properties; it plays an important role in absorbing and neutralizing free radicals, quenching singlet and triplet O₂ or decomposing peroxides⁹. Phenols contain plants has good antioxidant, anti-mutagenic and anti-cancer properties²⁶.

Determination of Total Flavonoid content

The total flavonoid content of the four different seed extract of *Artocarpus heterophyllus* was found to be 4.05±0.01mg/g for ethanolic fraction, 2.21±0.02mg/g for acetone fraction, 2.67±0.01mg/g for ethyl acetate fraction and 0.86±0.01mg/g for aqueous fraction of quercetin equivalent per 100mg seed extract, whereas in *Manilkara zapota*, it was found to be 3.98± 0.01mg/g (ethanolic), 2.19±0.01mg/g (acetone), 1.01±0.02mg/g (ethyl acetate) and 0.74± 0.01mg/g (aqueous) of Quercetin equivalent per 100mg (Table 1). The ethanolic extract of *Artocarpus heterophyllus* and *Manilkara zapota* seeds showed high flavonoid content than other three different solvents.

Flavonoids are a group of more than 4000 polyphenolic compounds that occur naturally in foods of plant origin. Flavonoids are a large class of phytochemicals which are omnipresent in human diets. Flavonoids are important for human health because of their high pharmacological activities as radical scavengers²⁷. The flavonoids are probably the most important natural phenolics due to their broad spectrum of chemical and biological activities, including antioxidant and free radical scavenging properties²⁸. Flavonoids have been reported as antioxidants, scavengers of a wide range of reactive oxygen species and inhibitors of lipid peroxidation, and also as potential therapeutic agents against a wide variety of diseases^{29, 30}.

Phenols and polyphenolic compounds, such as flavonoids, are widely found in food products derived from plant sources, and they have been shown to possess significant antioxidant activity³¹. The correlation between total phenol contents and antioxidant activity has been widely studied in different foodstuffs such as fruit and vegetables^{32, 33}.

Determination of Reduction potential

Reducing power is to measure the reductive ability of antioxidant, and it is evaluated by the transformation of Fe (III) to Fe (II) in the presence of the sample extracts³⁴. The results showed that reducing power of *Artocarpus heterophyllus* seed extracts was found to be 13.12±0.01µg/ml, 10.52±0.02µg/ml, 10.13±0.01µg/ml and 09.56±0.02µg/ml for ethanolic, acetone, ethyl acetate and aqueous extracts respectively, whereas reducing power of *Manilkara zapota* seed extracts was found to be 13.96±0.01µg/ml (ethanolic), 10.43±0.01µg/ml (acetone), 10.48±0.02µg/ml (ethyl acetate) and 09.41±0.02µg/ml (aqueous) and were showed in Table 1. The ethanolic extracts of *Artocarpus heterophyllus* and *Manilkara zapota* showed the highest reducing power and the values were comparable to that of phenolic content.

The reducing ability of a compound greatly depends on the presence of reductones, which have exhibit antioxidative potential by breaking the free radical chain by donating a hydrogen atom³⁵. The ability to reduce Fe (III) may be attributed from hydrogen donation from phenolic compounds³⁶, which is also related to presence of reductant agent. In addition, the number and position of hydroxyl group of phenolic compounds also rule their antioxidant activity³⁷.

The reducing capacity of a compound may serve as a significant indicator of its potential antioxidant activity. However, the activities of antioxidants have been attributed to various mechanisms such as prevention of chain initiation, decomposition of peroxides, reducing capacity and radical scavenging³⁸. Generally the reducing properties

are associated with the presence of compounds, which exert their action by breaking the free radicals chain through donating a hydrogen atom^{39, 40}.

DPPH radical scavenging assay

The DPPH system is a stable radical generating procedure. The DPPH assay is a simple method to measure the ability of antioxidants to trap free radicals. It is well known that the DPPH has ability to capture free radicals is due to the delocalization of the unpaired electron all over the molecule⁴¹.

The seed extracts of *Artocarpus heterophyllus* and *Manilkara zapota* showed promising free radical scavenging effect of DPPH in concentration dependent upto 500µg/ml. The free radicals scavenging activity of the *Artocarpus heterophyllus* seed extracts on DPPH radicals was found to have IC₅₀ of 50.09% at 300µg/ml for ethanolic, 51.90% at 400µg/ml for acetone, 52.28% at 500µg/ml for ethyl acetate and 51.80% at 500µg/ml for aqueous fraction (Table 2), whereas the *Manilkara zapota* seed extracts on DPPH radicals was found to have IC₅₀ of 51.08% at 300µg/ml for ethanolic, 50.28% at 400µg/ml for acetone, 51.99% at 500µg/ml for ethyl acetate and 50.46% at 500µg/ml for aqueous fraction (Table 2). Comparatively, ethanol extract is found to exhibit significant antioxidant activity in both seed extracts.

The DPPH radical has been widely used to test the ability of compounds as free-radical scavengers or hydrogen donors and to evaluate the antioxidative activity of plant extracts and foods^{42, 43}. The DPPH scavenging activity was found to be dose dependent. This radical scavenging activity of plants extracts could be related to the

nature of phenolics, thus contributing to their electron transfer/hydrogen donating ability.

ABTS radical scavenging assay

ABTS radical, a protonated radical has characteristic absorbance maxima at 734nm which decreases with the scavenging of the proton radicals⁴⁴. The ABTS radical cation scavenging activity of the *Artocarpus heterophyllus* seed extracts at 200 µg/ml was found to be IC₅₀ at 51.89% for ethanolic fraction, 52.09% at 300µg/ml for acetone, 50.36% at 400µg/ml for ethyl acetate and 52.04% at 500µg/ml for aqueous fraction (Table 3), whereas the *Manilkara zapota* seed extracts at 200µg/ml was found to be IC₅₀ at 50.32% for ethanolic fraction, 51.45% at 300µg/ml for acetone, 50.67% at 400µg/ml for ethyl acetate and 51.75% at 500µg/ml for aqueous fraction.

The methanolic extracts were fast and effective scavengers of the ABTS radicals⁴⁵. The high molecular weight phenolics have more ability to quench free radicals and their effectiveness depends on the molecular weight, the number of aromatic rings and nature of hydroxyl group substitution than the specific functional groups was reported by⁴⁶. Higher concentrations of the extracts were more effective in quenching free radicals in the system.

Phenols and flavonoids contribute to quality and nutritional value in terms of modifying colour, taste, aroma and flavour. The phenolic compounds act as antioxidant agents. As a whole the antioxidants are vital substances which possess the ability to protect body from damage by free radical induced oxidative stress.

Table 1: Total phenolic and flavonoid content, Reducing potential of different seeds extracts of *Artocarpus heterophyllus* and *Manilkara zapota*

Seed extracts	Different fractions	Total phenolic content, GAE (mg/g)	Total flavonoid content, QE (mg/g)	Reducing potential (µg/ml)
<i>Artocarpus heterophyllus</i>	Ethanolic	4.16±0.01	4.05±0.01	13.12±0.01
	Acetone	2.30±0.02	2.21±0.02	10.52±0.02
	Ethyl acetate	2.77±0.02	2.67±0.01	10.13±0.01
	Aqueous	1.18±0.01	0.86±0.01	09.56±0.02
<i>Manilkara zapota</i>	Ethanolic	4.00±0.01	3.98±0.01	13.96±0.01
	Acetone	2.26±0.01	2.19±0.01	10.43±0.01
	Ethyl acetate	2.54±0.02	1.01±0.02	10.48±0.02
	Aqueous	1.21±0.01	0.74±0.01	09.41±0.02

All values are expressed as Mean ± SD for three determinations; GAE- Gallic acid equivalent; QE -Quercetin equivalent

Table 2: DPPH radical scavenging assay of different seeds extracts of *Artocarpus heterophyllus* and *Manilkara zapota*

Seed extracts	Different fractions	Antioxidant activity - DPPH radical scavenging assay					IC ₅₀ Values (µg/ml)
		Different concentrations					
		100 µg/ml	200 µg/ml	300 µg/ml	400 µg/ml	500 µg/ml	
<i>Artocarpus heterophyllus</i>	Ethanolic	37.18±0.01	41.56±0.01	50.09±0.01	55.22±0.01	67.77±0.01	300
	Acetone	22.08±0.02	39.11±0.02	44.23±0.01	51.90±0.02	56.78±0.02	390
	Ethyl acetate	20.55±0.01	25.04±0.01	40.39±0.01	48.67±0.01	52.28±0.01	490
	Aqueous	31.12±0.02	33.45±0.02	42.75±0.02	45.33±0.04	51.80±0.02	480
<i>Manilkara zapota</i>	Ethanolic	36.07±0.01	49.52±0.01	51.08±0.01	56.13±0.01	64.49±0.01	290
	Acetone	22.37±0.01	37.91±0.01	46.83±0.01	50.28±0.01	52.83±0.01	400
	Ethyl acetate	19.14±0.02	28.27±0.02	33.38±0.02	44.93±0.02	51.99±0.02	490
	Aqueous	26.86±0.01	30.23±0.01	39.65±0.01	48.92±0.01	50.46±0.01	500

All values are expressed as mean ± SD for three determinations

Table 3: ABTS radical scavenging assay of different seeds extracts of *Artocarpus heterophyllus* and *Manilkara zapota*

Seed extracts	Different fractions	Antioxidant activity- ABTS radical scavenging assay					IC ₅₀ Values (µg/ml)
		Different concentrations					
		100 µg/ml	200 µg/ml	300 µg/ml	400 µg/ml	500 µg/ml	
<i>Artocarpus heterophyllus</i>	Ethanolic	31.26±0.02	39.93±0.01	51.89±0.02	57.20±0.02	62.01±0.01	290
	Acetone	19.98±0.02	27.54±0.02	41.78±0.01	52.09±0.01	52.78±0.01	380
	Ethyl acetate	21.02±0.01	27.37±0.01	39.23±0.02	46.23±0.01	50.36±0.01	500
	Aqueous	29.87±0.02	35.04±0.01	43.45±0.02	48.32±0.02	52.04±0.02	480
<i>Manilkara zapota</i>	Ethanolic	32.23±0.01	44.91±0.01	50.32±0.02	53.67±0.01	60.66±0.01	300
	Acetone	21.65±0.02	33.98±0.02	43.08±0.01	51.45±0.02	56.05±0.01	390
	Ethyl acetate	20.01±0.02	30.05±0.01	36.34±0.02	40.29±0.01	50.67±0.01	500
	Aqueous	28.46±0.01	29.78±0.01	40.39±0.01	45.67±0.02	51.75±0.01	490

All values are expressed as mean ± SD for three determinations

CONCLUSION

The results obtained in the present study, it was concluded that the seed extracts of both plants contains large amounts of flavonoids and phenolic compounds, exhibits high antioxidant and free radical scavenging activities in addition to having their medicinal properties. It also chelates iron and has reducing power. Both plant extract is a significant source of natural antioxidant, which might be helpful in preventing the progress of various oxidative stresses. Therefore, studies on these medicinal plants would be of great importance. Both plants are widely used in number of pharmacological actions with high content of flavonoids and phenol seems to have a high potential for antioxidant activity.

ACKNOWLEDGEMENT

The authors are grateful to School of Biotechnology, Dr. G.R. Damodaran College of Science, Coimbatore, Tamil Nadu, India for providing all facilities and for their encouragement.

REFERENCES

- Cheeseman KH, Scater TF. Free Radical in Medicine. In British medical bulletin. Churchill livingstone, London, 2003;9:479-724.
- Rao DB, Kiran CR, Madhavi Y, Roa PK, Roo TR. Evaluation of antioxidant potential of *Clitoria ternata*L and *Eclipta prostata*L. Ind J Biochem Biophy 2009; 46:247-252.
- Halliwell B, Gutteridge JMC. Oxidative stress and antioxidant protection: some special cases. In: Halliwell B, Gutteridge JMC, editors. Free Radicals in Biology and Medicine. 3rd edition. Oxford: Clarendon Press; 1999; 530-33.
- Sudha G, Sangeetha priya M, Indhu shree R, Vadivukkarasi S. Antioxidant activity of ripe pepino fruit (*solanum muricatum* aiton) Int J Pharm Pharm Sci, 2011;3: 257-261.
- Nakiboglu M, Urek OR, Kayali AH, Tarhan L. Antioxidant capacities of endemic *Sideritis sipylea* and *Origanum sipyleum* from Turkey. Food Chem 2007; 104: 630-635.
- Subhashini N, Thangathirupathi A, Lavanya N. Antioxidant activity of *Trigonella foenum graecum* using various in vitro and ex vivo models. Int J Pharm Pharm Sci, 2011; 3: 96-102.
- Halliwell B, Gutteridge JMC. Free radical in biology and medicine. 3rd Edition. London: Oxford. 1999; 36-40, 311-312.
- Daniel RS, Mathew BC, Devi KS. Antioxidant effect of two flavonoids from the bark of *Ficus bengalensis* Linn. in hyperlipidemic rats. Indian J Exp Bio 1998; 36: 902-906
- Zheng W, Wang SY. Antioxidant activity and phenolic compounds in selected herbs. J Agric Food Chem 2001; 49: 5165-5170.
- Seifried HE, Anderson DE, Fisher EI, Milner JA. A review of the interaction among dietary antioxidants and reactive oxygen species. J Nut Biochem 2007; 1: 1-8.
- Khopde SM, Priyadarshini KI, Guha SN, Satav JG, Venkatesan P, Rao MNA. Biosci Biotechnology Biochem 2000; 64: 503-509.
- Araceli S, Camen RM, Guillermo RS, Salvodar M, Rosa MG. Assessment of the anti-inflammatory activity and free radical scavenger activity of Tiliroside. Eur J Pharmacol 2003; 461:53-61.
- Rahman AM, Nahar N, Mian AJ, Mosihuzzaman M. Variation of carbohydrate composition of two forms of fruit from jack tree (*Artocarpus heterophyllus* L.) with maturity and climatic conditions. Food Chem 1999; 65: 91-97.
- Rama Rao AV, Mala V, Venkataraman. Colouring matter of the *A. hetetophyllus*. Indian J Chem 1973; 11: 298-299.
- Harbone JB. Phytochemical methods: A guide to modern techniques of plant analysis, 3rd Edition. Chapman and Hill publishers, London; 1998; 279.
- Siddhuraju P, Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agro climatic origins of Drumstick tree (*Moringa oleifera* Lam.) leaves. J Agric Food Chem 2003; 51: 2144-2155.
- Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. J Food Drug Anal 2002; 1:178-182.
- Oyaizu M. Studies on products of browning reaction prepared from glucosamine. Jpn. J Nutr 1986; 44:307-314.
- Mensor LL, Menezes FS, Leitao GG, Reis AS, Santos TC, Coube CS, Leitao SG. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. Phytotherapy Research 2001; 15:127-130.
- Shiwaiker A, Rajendran K, Kumar CD. In vitro antioxidant studies of *Annona squamosa* Linn leaves. Ind J Exp Biol 2006; 42:803-807.
- Shahidi F, Wanasundara PKJPD. Phenolic antioxidants. Critical Reviews in Food Science and Nutrition 1992; 32: 67-103.
- Duh PD, Tu YY, Yen GC. Antioxidant activity of water extract of harn jyu (*Chrysanthemum morifolium* Ramat). LWT-Food Sci Technol 1999; 32: 269-277.
- Hatano T, Edamatsu R, Hiramatsu M, Mori A, Fujita Y, Yasuhara D. Effects of interaction of tannins with co-existing substances. VI. Effects of tannins and related polyphenols on superoxide anion radical and on DPPH radical. Chem Pharm Bull 1989; 37: 2016-2021
- Cook NC, Samman S. Flavonoids- chemistry, metabolism, cardio protective effects, and dietary sources. Nutritional Biochemistry 1996; 7:66-76.
- Kessler M, Ubeaud G, Jung L. Anti- and pro-oxidant activity of rutin and quercetin derivatives. J Pharm Pharmacol 2003; 55: 131-142.
- Ahmad N, Mukhtar H. Green tea polyphenols and cancer: biological mechanisms and practical implications. Nutr Rev 1999; 57:78-83.
- Yao LH, Jiang YM, Shi J, Tomas-baeberan FA, Datta N, Singanusong R. Flavonoids in food and their health benefits. Plant Food Hum Nutr 2004; 59: 113-122.
- Kahkonen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS. Antioxidant activity of plant extracts containing phenolic compounds. J Agri Food Chem 1999; 47: 3954-3962.
- Ross JA, Kasum CM. Dietary flavonoids: Bioavailability, metabolic effects and safety. Annual Review of Nutrition 2002; 22: 19-34.
- Williams RJ, Spencer JP, Rice-Evans C. Flavonoids: Antioxidants or signalling molecules. Free Radical Biology and Medicine 2004; 36: 838-849.
- Van Acker SA, van Den Berg DJ, Tromp MN, Griffioen DH, Van Bennekom WP, van der Vijgh WJ and Bast A. Structural aspects of antioxidant activity of flavanoids. Free Radic Bio Med 1996; 20: 331-342.
- Kiselova Y, Ivanova D, Chervenkov T, Gerova D, Galunska B, Yankova T. Correlation between the *in vitro* antioxidant activity and polyphenol content of aqueous extracts from Bulgarian herbs. Phytother Res 2006; 20: 961-965.
- Kedage VV, Tilak JC, Dixit GB, Devasagayam TPA, Mhatre MA. Study of antioxidant properties of some varieties of grapes (*Vitis vinifera* L.). Crit Rev Food Sci Nutr 2007; 47: 175-185.
- Gulcin I, Oktay M, Kirecci E, Kufreviolu I. Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. Food Chem 2003; 83: 371-382.
- Duh PD. Antioxidant activity of Budrock (*Arctium laooa* Linn.) its scavenging effect on free radical and active oxygen. J Am Oil Chem Soc 1998; 75: 455-461.
- Shimada K, Fujikawa K, Yahara K, Nakamura T. Antioxidative properties of xanthan on the autoxidation of soyabean oil in cyclodextrin emulsion. J Agric Food Chem 1992; 40: 945-948.
- Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolics flavonoids. Free Radical Res 1995; 23: 375-383.
- Yildirim A, Mavi A, Oktay M, Kara AA, Algur OF, Bilaloglu V. Comparison of antioxidant and antimicrobial activities of *Tilia (Tilia argentea* Desf Ex DC), *Sage (Savia triloba* L.), and Black Tea (*Camellia sinensis*) extracts. J Agric Food Chem 2000; 48:5030-5034.
- Gordana R, Drazenka K, Sasa L, Dunja H, Maja K. Phenolic content and antioxidative capacity of green and white tea extracts depending on extraction conditions and the solvent used. Food chem 2008; 110:852-858.
- Dunja H, Drazenka K, Belscak A, Kovacevic GK, Damir I, Damir K. The composition of polyphenols and methylxanthines in teas and herbal infusions. Food chem 2009; 115:441-448.

41. Yokozawa T, Chen CP, Dong E, Tanka T, Nonaka GI, Nishioka I. Study of the inhibitory effect of tannins and flavanoids against the 1,1-Diphenyl-2-picrylhydrazyl radical. *Biochemical Pharmacology* 1998; 56: 13-22.
42. Porto CD, Calligaris S, Celloti E, Nicoli MC. Antiradical properties of commercial cognacs assessed by the DPPH test. *J Agric Food Chem* 2000; 48: 4241-4245.
43. Soares JR, Dinis TCP, Cunha AP, Almeida LM. Antioxidant activities of some extracts of *Thymus zygii*. *Free Rad Res* 1997; 26: 469-478.
44. Mathew S, Abraham ET. *In vitro* antioxidant activity and scavenging effects of *Cinnamomum verum* leaf extract assayed by different methodologies. *Food Chem Toxicol* 2004; 44:198-206.
45. Yu L, Haley S, Perret J, Harris M, Wilson J, Qian M. Free radical scavenging properties of wheat extracts. *J Agric Food Chem* 2002; 50: 1619-1624.
46. Hagerman AE, Riedl KM, Jones GA, Sovik KN, Ritchard NT, Hartzfeld PW. High molecular weight plant polyphenolics (Tannins) as biological antioxidants. *J Agric Food Chem* 1998; 46: 1887-1892.