

## IN VIVO AND IN VITRO PHYTOCHEMICAL, MICROBIAL AND ANTIOXIDANT EVALUATION OF SERICOSTOMA PAUCIFLORUM STOCKS EX WIGHT CALLUS

SATISH C. JAIN<sup>\*A</sup> BOSKEY PANCHOLI<sup>A</sup> AND RENUKA JAIN<sup>B</sup>

<sup>a</sup>Medicinal Plants and Biotechnology Laboratory, Department of Botany, University of Rajasthan, Jaipur, India. <sup>b</sup>Department of Chemistry, University of Rajasthan, Jaipur, India. Email: jainnatpro3@rediffmail.com.

Received: 29 Aug 2011, Revised and Accepted: 12 Nov 2011

### ABSTRACT

Cell cultures of *Sericostoma pauciflorum*, a promising antidiabetic plant, have been established using Murashige and Skoog's (MS) medium supplemented with different concentrations of 2, 4-dichlorophenoxyacetic acid (2, 4-D). Callus was harvested at different time intervals of 2, 4, 6, 8 and 10 weeks, and their antimicrobial and antioxidant potentials were investigated using established protocols. Methanol extracts demonstrated appreciable antifungal activity ( $17.33 \pm 0.34$  mm in 6 weeks-old and  $16.66 \pm 0.34$  mm in 4 weeks-old callus) against *P. chrysogenum* while antibacterial activity was recorded in 10 weeks-old callus against *B. subtilis* and *P. aeruginosa* ( $17.00 \pm 0.57$  mm and  $16.33 \pm 0.67$  mm respectively). Total levels of phenolics and antioxidant potentials were analyzed using Folin-Ciocalteu and DPPH, FRAP methods respectively. Methanol extracts of 8 weeks-old callus showed appreciable data (total phenolics  $114.00 \pm 1.80$  mg GAE/g extract; IC<sub>50</sub> 0.07 mg/ml, % inhibition 85.71 µg/ml), followed by 6 weeks-old callus (total phenolics  $110.32 \pm 0.50$  mg GAE/g extract; IC<sub>50</sub> 0.08 mg/ml, % inhibition 74.57 mg/ml). Later, the results were compared with those of *in vivo* studies.

**Keywords:** Antifungal, Antibacterial, Antioxidant, Phenolics.

### INTRODUCTION

*Sericostoma* is a small genus of family Boraginaceae. It comprises of eight species distributed through the tropical East and North East of Africa and North West India. *S. pauciflorum* Stocks ex Wight is a short straggling under-shrub growing widely throughout sea coast of Saurashtra and Maharashtra and used in making an important drug in Ayurveda named "Krishnavalli". It is anticancer, antidiabetic, used in dehydration, acidity, and health promoting drug (as described in "Nighantu Ratnakar"). Phytochemically, fernane, hopane and other type of triterpenoids<sup>1-4</sup> have been isolated.

The production of secondary metabolites through cell culture technology of renowned medicinal plants has been a challenging subject for many researchers. Various secondary metabolites were isolated from cell cultures of Boraginaceae family viz., from *Lithospermum erythrorhizon* and *Eritrichium sericeum* rosmarinic acid, which has biological activities including antioxidant, antiischemia reperfusion, antithrombosis, antihypertension, antifibrosis, antiviral and antitumor<sup>5-7</sup>; shikonin, a benzoquinone from *L. erythrorhizon*<sup>8,9</sup>; quinones and naphthoquinones from *Echium lycopsis* have been isolated and their antimicrobial activities were reported<sup>10-12</sup>.

Since *S. pauciflorum* extract is one of the important herbal formulations in ayurvedic medicine, as a tonic for treating many diseases and there is no report on its biological studies. Therefore, the present work deals with comparative studies of *S. pauciflorum* cell culture and plants extracts antibacterial, antifungal and antioxidant activities. This study adds further value for the possible use of this plant, especially the *in vitro*-induced callus extract, as a food additive.

### MATERIALS AND METHODS

#### Plant material

Whole plants of *Sericostoma pauciflorum* Stocks ex Wight (Boraginaceae), aerial parts and roots, were collected from the campus of University of Rajasthan, Jaipur during the months of July-October, 2009. The botanical identity was confirmed by Herbarium, Department of Botany, University of Rajasthan, Jaipur. Voucher specimen (No. 20155) of the plant has been deposited at the Herbarium and the Laboratory for further reference.

#### Callus induction and maintenance

Young nodal explants were washed under running tap water for 15 min, treated with 1% tween 20 (commercial detergent, Sigma) for 10 min and cut into small pieces of 1 inches each. Explants were

sterilized by successive treatment with abs. alcohol for 30 s, 0.1% (w/v) HgCl<sub>2</sub> for 4 min and later washing with sterilized distilled water (DW) four times. Sterilized explants were transferred on autoclaved Murashige and Skoog's basal culture medium<sup>13</sup> supplemented with different concentrations of 2,4-dichlorophenoxyacetic acid (2,4-D; 0.5, 1, 1.5 and 2 mg/l; Sigma), sucrose 30 g/l and 1% (w/v) agar (Merck). Biomass production in terms of growth index was recorded at different time interval of 2, 4, 6, 8 and 10 weeks<sup>14</sup>. Cultures were maintained on 25±2°C with the daily photoperiod of 16 hours. Out of the four 2, 4-D concentrations used, concentration yielding higher biomass of callus was selected and used for further study.

#### Preparation of extracts

Aerial parts with roots of *S. pauciflorum* was collected, dried and powdered. 50 g of this plant material was extracted with 100 ml of ethanol (3×18 hours). Similarly, callus was harvested at different growth periods of 2, 4, 6, 8 and 10 weeks, kept at 100°C for 5 min to inactivate the enzymatic activity and later, at 60°C till a constant weight is achieved. 20 g of callus of each growth period was separately powdered and extracted in 100 ml of ethanol (60°C, 3l×18 hours). Each sample was filtered using Whatman No. 1 filter paper in a Buchner funnel, the filtrate was freeze-dried and weighed. Later, the ethanolic extract was fractionated successively using pet. ether (60°- 80° C), methanol and water. Subsequently, each fraction was filtered, concentrated to dryness *in vacuo* and stored at appropriate temperature for further studies.

#### Phytochemical studies

Thin layer chromatography (TLC) profile of callus extracts was carried out on silica G plates using heptane-benzene-alcohol (100:100:1)(I) and butanol: 27% aqueous acetic acid (1:1 v/v)(II) as solvent systems. Plates were visualized by spraying with 10% SbCl<sub>3</sub> for solvent I and 10% methanolic AlCl<sub>3</sub> for solvent II. Several spots coinciding reference markers were scrapped from unsprayed plates, eluted with methanol, filtered, evaporated to dryness, reconstituted and crystallized in methanol. The melting point of the isolated compounds was determined in capillary tubes (Toshniwal Melting point apparatus) and subjected to ir spectrum (Perkin Elmer 337, Grating Infra red spectrophotometer).

The levels of terpenoids and phenolics acids were estimated colorimetrically<sup>15,7</sup>.

### Bioefficacies

The bacterial strains of *Bacillus subtilis* (MTCC 441), *Enterobacter aerogenes* (MTCC 111), *Escherichia coli* (ATCC 443), *Pseudomonas aeruginosa* (ATCC 741), *Raoutella planticola* (MTCC 530) and *Staphylococcus aureus* (ATCC 740) were obtained from IMTECH, Chandigarh, India, and maintained on Nutrient broth medium (NB) at 27°C for 48 hours. In fungus, *Aspergillus flavus* (ATCC 16870), *A. niger* (ATCC 322), *Candida albicans* (ATCC 4718), *Penicillium chrysogenum* (ATCC 5476) and *Tricophyton rubrum* (ATCC 2327) were obtained from IARI, New Delhi, India, and cultured on Sabouraud dextrose agar (SDA) at 37°C for 48 hours.

### Determination of antimicrobial activity

Antimicrobial tests were performed by agar well diffusion method<sup>16</sup>. Inoculum of each bacterial strain was suspended in nutrient broth and fungal strains in SDA broth at 25°C for period of 24 hours to the final cell concentration of 10<sup>6</sup>-10<sup>7</sup> CFU/ ml. Bacterial and fungal inoculums were inoculated in Müller-Hinton and SDA medium respectively. 4 mg extract in methanol was delivered to each well. To ensure diffusion of sample into agar, plates were incubated at 4°C for 1 hour, which were then incubated overnight at 25°C. The diameter of the inhibition zone around each hole was measured and recorded (Inhibition zone recorder, HiMedia, India). Streptomycin (10 mcg/disc) for bacteria and ketonocazole (10 mcg/disc) for fungus were used as positive controls.

### Total phenolic contents

The total phenolic contents were determined with Folin-Ciocalteu reagent<sup>17</sup> by measuring optical density (OD) at 750 nm (Shimadzu, Pharmaspec UV- Vis spectrophotometer). A standard calibration curve of gallic acid (1 - 50 mg/l) was prepared and total phenolics were expressed in mg of gallic acid equivalents (mg GAE/g of extract).

### Antioxidant activity

#### Free radical scavenging activity by 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH)

The effect on DPPH radical was determined using the method by Fogliano et al.<sup>18</sup>. Different concentrations of extract (0.8, 0.6, 0.4, 0.2, 0.1 mg/ml) were prepared in methanol and mixed with 2.5 mL of DPPH (2 mg/10 ml methanol). After 30 min of incubation time, absorbance was measured at 517 nm using UV-Vis spectrophotometer. Negative control (methanol) and positive

control (quercetin) were also used. Capability to scavenge the DPPH radical was calculated using following equation:

$$\text{DPPH scavenging effect (\%)} = [(A_0 - A_1 / A_0) \times 100]$$

Where, A<sub>0</sub> was the absorbance of the control reaction and A<sub>1</sub> was the absorbance in presence of the sample of given extract.


#### Total reduction capability by Ferric ion Reducing Antioxidant Potentials (FRAP).


Total reducing power of extracts was determined according to FRAP method<sup>19</sup>. Specific concentration of extracts (ascorbic acid) and extract (62.5-1000 µg/ml) was prepared in 1 mL ethanol separately, mixed with phosphate buffer (2.5 ml, 0.2 M, pH 6.6) and K<sub>3</sub>Fe(CN)<sub>6</sub> (2.5 ml, 1%). After incubation at 50°C (20 min), 2.5 ml of 10% tri chloroacetic acid, 2.5 ml of distilled water and FeCl<sub>3</sub> (0.5 ml, 1%; chromogenic reagent) were added in the sequence and absorbance was measured at 700 nm. A standard calibration curve of ascorbic acid was prepared and antioxidant activity was expressed in mg of ascorbic acid equivalents (mg AAE/g) of extract. All determinations were carried out in triplicate and statistically analyzed.

### RESULTS

Callus was initiated within 15 days, where MS medium with 2,4-D, 1.5 mg/l proved better for the callus biomass production and it was used for further experimentation. Colour and texture of the callus at different growth stages was also recorded. From callus cultures, several triterpenoidal compounds viz. friedelin, α- amyryrin, β- amyryrin, β- sitosterol and a phenolic compound-caffeic acid, have been isolated and identified on the basis of their chromatographic behaviour, melting points and spectral analysis (Table 1; Figure 1)

From the data, appreciable antimicrobial activity was demonstrated by pet. ether extract of 10 weeks- old callus as it is effective against *B. subtilis*, *P. aeruginosa*, *E. coli* (IZ 17.00±0.57, 16.33±0.67 and 15.66±0.34 mm respectively; Table 2), followed by its methanol extract against *S. aureus* (16.00±0.57 mm) and *E. coli* (15.33±0.34 mm). In antifungal activity, methanol extract of 4 weeks-old callus was more effective, showing appreciable activity against *P. chrysogenum*, *A. niger* (16.66±0.34 and 15.66±0.33 mm respectively), followed by 6 weeks-old callus methanol extract against *C. albicans*, *P. chrysogenum* and *T. rubrum* (17.33±0.34 and 14.66 mm in last two cases). Aqueous extract of callus showed negligible activity against most of the used microbes as compared to *in vivo* plant.





Caffeic acid

Fig. 1: Isolated compounds from *S. pauciflorum* cell culturesTable 1: Chromatographic behavior and chemical characteristics of isolated compounds from *S. pauciflorum* stem callus

Isolated compounds	R <sub>F</sub> (× 100)		Color after spray		m.p. (°C)	IR (ν <sub>max</sub> ) cm <sup>-1</sup> (KBr)
	I	II	I	II		
Friedelin	81	-	Pink	-	198-200	1720, 1380, 1365, 1255, 1230, 1200, 920
α-Amyrin	26	-	Pink	-	183-184	3350, 1640, 1480, 1360, 1130, 1050, 930
β- Amyrin	20	-	Pink	-	197-198	3350, 1650, 1190, 1140, 1100, 1050
β - Sitosterol	06	-	Blue	-	136-137	1730, 1640, 1240, 735, 725
Caffeic acid	-	76	-	Yellow	195-198	812, 849, 899, 972, 1118, 1172, 1212, 1448, 1640, 3440

I: Heptane-benzene-alcohol (100:100:1), spray with 10% SbCl<sub>3</sub>; II: Butanol: 27% aqueous acetic acid (1:1 v/v); spray with 10% methanolic AlCl<sub>3</sub>.

Table 2: Antimicrobial activity\* of *S. pauciflorum* stem callus at different growth stages

Age of callus	Type of extract	Bacteria					Fungi					
		<i>B. subtilis</i>	<i>E. aerogenes</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>R. planticola</i>	<i>S. aureus</i>	<i>A. flavus</i>	<i>A. niger</i>	<i>C. albicans</i>	<i>P. crysogenum</i>	<i>T. rubrum</i>
2 week	PE	14.33±0.34	13.33±0.67	12.67±0.3	15.00±0.00	10.66±0.66	11.00±0.57	15.66±0.66	12.33±0.33	10.00±0.00	12.66±0.33	12.00±0.5
	Met	13.33±0.00	10.00±0.57	10.66±0.3	13.00±0.57	10.33±0.67	10.00±0.00	15.00±1.00	11.33±0.33	13.66±0.67	16.00±0.00	13.00±0.0
	Aq	11.66±1.37	14.00±0.57	8.66±0.67	13.30±1.00	12.66±0.66	-	9.33±0.67	11.33±0.33	-	-	-
4 week	PE	14.33±0.34	10.00±0.00	12.67±0.3	12.33±0.76	10.66±0.66	-	9.33±0.67	13.33±0.33	9.66±0.88	13.66±0.67	12.66±0.3
	Met	13.33±0.00	12.00±0.57	10.66±0.3	13.00±0.00	10.00±0.57	10.66±0.66	15.00±0.57	15.66±0.33	13.66±0.34	16.66±0.37	15.00±0.0
	Aq	11.66±1.37	11.66±0.34	8.66±0.67	13.33±0.76	-	10.00±0.00	13.66±0.66	8.00±0.00	-	-	-
6 week	PE	11.66±0.67	8.66±0.66	10.66±0.3	15.00±0.00	10.00±0.00	10.00±0.00	13.00±0.66	12.66±0.67	11.33±0.33	13.33±0.67	13.00±0.5
	Met	14.33±0.34	13.33±0.66	12.66±0.6	11.00±0.00	11.00±0.00	13.66±0.66	13.00±0.57	13.66±0.34	17.33±0.34	14.66±0.67	14.66±0.3
	Aq	8.00±0.00	13.33±0.76	8.00±0.00	15.33±0.32	9.00±0.57	12.66±0.34	-	-	-	-	-
8 week	PE	15.66±0.67	10.66±0.67	14.66±0.6	15.33±0.32	9.66±0.67	10.00±0.00	11.66±0.87	14.66±0.67	8.66±0.66	12.66±0.66	14.33±0.3
	Met	15.66±0.67	-	15.00±0.0	15.33±0.32	12.66±0.66	16.00±0.76	13.66±0.66	13.66±0.34	12.33±0.34	13.66±0.66	15.66±0.6
	Aq	12.33±0.33	13.00±0.57	10.00±0.0	10.33±0.33	-	8.00±0.57	-	-	-	-	-
10 week	PE	17.00±0.51	-	15.66±0.3	16.33±0.67	15.00±0.00	10.33±0.76	-	-	-	12.33±0.67	13.33±0.3
	Met	15.00±0.00	14.33±0.34	15.33±0.3	15.00±0.00	13.00±0.57	16.00±0.57	-	-	-	14.00±0.57	15.66±0.6
	Aq	8.66±0.34	12.00±0.57	8.00±0.00	11.66±0.67	11.33±0.57	-	-	-	-	-	15.66±0.6
<i>In vivo</i> plant	PE	12.33±0.67	13.00±0.57	11.66±0.6	14.66±0.34	11.33±0.34	8.00±0.00	12.66±0.66	11.00±0.33	12.66±0.66	11.66±0.32	10.00±0.0
	Met	12.33±0.34	13.00±0.57	10.66±0.3	13.33±0.34	8.66±0.67	12.00±0.57	10.00±1.00	10.00±0.00	11.33±0.66	11.66±0.32	10.00±0.0
	Aq	15.00±0.57	10.33±0.32	10.33±0.3	15.66±0.66	10.33±0.32	10.33±0.32	10.00±0.00	12.66±0.74	11.66±0.32	13.33±0.32	9.33±0.34

Antimicrobial activity (in terms of inhibition zone in mm including the diameter of well; 6 mm); Mean  $\pm$  S.E. (Standard error); Standard: Streptomycin (10 mcg/ml) for bacteria, Ketocozole, (10 mcg/disc) for fungi; PE = pet. ether, Met = methanol, Aq = aqueous.

In methanol extracts, 114.00 $\pm$ 1.80 mg GAE/g of total phenolics was recorded in 8 weeks-old callus followed by 6 weeks-old callus (110.32  $\pm$  0.50 mg GAE/g). In the present study, enhanced

antioxidant activity was demonstrated by methanol extract of 8 weeks-old callus with lowest IC<sub>50</sub> value (0.07 mg/ml, % inhibition 85.71 on the concentration of 0.8 mg/ml; Table 3) followed by 6 weeks old-callus methanol (IC<sub>50</sub> 0.08 mg/ml, % inhibition of 75.51 in 0.8 mg/ml concentration). In FRAP method, methanol extract 10 weeks-old callus showed higher values of 60.00  $\pm$  0.00 AAE/g dw antioxidant potentials (Table 3).

**Table 3: Total phenolics and antioxidant assay by DPPH and FRAP method**

Age of callus	Type of extract	Total phenolics (mg GAE/g)	IC <sub>50</sub>	% Inhibition in DPPH method (concentration in mg/ml)					Antioxidant activity in FRAP method (in terms of AAE/g dw)				
				0.1	0.2	0.4	0.6	0.8	62.5	125	250	500	1000
2 week	PE	45.00 $\pm$ 0.61	0.47	45.51	46.50	48.07	55.25	55.12	14.66 $\pm$ 1.34	21.66 $\pm$ 2.79	23.33 $\pm$ 1.97	26.00 $\pm$ 1.66	30.00 $\pm$ 0.00
	Met	80.00 $\pm$ 1.25	0.62	32.68	34.78	46.10	48.76	62.27	43.33 $\pm$ 2.02	194.33 $\pm$ 1.49	214.33 $\pm$ 4.67	230.00 $\pm$ 5.78	300.00 $\pm$ 3.47
	Aq	59.66 $\pm$ 0.32	-	35.26	55.78	60.82	61.30	65.29	20.66 $\pm$ 3.33	20.33 $\pm$ 1.44	21.66 $\pm$ 5.78	25.66 $\pm$ 5.41	23.33 $\pm$ 1.72
4 week	PE	51.32 $\pm$ 0.32	0.11	49.16	54.40	60.00	60.36	60.61	21.66 $\pm$ 3.33	20.00 $\pm$ 0.00	25.00 $\pm$ 0.00	26.66 $\pm$ 5.78	36.66 $\pm$ 4.41
	Met	102.66 $\pm$ 0.72	0.09	54.97	63.55	65.90	69.15	74.62	95.00 $\pm$ 7.67	153.33 $\pm$ 1.49	165.00 $\pm$ 0.00	178.33 $\pm$ 1.67	320.00 $\pm$ 0.00
	Aq	78.00 $\pm$ 0.73	0.55	70.43	70.89	73.88	74.82	77.58	18.33 $\pm$ 0.02	21.66 $\pm$ 0.57	23.33 $\pm$ 1.98	23.33 $\pm$ 1.66	35.00 $\pm$ 0.00
6 week	PE	52.32 $\pm$ 1.58	0.085	58.29	58.85	59.59	60.31	61.46	18.33 $\pm$ 1.61	23.33 $\pm$ 5.78	25.00 $\pm$ 0.00	33.33 $\pm$ 6.62	36.66 $\pm$ 1.61
	Met	110.32 $\pm$ 0.50	0.08	53.41	58.77	63.40	66.57	75.58	121.66 $\pm$ 1.72	126.66 $\pm$ 2.97	226.66 $\pm$ 1.97	231.66 $\pm$ 1.66	340.66 $\pm$ 3.33
	Aq	78.00 $\pm$ 0.73	0.17	34.67	46.10	51.82	50.16	52.59	19.66 $\pm$ 7.55	24.00 $\pm$ 0.00	26.00 $\pm$ 0.00	28.66 $\pm$ 1.97	38.66 $\pm$ 0.33
8 week	PE	53.66 $\pm$ 0.15	0.14	47.17	53.02	60.51	61.28	66.36	19.00 $\pm$ 0.88	20.00 $\pm$ 1.51	25.66 $\pm$ 8.68	20.66 $\pm$ 7.64	41.66 $\pm$ 3.35
	Met	114.00 $\pm$ 1.80	0.07	68.94	69.79	74.87	83.00	85.71	25.00 $\pm$ 0.00	126.66 $\pm$ 7.27	235.00 $\pm$ 0.00	236.66 $\pm$ 5.78	360.00 $\pm$ 0.00
	Aq	73.00 $\pm$ 0.73	0.08	57.78	60.84	63.86	68.23	73.01	28.33 $\pm$ 1.66	31.66 $\pm$ 6.12	34.33 $\pm$ 4.94	35.00 $\pm$ 0.00	43.33 $\pm$ 5.79
10 week	PE	47.32 $\pm$ 0.72	0.095	51.98	52.64	55.22	55.50	56.04	20.00 $\pm$ 0.00	23.33 $\pm$ 7.82	23.33 $\pm$ 4.23	28.33 $\pm$ 3.78	42.33 $\pm$ 7.72
	Met	75.00 $\pm$ 0.60	0.10	49.60	63.25	65.06	67.33	74.57	125.00 $\pm$ 0.00	126.66 $\pm$ 7.27	235.00 $\pm$ 0.00	236.66 $\pm$ 5.78	360.00 $\pm$ 0.00
	Aq	71.00 $\pm$ 0.73	-	28.36	30.13	34.98	37.64	89.99	30.00 $\pm$ 0.00	35.00 $\pm$ 0.00	46.66 $\pm$ 3.33	53.33 $\pm$ 6.67	54.33 $\pm$ 6.01
In vivo plant	PE	31.5 $\pm$ 1.67	-	68.61	70.61	70.61	74.40	73.75	18.33 $\pm$ 4.41	21.00 $\pm$ 6.69	25.00 $\pm$ 0.00	31.66 $\pm$ 3.78	46.66 $\pm$ 3.33
	Met	84.93 $\pm$ 0.15	0.15	47.32	51.80	72.60	72.93	80.04	110.00 $\pm$ 3.33	211.66 $\pm$ 7.24	220.00 $\pm$ 1.95	275.00 $\pm$ 0.00	351.66 $\pm$ 9.36
	Aq	104 $\pm$ 0.44	0.075	68.48	76.92	92.51	93.40	93.78	32.56 $\pm$ 5.66	32.66 $\pm$ 4.56	43.99 $\pm$ 3.33	51.99 $\pm$ 3.48	51.11 $\pm$ 5.69

PE = pet. ether, Met = methanol, Aq = aqueous.

On quantification, caffeic acid was estimated to be maximum in the methanolic extract of 6 weeks- and 8 weeks-old callus (4.54  $\pm$  1.56 mg/g dw and 4.42  $\pm$  3.15 mg/g dw respectively). Out of the

terpenoids extracted, levels of  $\beta$ -sitosterol was found to be maximum (12.8  $\pm$  2.71 mg/g dw) as compared to other compounds (Table 4).

**Table 4: Quantification of isolated secondary metabolites**

Nature of extract		Concentration ( mg/g dw)	
		$\beta$ - Sitosterol	Caffeic acid
Callus	2 weeks-old	3.00 $\pm$ 4.42	1.58 $\pm$ 2.78
	4 weeks-old	5.70 $\pm$ 0.71	2.56 $\pm$ 0.25
	6 weeks-old	7.00 $\pm$ 3.94	4.54 $\pm$ 1.56
	8 weeks-old	7.30 $\pm$ 1.44	4.42 $\pm$ 3.15
	10 weeks-old	12.8 $\pm$ 2.71	3.32 $\pm$ 1.33
Whole plant		14.0 $\pm$ 1.66	2.24 $\pm$ 5.64

## DISCUSSION

In the developing countries, infectious diseases due to development of antibiotic resistance related to several health problems<sup>20</sup>. Epidemiological studies suggested that phytosterols reduces the risk

of colon cancer<sup>21, 22</sup>.  $\beta$ -Sitosterol found in almost all plants and work against human breast cancer cells<sup>23</sup>. Gram +ve bacteria were more susceptible to phenolic acids (*p*-coumaric-, sinapic- and caffeic acids) than Gram -ve bacteria, possibly due to the presence of their outer

membrane spaces<sup>24, 25</sup>. Moreover, *S. aureus* and *B. subtilis* were the most susceptible Gram +ve bacteria, an observation that may be attributed to the presence of a single membrane of the organisms which makes it more accessible to permeation by active principles of *S. pauciflorum* active extracts.

In the present study, methanol extract demonstrated greater antioxidant efficacy as compared to pet. ether and water extracts, where the levels of phenolic contents were more than the others. Similar results were obtained by FRAP method also which were better than the *in vivo* plant. Free radicals are responsible for several disorders in humans such as atherosclerosis, arthritis, ischemia and central nervous system injury<sup>26, 27</sup>. It is known that compounds called as antioxidants, scavenge these free radicals and protect our body system. In recent years due to toxic effect of commercially available antioxidants, medicinal plants as antioxidants have been used and they are found more effective than their isolated forms<sup>28</sup>. Phytochemical screening indicated the presence of phenolic acids, which are mainly responsible for the antimicrobial and antioxidant effect of this plant callus.

## CONCLUSION

The successful establishment of callus cultures of *S. pauciflorum* for the secondary metabolites production, antimicrobial and antioxidant potentials has been demonstrated. Terpenoids and phenolic metabolites play an important role in human health and can be strong scientific evidence to use this plant as a useful source of both biological and pharmacological references. Establishment of standard protocols for the production of these potent antimicrobial and antioxidant compounds could give rise to the commercial production of valuable metabolites. Further studies on phytochemical investigation are underway to isolate the bioactive phytochemicals leading to isolation of effective drug(s) from this plant callus.

## ACKNOWLEDGEMENT

Authors are thankful to the Indian Council of Medical Research, New Delhi, India, for partial financial support.

## REFERENCES

- Afza N, Bader A, Malik A, Ayatollahi AM, Ahmed Z, Khan AQ. Structural studies of triterpenoid isolated from some *Euphorbia* species and *Sericostoma pauciflorum*. Proc 1<sup>st</sup> International Conf Pharm Sci 1992; 36-56.
- Ayatollahi SAM, Ahmed Z, Afza N, Malik A. A triterpene from *Sericostoma pauciflorum*. Phytochem 1992; 31: 2899-2901.
- Ayatollahi SAM, Ahmed Z, Malik A. A fernane-type triterpene from *Sericostoma pauciflorum*. J Nat Prod 1991; 54: 570-572.
- Ayatollahi SAM, Ahmed Z, Malik A, Afza N, Bader A. A Hopane type triterpenoid from *Sericostoma pauciflorum*. Fitoterapia 1992; LXIII: 304-307.
- Bryukhanov VM, Bulgakov VP, Zverev YF, Fedoreev SA, Lampatov VV, Veselovam MV et al. An *Eritrichium sericeum* Lehm. (Boraginaceae) cell culture-a source of polyphenol compounds with pharmacological activity. Pharm Chem J 2008; 42: 344-347.
- Jianga RW, Laua KM, Hona PM, Makb TCW, Wooc KS, Fung KP. Chemistry and biological activities of caffeic acid derivatives from *Salvia Miltiorrhiza*. Current Med Chem 2005; 1: 237-246.
- Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL. Lipophilic and hydrophilic antioxidant capacities of common foods in the United States. J Agric Food Chem 2004; 52: 4026-4037.
- Bulgakov VP, Kozyrenko MM, Fedoreyev SA, Zhuravlev YN. Shikonin production by para-fluorophenylalanine resistant cells of *Lithospermum erythrorhizon*. Fitoterapia 2001; 72: 394-401.
- Fukui H, Feroj HAFM, Tomohiro U, Masaharu K. Formation and secretion of a new brown benzoquinone by hairy root culture of *Lithospermum erythrorhizon*. Phytochem 1998; 47: 1037-1039.
- Fukui H, Tsukada M, Hajime M, Tabata M. Formation of stereoisomeric mixture of naphthoquinone derivatives in *Echium lycopsis* callus culture. Phytochem 1983; 22: 453-456.
- Inouye H, Matsumura H, Kawasaki M, Inoue K, Tsukada M, Tabata M. Two quinone from callus culture of *Echium lycopsis*. Phytochem 1981; 20: 1701-1706.
- Tabata M, Tsukada M, Fukui H. Antimicrobial activity of quinone derivatives from callus culture of *Echium lycopsis*. Planta Med 1982; 44: 234-236.
- Murashige T, Skoog F. A revised medium for rapid growth and bioassay with tobacco tissue cultures. Physiol Plant 1962; 15: 473-497.
- Nezbedova L, Hesse M, Dusek J, Werner C. Chemical potential of *Aphelandra* sp. cell cultures. Plant Cell Tissue Organ Cult 1999; 58: 133-140.
- Das AK, Banerjee AB. A rapid method for quantification of sterols after thin-layer chromatography. Indian J Exp Biol 1980; 18: 969-971.
- Boyanava L, Gergova G, Nikolov R, Derejian S, Lazarova E, Katsarov N et al. Activity of Bulgarian propolis against 94 *Helicobacter pylori* strains *in vitro* by agar well diffusion, agar dilution and disc diffusion methods. J Med Microbiol 2005; 54: 481-483.
- Bray HG, Thorpe WV. IN Glick, D, editors. Methods of Biochemical Analysis. New York: John Wiley & Sons; 1954. p. 27-57.
- Fogliano V, Verde V, Randazzo G, Ritieni A. A method for measuring antioxidant activity and its application to monitoring the antioxidant capacity of wines. J Agric Food Chem 1999; 47: 1035-1040.
- Yen GC, Chen HY. Antioxidant activity of various tea extracts in relation to their antimutagenicity. J Agric Food Chem 1995; 43: 27-32.
- Murugan S, Uma Devi P, Parameswari NK, Mani KR. Antimicrobial activity of *Syzygium jambos* against selected human pathogens Intl J Pharm Pharm Sci 2011; 3(2): 44-47.
- Hirai K, Shimazu C, Takezoe R, Ozek Y. Cholesterol, phytosterol and polyunsaturated fatty acid levels in 1982 and 1957 Japanese diets. J Nutr Sci 1986; 32: 363-372.
- Nair P, Turjman N, Kessie G, Calkins B, Goodman G. Diet, nutrition intake, and metabolism in populations at high and low risk for colon cancer. Dietary cholesterol, beta-sitosterol, and stigmaterol. American J Clin Nutr 1984; 40: 927-930.
- Awad BA, Roy R, Fink CS. Beta-sitosterol, a plant sterol, induces apoptosis and activates key caspases in MDA-MB-231 human breast cancer cells. Oncol Rep 2003; 10: 497-500.
- Adesokan AA, Akanji MA, Yakubu MT. Antibacterial potentials of aqueous extract of *Enantia chlorantha* stem bark. Afr J Biotechnol 2007; 6: 2502-2505.
- Nikaido H. Microdermatology Cell surface in the interaction of microbes with the external world. J Bacteriol 1999; 181: 4-8.
- Nehete J, Bhatia M. Correlation of antioxidant activity with phenolic content and isolation of antioxidant compound from *Lygodium flexuosum* (L.) sw. extracts. Intl J Pharm Pharm Sci 2011; 3(2): 48-52.
- Kumpulainen JT, Salonen JT. Natural Antioxidants and Anticarcinogens in Nutrition, Health and Disease. UK: The Royal Society of Chemistry; 1999. p. 178-187.
- Khopde SM, Priyadarsini KI, Mohan H, Gawandi VB, Satav JG, Yakhmi JV et al. Characterizing the antioxidant activity of amla (*Phyllanthus emblica*) extract. Curr Sci 2001; 81: 185-190.