

STUDIES ON STANDARDISATION OF *ANDROGRAPHIS PANICULATA* NEES AND IDENTIFICATION BY HPTLC USING ANDROGRAPHOLIDE AS MARKER COMPOUND

ARJUN SINGH^{*1}, A K MEENA², SUDEEP¹, MEENA³, P. PANT¹, M M PADHI¹

¹Central Council for Research in Ayurvedic Sciences, Janakpuri, Delhi-110058, ²National Institute of Ayurvedic Pharmaceutical Research, Patiala - 47001, Punjab, ³Doon College of Engineering & Technology, Sunderpur, Saharanpur, India.
Email: arjuncrcas@gmail.com; ajaysheera@yahoo.com

Received: 17 Oct 2011, Revised and Accepted: 30 Nov 2011

ABSTRACT

The present communication attempts to evaluate the physicochemical and preliminary phytochemical studies on the extract (Hydro-alcoholic) of *Andrographis paniculata* Nees. (*Acanthaceae*). It is widely found and cultivated in tropical and subtropical Asia, south-east Asia and India. It is wide range of pharmacological effects and some of them extremely beneficial such as anti-inflammatory, anti-diabetes, anti-diarrhoeal, antiviral, antimalarial, hepatoprotective, anticancer, antihuman immunodeficiency virus (HIV), immune stimulatory and antsnakebite activity. In Physicochemical parameters studies LOD was found to be 4.54 %, Total ash content was 28.57 % and acid-insoluble was 5.45%. The water-soluble and alcohol soluble extractive values were found to be 88.27, 55.47% respectively. Other parameters like density, solvent residue were also analysed. The levels of toxic heavy metals and microbial contamination was indicated in such herbal drugs was in permissible limit as per WHO specification. The data indicated suggest that there is requirement of in process improvement to provide better quality for consumer health in order to be competitive in international markets.

Keywords: *Andrographis paniculata* Nees., Toxic metals, Physicochemical studies.

INTRODUCTION

Andrographis paniculata Nees. is a slender upright annual varying in height from 30 to 100 cm (1 to 3 feet), with a square stem and "lanceolate" leaves (i.e., shaped like a lance, sharp at the ends and curved in the middle). *Andrographis paniculata* Nees. is an erect annual herb extremely bitter in taste in all parts of the plant body. The plant is known in north-eastern India as *Maha-tita*, literally "king of bitters", and known by various vernacular names like Chiorta in Assamese, Oli-kiryata in Marathi, Kalmegh in Bengali, Bhuinimba in Oriya, Green chirayta, creat, andrographis, India Echinacea in English, Kariyatu in Gujarati, Kālamegha, Bhūnimba in Sanskrit¹, Kirayat in Hindi, Nilavembu in Tamil, Nelaberu in Kannada, Nelavemaa or Nelavepu meaning "Neem of the ground" in Telugu. Nilavembu in Malayalam.

Since ancient times, *Andrographis paniculata* Nees. is used in traditional Siddha and Ayurvedic² systems of medicine as well as in tribal medicine in India and some other countries for multiple clinical applications. As an Ayurveda herb it is known as *Kalmegh* or *Kalamegha*, meaning "dark cloud". It is also known as *Bhui-neem*, meaning "neem of the ground", since the plant, though being a small annual herb, has a similar strong bitter taste as that of the large Neem tree (*Azadirachta indica*). The genus *Andrographis* consists of 28 species of small annual shrubs essentially distributed in tropical Asia. Only a few species are medicinal, of which *Andrographis paniculata* Nees. is the most popular.

Hydro-alcoholic extract of *Andrographis paniculata* Nees. is traditionally used as a medicine to treat different diseases in India, China and Southeast Asia including Malaysia. The leaves and roots have traditionally been used over the centuries in Asia and Europe as a folklore medicine for a wide variety of ailments or as herbal supplements for health promotion. In traditional Chinese medicine, it is widely used to get rid of body heat, as in fevers and to dispel toxins from the body. In Scandinavian countries, it is commonly used to prevent and treat common cold³. Previous studies have explicitly revealed that *Andrographis paniculata* Nees. has a wide range of pharmacological effects and some of them extremely beneficial such as anti-inflammatory⁴, anti-diabetes⁵, anti-diarrhoeal⁶, antiviral⁷, antimalarial⁸, hepatoprotective⁹, anticancer¹⁰, antihuman immunodeficiency virus (HIV)¹¹, immune stimulatory¹², and antsnakebite activity¹³. Diterpenoids and flavonoids are the main chemical constituents of *A. paniculata* which are believed to be responsible for the most biological activities of this plant¹⁴. Andrographolide is the major constituent extracted from the leaves

of the plant which is a bicyclic diterpenoid lactone. Bitter diterpenoid lactones, especially 14-Deoxy-11-dehydroandrographolide, 14-Deoxy-11-oxoandrographolide, 5-Hydroxy-7, 8, 2', 3'-Tetramethoxy flavones, Andrographolide, Neoandrographolide, Paniculide-A, Paniculide-B, Paniculide-C, have been isolated from the whole plant and leaves. Andrographine, Panicoline, Diterpene dimers Flavonoids available in the roots. The main active constituents and marker compounds are considered to be the andrographolides and andrographis extracts are often standardized to these compounds¹⁵.

Important actions according to Ayurveda of *Andrographis paniculata* Nees. (*Kalmegh*) balances both Kapha and Pitta Doshas; so it can be used in all health problems originating from aggravation of Pitta or Kapha or both. *Andrographis paniculata* Nees. ignites the digestive fire in stomach, stimulates perspiration and removes toxins from the body via sweat. *Andrographis paniculata* Nees. works well against parasites and microbes, stimulates expulsion of bile from liver. *Andrographis paniculata* Nees. is indicated in all disorders of liver. *Andrographis paniculata* Nees. is useful in skin diseases. The Tamils have been using Nilavempu - as it is called in Tamil - for centuries. In Siddha medicine, *Andrographis paniculata* Nees. is used widely to treat fevers like chikenguinea, swine-flu, typhoid etc¹⁶.

MATERIALS AND METHODS

Extract material

The extract (Hydro: Alcoholic-60:40) of *Andrographis paniculata* Nees was procure from GMP certified firm. Sample was studied for physicochemical evaluation and identification.

Methods for Physicochemical Parameter

Moisture content

4 g of the sample was taken and heated in an oven at 105°C for 5 hour in a previously weighed 100 ml beaker. It was cooled in desiccators and weighed. The procedure was repeated till constant weight is obtained. The percentage of loss in weight of the sample was calculated.

Deterioration time of the plant material depends upon the amount of water present in plant material. If the water content is high, the plant can be easily deteriorated due to fungal attack. The loss on drying at 105°C of *Andrographis paniculata* extract was found to be 4.54% w/w.

Determination of Total ash

2 g of the sample was taken accurately in a previously ignited and tarred Silica dish. The material was spread evenly and ignited in a muffle furnace by gradually increasing the temperature to 600°C until it is white, indicating the absence of carbon. The crucible was cooled in desiccators and allowed to stand for 30 minutes and weighed.

Total ash value of plant material indicated the amount of minerals and earthy materials attached to the plant material. Analytical results showed total ash value of *Andrographis paniculata* extract was 28.57 % w/w.

Determination of Acid insoluble ash

To the dish containing the total ash, 25 ml of 20 % Hydrochloric acid was added covered with a watch glass and boiled gently for 5 minutes. The watch was rinsed with a hot water and added to the crucible. The residue was washed with the hot water till the washings were neutral to the litmus. The insoluble material was collected and again placed in a same crucible and again ignited for 6 hr. to constant weight. The residue was cooled a desiccators for 30 minutes and weighed.

Percentage of acid insoluble as was calculated. The amount of acid-insoluble siliceous matter present in the *Andrographis paniculata* extract was 5.45% w/w.

Determination of Water soluble extractive value

4 g of the sample was taken in a glass stoppered flask. 100 ml of distilled water was added. The flasks were shaken occasionally for 6 hours and then allowed to stand for 18 hours. The extract was filtered and 25 ml of the filtrate was pipette out in a pre-weighed 100 ml beaker and evaporated to dryness on a water bath. It was kept in a hot air oven for 5 hr at 105°C, cooled in desiccators for 30 minutes and weighed. The procedure was repeated till constant weight.

The water-soluble extractive value indicated the presence of sugar, acids and inorganic compounds. The water soluble extractive value in the *Andrographis paniculata* extract sample was found to be 88.27% w/w.

Determination of Alcohol soluble extractive Value

Same procedure as for the water soluble extractive value was followed. Instead of water, rectified spirit was taken as a solvent.

The alcohol soluble extractive values indicated the presence of polar constituents like phenols, alkaloids, steroids, glycosides, flavonoids and secondary metabolites present in the plant sample. The alcohol soluble extractive value was found to be 55.47% w/w in the *Andrographis paniculata* extract.

Determination of pH Value

10% aqueous solution of sample was prepared and used for determining the pH value by pH meter. The pH value of *Andrographis paniculata* extract was found to be 7.33.

HPTLC analysis was performed on HPTLC apparatus (CAMAG, Swtzerland) and Aluminium plate pre-coated with silica gel 60 F₂₅₄ of 0.2 mm thickness (Merck, Germany) was used for HPTLC analysis.

Analytical Procedure

2 g each of extract was soaked overnight in 20 ml of 90% ethanol. The solution was continuously stirred for 6 hrs and kept for next 18 hrs. Next day filtered the samples, dried and Made 10% solution. The solution was applied as 10 mm band together with a separate band of standard on Merck Aluminium plate pre-coated with silica gel 60 F₂₅₄ of 0.2 mm thickness. The plates were developed in Toluene: Ethylacetate: Acetic acid (4.0: 6.0: 0.5). The plate was dried and visualized under UV 254 & 366 nm. The plate was then dipped in Anisaldehyde-Sulphuric acid, heated at 105° C till the colour of the resolved bands appeared and visualized under white light. Marker compound andrographolide was used for the conformity of the samples used for the analysis.

RESULTS AND DISCUSSION

Hydro-alcoholic extract of the *Andrographis paniculata* Nees. was collected and analysed for the various standardisation parameters. Preliminary phytochemical results showed the presence or absence of certain phytochemical in the drug. Phytochemical test revealed the presence, Alkaloid, triterpene, saponins, flavonoids, polysaccharides, Steroid and Tannin results are given in Table1.

Table 1: Qualitative analysis of the phytochemical in the extracts of *Andrographis paniculata* Nees.

S. No.	Phytochemicals Constituents	Test performed	Result
1.	Alkaloid	Dragendorff's test	+ve
2.	Flavonoids	Shinoda test	+ve
3.	Steroid	Liebermann-Burchard reagent	+ve
4.	Tannin	Neutral FeCl ₃	+ve
5.	Glycoside/Sugar	Molisch's test	+ve
6.	Terpenoid	Noller's test	+ve
7.	Saponin	NaOH solution	+ve

The presence of three heavy metals namely Arsenic, Cadmium, Lead and microbial contamination were analysed in the sample and the results are shown in Table 2. The concentration of all the heavy metals and microbial contamination were below the WHO/FDA permissible limits^{17,18}.


Physico-chemical parameters of the *Andrographis paniculata* Nees. (Hydro-alcoholic extract) are tabulated in Table 2. The pH value of 10% w/v aqueous solution is 7.33. Deterioration time of the plant material depends upon the amount of water present in plant material. If the water content is high, the plant can be easily deteriorated due to fungus. The loss on drying at 105°C was found to be 4.54 %. Total ash value of plant material indicated the amount of minerals and earthy materials attached to the plant material. Analytical results showed total ash content was 28.57 %. The amount of acid-insoluble /siliceous matter present in the plant was 5.45%. The water-soluble extractive value was found to be 88.27; it is indicating the presence of sugar, acids, inorganic compounds and other water soluble matter. The amount of alcohol soluble extractive

values was 55.47%; it indicated the presence of polar constituents like phenols, alkaloids, steroids, glycosides, flavonoids. The results of other parameters like density, solvent residue, assay etc are given in Table 2.

Thin layer chromatographic technique was used to separate the chemical compounds present in the drug. Various solvent systems were checked to separate the maximum number of chemical compounds in the drug. HPTLC of the ethanolic solution developed in the mobile phase of Toluene: Ethyl acetate: Acetic Acid (4.0: 6.0: 0.5) (Figure. 1) and on observation under UV 254 nm 09 bands of sample at R_f value 0.31, 0.38, 0.45, 0.55, 0.61, 0.66, 0.77, 0.82, 0.96 and 01 band of standard R_f 0.38, (all bands are dark green colour) ; under UV 366 nm 09 bands of sample at R_f 0.07, 0.18, 0.39, 0.46, 0.55, 0.60, 0.67, 0.73, 0.77 and no visible band of standard and after derivatisation with Anisaldehyde-sulphuric acid under white light 10 bands of sample at R_f 0.11, 0.15, 0.32, 0.38, 0.44, 0.55, 0.66, 0.74, 0.83, 0.88 and one band of standard at R_f 0.38 were observed.

Table 2: Physico-chemical parameters of *Andrographis paniculata* Nees.

S. No.	Parameters	Results
1.	Description	Brown Colour Dry Powder
2.	Extraction Medium	Ethanol : Water (40 : 60)
3.	Loss on drying at 105 °C	4.54 % w/w
4.	Total Ash	28.57 % w/w
5.	Acid-insoluble ash	5.45 % w/w
6.	Water-soluble extractive	88.27 % w/w
7.	Alcohol-soluble extractive	55.47 % w/w
8.	pH (10 % w/v aqueous suspension)	7.33
9.	Passing through 40 mesh size sieve	95.28 % w/w
10.	Bulk density	0.68gm/ml
11.	Tap density	1.04 gm/ml
12.	Solvent residue (Ethanol)	<500 ppm
13.	Heavy metals	
a)	Lead	< 5 ppm
b)	Cadmium	<1 ppm
c)	Arsenic	<2 ppm
14.	Microbial contamination	
	Test for E.coli/g	Absent
	Test for Salemonella/g	Absent
	Test for S. aureus/g	Absent
	Total viable count/g	300cfu/gm
	Total fungal count/g	<10 cfu/gm
	Total enterobacteriaceae/g	Absent

Fig. 1: HPTLC Profile *Andrographis paniculata* Nees.

CONCLUSION

Preliminary phytochemical as well as various aspects of the *Andrographis paniculata* Nees. extract (Hydro: Alcoholic-60:40) was studied and described along with, physico-chemical, toxic/heavy metal, microbial contaminants and HPTLC studies to authenticate the adulteration for quality control of herbal drugs. HPTLC profile together with marker compound may economically verify the identity of this plant material in any formulation. *Andrographis paniculata* Nees. extract exhibits a set of diagnostic characters, which will help to identify the drug in dried condition and in other formulations.

It has been concluded from this study that estimation of heavy metals and microbial contamination is highly essential for raw drugs or plant parts used for the preparation of compound formulation drugs. The periodic assessment is essential for quality assurance and safer use of herbal drugs.

ACKNOWLEDGEMENT

The authors are very grateful to Director General, CCRAS, New Delhi and for providing encouragement and facilities for carrying out this work.

REFERENCE

1. Sanskrit synonyms of bhunimb Amarkosha ch. 2, section - Forest medicinal plants, verse, 143.
2. Nighantu Adarsh. Medicinal properties of bhunimb.
3. Cacer DD, Hancke JL, Burgos RLA & Wickman JK. Prevention of common cold with *Andrographis paniculata* Nees. dried extract. A pilot double blind trial. *Phytomedicine*, 1997; 4: 101-104.
4. Shen YC, Chen CF & Chiou WF. Andrographolide prevents oxygen radical production by human neutrophils: possible mechanism(s) involved in its anti-inflammatory effect. *British Journal of Pharmacology*. 2002; 135: 399-406.
5. Syahrin A, Amrah S, Chan K, Lim B, Hasenan N, Hasnan J & Mohsin S. Effect of spray dried ethanolic extract of *Andrographis paniculata* (Burms F.) Nees on strepto - zotocin induced Diabetic Rats. *International Journal of Diabetes in Developing Countries*. 2006; 26: 163-168.
6. Gupta S, Ahmad MC, Yadava JNS, Srivastava V & Tandon JS. Antidiarrhoeal activity of diterpenes of *Andrographis paniculata* Nees. (Kal-Megh) against *Escherichia coli* enterotoxin *in vivo* Models. *Journal of Pharmaceutical Biology*, 1990; 28: 273-283.
7. Wiart C, Kumar K, Yusof MY, Hamimah H, Fauzi ZM & Sulaiman M. Anti-viral properties of ent-labdene diterpenes of *Andrographis paniculata* Nees. *Phytotherapy Research*. 2005; 19: 1069-1070.
8. Rahman NN, Furuta T, Kojima S, Takane K & Mohd MA. Antimalarial activity of extracts of Malaysian medicinal plants. *Journal of Ethnopharmacology*. 1999; 64: 249-254.
9. Trivedi N & Rawal UM. Hepatoprotective and toxicological evaluation of *Andrographis paniculata* on severe liver damage. *Indian Journal of Pharmacology*, 2005; 32:288-293.
10. Zhou J, Zhang S, Ong CN & Shen HM. Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells. *Biochemical Pharmacology*. 2006; 72: 132-144.
11. Calabrese C, Berman SH & Babish JG. A phase I trial of andrographolide in HIV positive patients and normal volunteers. *Phytotherapy Research*. 2000;14: 333-338.
12. Iruretagoyena MI, Tobar JA & Gonzalez PA. Andrographolide interferes with T-cell activation and reduces experimental autoimmune encephalomyelitis in the mouse. *Journal of Pharmacology & Experimental Therapeutics*, 2005; 312: 366-372.
13. Samy RP, Thwin MM, Gopalakrishnakone P & Ignacimuthu S. Ethnobotanical survey of folk plants for the treatment of snake bites in southern part of Tamilnadu, *Indian Journal of Ethnopharmacology*. 2008; 115: 302-312.
14. Tang W & Eisenbrand G. Chinese drugs of plant origin, chemistry, pharmacology and use in traditional and modern medicine. Springer-Verlag. 1992; 97-103.
15. Chao WW & Lin BF. Isolation and identification of bioactive compounds in *Andrographis paniculata* Nees. (Chuanxinlian) *Chinese Medicine*. 2010; 5: 17.
16. Chopra R N, Nayar S L & Chopra IC. Glossary of Indian medicinal plants. NISCIR, CSIR, Delhi, 2002.
17. Official methods of analysis of AOAC International, 16th edition. 1995, AOAC International, Suite 400, 2200 Wilson Boulevard, Arlington, Virginia, USA.
18. World Health Organization, Quality control methods for medicinal plant materials. Published by WHO, Geneva, 1998.