

PROTECTIVE EFFECT OF *SOLANUM TORVUM* AGAINST TESTICULAR TOXICITY IN MALE WISTAR RATS

MAHALAXMI MOHAN*, SANGITA BHANDARE

*Department of Pharmacology, M. G. V's Pharmacy College, Panchavati, Nashik, Maharashtra 422003, Priyadarshini College of Pharmaceutical Sciences, Nampally, Chowdaryguda, Ghatkesar, Ranga Reddy District, Hyderabad 500088 India.
Email: mm_nasik@yahoo.co.in

Received: 29 Jan 2012, Revised and Accepted: 18 Mar 2012

ABSTRACT

Oxidative stress is the main factor in Doxorubicin (DXR) induced toxicity. Rats received either DXR (3 mg/kg, i.p.) on day 1, 7, 14, 21, 28 or *Solanum torvum* extract (100 mg/kg and 300 mg/kg, p.o.) prior to DXR or *S. torvum* (100 mg/kg and 300 mg/kg, p.o.) extract alone for 4 weeks. Testicular toxicity was assessed by recording changes in body weight, testis weight, epididymal sperm count and measuring the levels of marker enzymes-LDH, SGOT and CK-NAC; antioxidant defence enzymes SOD, CAT and GSH; membrane bound enzymes such as Na⁺ K⁺ ATPase, Ca²⁺ ATPase and Mg²⁺ ATPase; total proteins and histopathological changes. *S. torvum* (100 mg/kg and 300 mg/kg) significantly (p<0.05) reversed the changes induced by DXR and showed a lesser degree of cellular infiltration in histopathological studies. The results suggest that *S. torvum* has the potential in preventing the testicular toxicity induced by Doxorubicin.

Keywords: *S.torvum*, Doxorubicin, Testicular toxicity.

INTRODUCTION

Doxorubicin is well known anthracycline glycoside antibiotic that possesses a potent broad spectrum antitumour activity against a variety of cancers including severe leukemias, lymphomas, human solid tumours and haematological malignancies¹⁻³. However its use in chemotherapy has been limited largely due to its diverse toxicities, including kidney, renal, hematological and testicular toxicity⁴⁻⁶. Oxidative stress has been associated with DXR induced tissue damage⁷. Although the exact mechanism of DXR induced toxicity remains unknown, it is believed to be mediated through free radical formation, iron-dependent oxidative damage of biological macromolecules and membrane lipid peroxidation⁸. The production of free radicals as a byproduct of DXR metabolism is considered to be the primary mechanism of DXR induced toxicity, which can be ameliorated by the use of natural antioxidants⁹.

Solanum torvum Sw. (Solanaceae), commonly known, as Turkey berry is an erect spiny shrub of about 4 m tall, evergreen and widely branched. It is native and found cultivated in Africa and West Indies¹⁰. The fruits and leaves are widely used in Camerooninan folk medicine. *Solanum torvum* contains a number of potentially pharmacologically active chemicals like isoflavonoid sulfate and steroidal glycosides^{11,12}, chlorogenone and neochlorogenone¹³, triacontane derivatives^{14,15}, 22- β -O-spirostanol oligoglycosides¹⁶, 26-O- β -glucosidase¹⁷.

Several natural compounds with antioxidant activities are known to exhibit protection against DXR induced testicular toxicities. *Citrus paradisi*¹⁸, Grape fruit seed extract¹⁹ and melatonin²⁰ was found to protect against DXR induced testicular toxicity. The antioxidant properties of flavonoids and their ability to chelate free iron could be effective in reducing toxicity of DXR²¹. In view of this, though the antioxidant activity of *S.torvum* is well known, its protection against DXR- induced testicular damage is not reported. Therefore the present study was aimed at investigating the possible protective effect of *Solanum torvum*, against DXR-induced testicular toxicity in rats.

MATERIAL AND METHODS

Animals

Albino rats (Wistar strain) of either sex weighing between 200-250 g, were obtained from Serum Institute, Pune. Animals were housed into groups of five under standard laboratory conditions of temperature 25 \pm 1°C with free access to food (Amrut rat and mice feed, Sangli, India.) and water. The experiments were performed during the light portion. The experiments were carried out

according to the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), New Delhi, India, and approved by the Institutional Animal Ethical Committee.

Drugs and chemicals

Doxorubicin (ADRM, 50 mg) was purchased from Dabur Pharma Ltd. Baddi, India. ATP (Adenosine-5' triphosphate, disodium salt was purchased from Hi media, Mumbai). All the chemicals used were of analytical grade and chemicals required for sensitive biochemical assays were obtained from Merck, India. CK-NAC and SGOT kits were purchased from Aspen laboratories, Baddi, India. LDH kit was purchased from Reckon Diagnostics Pvt. Ltd. Vadodara, India. *Solanum torvum* extract was dissolved in distilled water and administered orally.

Preparation of extract

Dried fruits of *Solanum torvum* sw. (Solanaceae) were purchased from local market and authenticated by Dr. Dasari, from Ayurvedic Seva Sangh, Panchavati, Nashik, India. Mature fruits were collected, sun dried and grounded. The powder obtained (950gm) was defatted using petroleum ether (60-80°C). The marc was macerated in methanol for 3-4 days at room temperature. The filtrate was air dried and concentrated under reduced pressure to obtain 113 g, corresponding to a yield of 11.3 % w/w. Appropriate concentrations of the methanolic extract of *Solanum torvum* were made in distilled water. The chemical tests for phytoconstituents present in the crude extract such as flavonoids, alkaloids, tannins, saponins, were carried out²².

Experimental Protocol

The animals were divided into six groups each consisting of five rats and received the following treatment.

Group I (Control): Animals received distilled water for injection.

Group II (DXR): Animals received DXR injection (3 mg/kg, i.p.) on day 1, 7, 14, 21, 28.

Group III (DXR + ST): Animals received ST (100 mg/kg /day p.o. for 28 days) and DXR injection (3 mg/kg, i.p.) on day 1, 7, 14, 21, 28. Group IV (DXR + ST): Animals received ST (300 mg/kg /day, p.o. for 28 days) and DXR injection (3 mg/kg i.p.) on day 1, 7,14,21,28. Group V (ST): Animals received ST (100 mg/kg /day, p.o. for 28 days)

Group VI (ST): Animals received ST (300 mg/kg /day, p.o. for 28 days)

Estimation of body and organ weight

In each group, body weight of rats was taken before and after DXR treatment. The testes was weighed after keeping them in ice-cold saline.

Preparation of serum and tissue homogenate

After 48 hours of the last injection of either DXR or vehicle, blood was collected retro-orbitally from the inner canthus of the eye (under light ether anesthesia) using capillary tubes for serological analyses. Serum was separated using R-24 research centrifuge (Remi Instruments Ltd., Mumbai) at 3000 rpm for 15 min. and used for estimation of CK-NAC²³, LDH²⁴, SGOT²⁵.

The animals were sacrificed under euthanasia. Epididymis was removed, after the adhering tissues were cleared. The epididymal sperm count was done immediately. One of the testes was excised in chilled Tris buffer (10 mM pH 7.4) for measurement of tissue markers of oxidative stress and the other one was collected for histopathology. The excised testes were then weighed and homogenized in chilled Tris buffer (10 mM, pH 7.4) at a concentration of 10% (w/v). The homogenates were centrifuged at 10,000 rpm for 20 min using Remi centrifuge machine. The clear supernatant was used for the assays of lipid peroxidation (TBARS)²⁶; endogenous antioxidant enzymes, superoxide dismutase (SOD)²⁷, catalase (CAT)²⁸ and reduced glutathione (GSH)²⁹. The sediment after centrifugation of tissue homogenate were resuspended in ice cold Tris buffer (10mM, pH 7.4) to get a final concentration of 10% and was used for the estimation of different membrane bound enzymes such as Na⁺ K⁺ ATPase³⁰, Ca²⁺ ATPase³¹, Mg²⁺ ATPase³² and total proteins³³.

Epididymal sperm count

Epididymal sperm was collected by slicing the epididymis in 5 ml phosphate buffer saline (pH 7.2). An aliquot of the epididymal sperm suspension was used for spermatozoa count using Neubauer hemocytometer³⁴.

Histopathological studies

The testes were fixed in 10% formalin. The specimens were then processed for standard procedure and were embedded in paraffin

wax. The blocks were then sectioned according to hematoxylin and eosin method. Five-micrometer thick histological sections were obtained from the paraffin blocks. The sections were examined under the light microscope and photographs were taken under 10X using Moti camera.

Statistics

The mean \pm SEM values were calculated for each group. One-way ANOVA followed by Dunnett's multiple comparison tests were used for statistical analysis. Values of $p < 0.05$ was considered statistically significant.

RESULTS

Testis weight and sperm count

Control animals showed 1.46 ± 0.02 gm and 44.76 ± 1.79 millions/ml as isolated testes weight and sperm count. DXR treated group (3 mg/kg, i.p., on day 1, 7, 14, 21, 28) significantly ($p < 0.05$) decreased isolated testes weight and sperm count as compared to the control group. Co-treatment with methanolic extract of *Solanum torvum* (100 and 300 mg/kg/day, p.o.) in DXR treated (3 mg/kg, i.p. on day 1, 7, 14, 21, 28) animals significantly ($p < 0.05$) restored the isolated testes weight and sperm count as compared to DXR treated group (Table 1)

Antioxidant enzymes

The antioxidant level of SOD, CAT, GSH and TBARS enzyme of testes in control rats was found to be 0.23 ± 0.02 U/g, 7.58 ± 0.24 U/g, 42.77 ± 0.76 nM/mg and 2.07 ± 0.11 nM/mg of wet tissue. The SOD, CAT and GSH levels were significantly ($p < 0.05$) decreased and TBARS levels were significantly increased in testes tissue of DXR (3 mg/kg, i.p. on day 1, 7, 14, 21, 28) treated animals as compared to control group. Co-treatment with methanolic extract of *Solanum torvum* (100 and 300 mg/kg/day, p.o.) in DXR treated (3 mg/kg, i.p. on day 1, 7, 14, 21, 28) animals showed significant ($p < 0.05$) increase in SOD, CAT and GSH levels and significant decrease in TBARS levels as compared to DXR treated animals (Table 2).

Table 1: The Effect of methanolic extract of *Solanum torvum* on body weight, testes weight and sperm count in DXR treated animals.

S. No.	Treatment groups (mg/kg)	Final Body Weight (gm)	Absolute testes weight (gm)	Sperm count (millions/ml)
1	Control	239.4 \pm 2.34	1.46 \pm 0.02	44.76 \pm 1.79
2	DXR (3)	197.9 \pm 4.93*	0.89 \pm 0.01*	10.80 \pm 0.44*
3	ST (100) + DXR (3)	219.8 \pm 4.54#	1.09 \pm 0.02#	27.44 \pm 0.66#
4	ST (300) + DXR (3)	225.2 \pm 4.05#	1.15 \pm 0.06#	30.80 \pm 0.59#
5	ST (100)	211.4 \pm 5.65*	1.34 \pm 0.08	36.01 \pm 1.25*
6	ST (300)	220.6 \pm 5.56*	1.37 \pm 0.03	39.92 \pm 1.21*
7	F value	8.89	22.89	118.61

N = 5, All values are expressed as mean \pm SEM. One way ANOVA followed by Dunnett's test. * $P < 0.05$ against control group, # $P < 0.05$ against DXR group.

ST = methanolic extract of *Solanum torvum*, DXR = Doxorubicin.

Table 2: The Effect of methanolic extract of *Solanum torvum* on enzymes of oxidative stress in DXR treated animals.

S. No.	Treatment groups (mg/kg)	SOD (U/gm of wet tissue)	CAT (U/gm of wet tissue)	GSH (nm/mg of tissue)	TBARS (nm/mg of tissue)
1	Control	0.23 \pm 0.02	7.58 \pm 0.24	42.77 \pm 0.76	2.07 \pm 0.11
2	DXR (3)	0.15 \pm 0.01*	2.35 \pm 0.06*	16.43 \pm 1.33*	7.46 \pm 0.11*
3	ST (100) + DXR (3)	0.19 \pm 0.01#	3.22 \pm 0.08#	25.53 \pm 1.76#	5.79 \pm 0.24#
4	ST (300) + DXR (3)	0.23 \pm 0.01#	4.43 \pm 0.14#	34.85 \pm 0.79#	4.31 \pm 0.42#
5	ST (100)	0.19 \pm 0.01*	7.00 \pm 0.09*	34.68 \pm 0.92*	3.51 \pm 0.16*
6	ST (300)	0.27 \pm 0.01*	6.99 \pm 0.22*	37.82 \pm 0.39*	3.04 \pm 0.08*
7	F value	16.0	210.40	76.32	80.08

N = 5, All values are expressed as mean \pm SEM. One way ANOVA followed by Dunnett's test. * $P < 0.05$ against control group, # $P < 0.05$ against DXR group.

ST = methanolic extract of *Solanum torvum*, DXR = Doxorubicin.

Marker enzymes

Control animals showed (12.82 ± 0.56IU/L), (11.21 ± 0.47IU/L) and (4.81 ± 0.38 IU/L) as serum marker enzyme level of LDH, CK-NAC and SGOT respectively. DXR treated group (3 mg/kg, i.p., on day 1, 7, 14, 21, 28) significantly (p<0.05) increased the levels of marker

enzymes as compared to the control group. Co-treatment with methanolic extract of *Solanum torvum* (100 and 300 mg/kg/day, p.o.) in DXR treated (3 mg/kg, i.p., on day 1, 7, 14, 21, 28) animals showed significant (p<0.05) decrease in levels of marker enzymes as compared to the DXR treated group (Figure 1).




Fig. 1: The Effect of ST extract (100, 300 mg/kg/day, p.o., for 4 weeks) on biomarkers enzyme in DXR treated male rats.

All values are expressed as mean ± SEM, n=5. All data are subjected to One Way ANOVA followed by Dunnett’s test. * p<0.05 when compared to control and # p<0.05 when compared to DXR group. Vertical lines represent S.E.M. ST-*Solanum torvum*, DXR-Doxorubicin.

Membrane bound enzymes

The membrane bound enzyme level of Na⁺ - K⁺ ATPase, Ca²⁺ - ATPase and Mg²⁺ - ATPase of testes in control rats was found to be 6.88 ± 0.07, 4.33 ± 0.08 and 7.03 ± 0.26 μmoles of inorganic phosphorus liberated/min/mg protein. These membrane bound enzyme levels were significantly (p<0.05) decreased in testes tissue of DXR (3

mg/kg, i.p. on day 1, 7, 14, 21, 28) treated animals when compared to control group. Cotreatment with methanolic extract of *Solanum torvum* (100 and 300 mg/kg/day, p.o.) for 4 weeks in DXR treated (3 mg/kg, i.p. on day 1, 7, 14, 21, 28) animals showed significant (p<0.05) increase in membrane bound enzyme levels level in testes tissue as compared to DXR treated animals (Figure 2).




Fig. 2: The Effect of ST extract (100, 300 mg/kg/day, p.o., for 4 weeks) on membrane bound enzyme in DXR treated male rats.

All values are expressed as mean ± SEM, n=5. All data are subjected to One Way ANOVA followed by Dunnett’s test. * p<0.05 when compared to control and # p<0.05 when compared to DXR group. Vertical lines represent S.E.M. ST-*Solanum torvum*, DXR-Doxorubicin.

Histopathology

Control group treated with vehicle showed normal architecture, normal feature of seminiferous tubules and interstitial tissue. Spermatocytes, sertoli cells, mature sperms are were seen. Group treated with DXR (3 mg/kg, i.p.on day 1, 7, 14, 21, 28) showed degenerative changes in the seminiferous tubules, the polarity of

spermatocyte and basement membrane were disturbed. No mature sperms were present and the normal architecture of cells was disturbed.

Cotreatment with methanolic extract of *Solanum torvum* (100 and 300 mg/kg/day, p.o.) for 4 weeks in DXR treated (3 mg/kg, i.p. on day 1, 7, 14, 21, 28) animals showed near normal architecture (Figure 3).




Fig. 3: The Photomicrographs of histopathological examination (10 X) of testes tissue.

A) Control group treated with vehicle shows normal architecture, normal feature of seminiferous tubule and interstitial tissue. Spermatocytes, sertoli cell, mature sperms are present.

B) Group treated with DXR (3 mg/kg) shows degenerative changes in the seminiferous tubule. It disturbs the polarity of spermatocyte. Distruption of basement membrane. No mature sperms are present. Normal architecture of cell gets disturb.

C) Group treated with ST (100mg/kg) and ST (300mg/kg) shows normal architecture, shows normal feature of seminiferous tubule and interstitial tissue. Spermatocytes, sertoli cell, mature sperms are present similar to control group.

D) Group treated with ST (100mg/kg) and DXR 3 mg/kg) and ST (300mg/kg) and DXR (3 mg/kg) shows significant increase in mature sperms, try to gain the basement membrane as compare to DXR, restored these changes towards normalcy.

DISCUSSION

Doxorubicin a widely used anticancer drug is associated with multiple organ toxicity. A strategy to preserve the anticancer properties with minimum side effects is desired. Doxorubicin is known to disturb spermatogenesis in a dose dependent manner in animal studies [6]. Ward and his co-workers have also reported doxorubicin induced reductions in testicular sperm count [35]. Administration of DXR to rats significantly increased epididimal sperm count, TBARS levels, and levels of LDH, SGOT and CK-NAC; and significantly decreased testes weight, antioxidant defence enzymes SOD, CAT, GSH; membrane bound enzymes such as Na⁺ K⁺ ATPase, Ca²⁺ ATPase, Mg²⁺ ATPase; and total proteins. Our results showed that DXR induced biochemical and histopathological

changes which are indicative of toxicity are in good agreement with those previously reported [20, 36]. However treatment with *S. torvum* extract (100 mg/kg and 300 mg/kg) has resulted in significant ($p < 0.05$) change in all parameters as observed in DXR treated animals thus offering considerable protection against testicular toxicity. Free radical scavenging enzymes such as catalase, superoxide dismutase are the first line cellular defense enzymes against oxidative injury, decomposing O₂ and H₂O₂ before their interaction to form the more reactive hydroxyl radical (OH·). The equilibrium between these enzymes is an important process for the effective removal of oxygen stress in intracellular organelles [37]. In our study, a decrease in concentration of antioxidant levels in DXR treated group was observed. Treatment with *S. torvum* extract (100 mg/kg and 300 mg/kg) significantly increased the antioxidant levels. There is a general agreement that flavonoids act as scavengers of reactive oxygen species [38]. The antioxidant properties of *Solanum torvum* could be attributed to the presence of flavonoid phytoconstituent in it. Elevated serum levels of CK-NAC, LDH and SGOT are indicative of toxicity as seen in DXR group, which was ameliorated by concurrent treatment with *S. torvum* extract. The restoration of membrane bound enzymes like in *S. torvum* treated rats was due to the membrane stabilizing protective effect of *S. torvum*. Moreover, the histopathology data has revealed that treatment with *S. torvum* has protected the testis from degenerative changes in the seminiferous tubule, disruption of basement membrane and normal architecture induced by DXR. Thus, this study revealed that pretreatment with *Solanum torvum* protected against DXR induced testicular toxicity.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

ACKNOWLEDGEMENT

The authors acknowledge the help provided by Dr. Rajendra Thigale, Ashirvad Pathology Laboratory, Nashik; Mr. Pradeep for the technical assistance provided during histopath study and Dr. Meena Kulkarni, HOD, Dept of histopathology, KBH Dental College, Nashik.

REFERENCES

- Priestman T. 2008. Cancer Chemotherapy in Clinical Practice. Springer- Verlag, London, UK, pp.130-136.
- Blum RH, Carter SK. A new drug with significant clinical activity. Ann. Intern. Med. 1974; 80: 249-256.
- Calabresi P, Chabner BA. Chemotherapy of neoplastic diseases, In: Gilman AG, Rall TW, Nies AS, Taylor P. (eds.), The Pharmacological Basis of Therapeutics. NY: Pergamon Press Inc. 1990; 1203-1263.
- Gillick J, Giles S, Bannigan J, Puri P. Cell death in the early adriamycin rat model. Pediatr. Surg. Int. 2002; 18: 576-580.
- Yilmaz S, Atessahin A, Sahna E, Karahan I, Ozer S. Protective effect of lycopene on adriamycin- induced nephrotoxicity. Toxicology 2006; 218: 164-171.
- Lui RC, Laregina MC, Herbold DR, Johnson FE. Testicular cytotoxicity of intravenous doxorubicin in rats. J. Urol. 1986; 136: 940-943.
- Fadilloglue E, Oztas E, Erdogan H, Yagmurca M, Sogut S, Ucar M, Irmac MK. Protective effect of caffeic acid phenyl ethyl ester of doxorubicin induced nephrotoxicity in rats. J Appl Toxicol. 2004; 24: 47-52.
- Pritsos CA and Ma J. Basal and drug-induced antioxidant enzyme activities correlate with age-dependent doxorubicin oxidative toxicity. Chem. Biol. Interact. 2000; 127: 1-11.
- Quiles J, Huertas J, Battino M, Mataix J, Ramirez-Tortosa, M. Antioxidant nutrients and adriamycin toxicity. Toxicology.2002; 180:79-95.
- Adjanohoun JE, Aboubakar N, Dramane K, Ebot ME, Ekper JA, Enoworock EG, Foncho D, Gbile ZO, Kamanyi A, Kamoukom Jr, Keeta A, Mbenkum T, Mbi CM, Mbielle AL, Mbome IL, Mubiru NK, Naney WL, Nkongmeneck B, Satabie B, Sofowa A, Tanze V, Wirmum CK. Traditional medicine and pharmacopeia-contribution to ethnobotanical and floristic studies in Cameroon. In: CNPMS. Porto-Novo, Benin; 1996, 50 -52.
- Yahara S, Yamashita T, Nozawa N, Nohara T. Steroidal glycosides from *Solanum torvum*. Phytochemistry. 1996; 43, 1069-1074.
- Arthan D, Svasti J, Kittakoop P, Pittayakhachonwut D, Tanticharoen M,
- Thebtaranonth Y. Antiviral isoflavonoid sulfate and steroidal glycosides from the fruits of *Solanum torvum*. Phytochemistry. 2002; 59: 459-463.
- Carabot CA, Blunden G, Patel VA. Chlorogenone and neochlorogenone from the Unripe fruits of *Solanum torvum*. Phytochemistry. 1991; 30: 1339- 1341.
- Mahmood U, Shukla, YN, Thakur, RS. Non-alkaloidal constituents from *Solanum torvum* leaves. Phytochemistry. 22, 1983; 167-170.
- Mahmood U, Agrawal PK, Thakur RS. Torvonin-A, a spirostane Saponin from *Solanum torvum* leaves. Phytochemistry. 1985; 24: 2456-2457.
- Iida Y, Yanai Y, Ono M, Ikeda T, Nohara T. Three unusual 22- β -O-23-hydroxy-(5 α)-spirostanol glycosides from the fruits of *Solanum torvum*. Chemical and Pharmaceutical Bulletin. 2005; 53: 1122- 1125.
- Arthan D, Kittakoop P, Esen A, Svasti J. Furostanol glycoside 26-O- β glucosidase from the leaves of *Solanum torvum*. Phytochemistry. 2006; 67: 27- 33.
- Saalu LC, Osinubi AA, Jewo PI, Jewo AO, Oyewopo GO. An evaluation of influence of Citrus paradise seed extract on doxorubicin induced testicular oxidative stress and impaired spermatogenesis. Asian J. Sci. Res. 2010a; 1-11.
- Saalu LC, Osinubi AA, Oguntola JA, Adeneye IO, Benebo AS. The delayed testicular morphologic effects of doxorubicin and the rejuvenating role of grapefruit seed extract. Int. J. Pharmacol. 2010b; 6 (3): 192-199.
- Patil L, Balaraman R. Effect of melatonin on doxorubicin induced testicular damage in rats. Int. J. Pharm Tech Res. 2009; 1(3): 879-884.
- Vaclavikova R, Kondrova E, Ehrlichova D M, Boumendjel A, Kovar J, Stopka, P,et al. The effect of flavonoid derivatives on doxorubicin transport and metabolism. Bioorg. Med. Chem. 2008; 16: 2034- 2042.
- Kokate CK. In: Practical Pharmacognosy, 3rd Edn., Vallabh Prakashan, New Delhi; 1994; pp.107-109.
- Szasz G, Gruber W, Bernst E. Creatine kinase in serum: 1. Determination of optimum reaction conditions. Clin. Chem. 1976; 22: 650-656.
- Gay RJ, Comb RB, Bowers GN. Optimum reaction condition for human lactate dehydrogenase isoenzyme as they affect total lactate dehydrogenase activity. Clin. Chem.1968; 14: 740 -753.
- Karmen A, Wrblewski F, Ladue JS. Transaminase activity in human blood. J. Clin. Invest. 1955; 34: 126.
- Niehaus WG, Samuelsson B. Formation of malondialdehyde from phospholipids arachidonate during microsomal lipid peroxidation. Eur. J. of Biochem. 6, 1968; 126-130.
- Saggu H, Cooksey J, Dexter DA. Selective increase in particulate Superoxide dismutase activity in Parkinsonian substantia nigra. J. Neurochem. 1989; 53: 692- 697.
- Beers JR, Sizer IW. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. J Biol Chem. 1952; 115:133-140.
- Ellman GL. Tissue sulphhydryl groups. Archives of Biochemistry and Biophysics 1959; 82: 70-77.
- Bonting SL. Presence of enzyme system in mammalian tissues: Membrane and Ion Transport, Wiley Inter Science, London; 1970: 257- 263.
- Hjerten S, Pan, H. Purification and characterization of two forms of a low affinity Ca²⁺-ATPase from erythrocyte membranes. Biochimica et Biophysica ACTA.1983; 281-288.
- Ohnishi T, Suzuki T, Suzuki Y, Ozawa K. A comparative study of plasmamembrane Mg²⁺-ATPase activities in normal, regenerating and malignant cells. Biochimica et Biophysica ACTA. 1982; 684, 67-74.
- Lowry OH, Rosenbrough NJ, Farr AC, Randell RJ. Protein measurement with folinphenol reagent. J. Biol. Chem.1975; 193: 265- 275.
- Narayana K, D'Souza UJ, Rao KP. Effect of ribavirin on epididymal spermcount in rat. Indian J. Physiol Pharmacol. 2002; 46: 97-101.
- Ward JA, Bardin CW, Knight M, Robinson J, Gunsalus G, Morris ID. Delayed effects of doxorubicin on spermatogenesis and endocrine function in rats. Reprod. Toxicol. 1988; 2:117-126.
- Yeh YC, Hui CL, Chih TT, Lieng LW, Chuan LW, Yang WK, Chun HL. Protection by doxycyclin against doxorubicin - induced oxidative stress and apoptosis in mouse testes.2007; 74: 969-980.
- Singal PK, Li T, Kumar D, Danelisen I, Hiskovic, N. Adriamycin-induced heart failure: mechanism and modulation. Mol. Cell. Biochem. 2000; 207: 77-86.
- Haenen MM, Janssen FP, Bast A. The antioxidant properties of five (O- β hydroxyethyl) rutosides of the flavonoid mixture Venoruton. Phlebol Suppl.1993; 1: 10-17.