

CHEMICAL ANALYSIS OF PRIMARY AND SECONDARY METABOLITES OF *NARAVELIA ZEYLANICA*LALITHA EASWARAN¹ AND V. ALEX RAMANI²¹Department of Chemistry, MAM College of Engineering, Siruganur, Tiruchirapalli, India. ²Department of Chemistry, St. Joseph's College (Autonomous), Tiruchirapalli, India. Email: lalithaeaswaran71@yahoo.com

Received: 18 Feb 2012, Revised and Accepted: 30 Mar 2012

ABSTRACT

Minerals are very essential nutrient for human life. The present study deals with the study of phytoconstituents from the leaves of *Naravelia zeylanica*. The plant *Naravelia zeylanica* was analyzed quantitatively for the primary metabolites -moisture (65%), protein (16.6%), fat (2.2%), fiber (24.4 %), carbohydrate (8.8%), vitamin A(0.125ppm), vitamin B (0.648ppm) and vitamin C (0.263ppm).The macro elements such as nitrogen, phosphorus, potassium, calcium and magnesium constitute 16.04% and microelements such as sodium, sulphur, zinc, copper, iron, manganese, boron and molybdenum constitute 98.84ppm. The secondary metabolites such as alkaloid, flavonoid, tannin and glycoside constitute 2.23ppm and trace amount of serpentine, terpenoid, saponin and phenol were estimated. The nutritive value of the plant is 121.4 cal/100g.

Keywords: Metabolites, *Naravelia zeylanica*

INTRODUCTION

Naravelia zeylanica (Ranunculaceae) is a genus of woody climbers distributed in Himalayas. The crushed stem was used as a cure for rhinitis¹. The crushed roots were found to relieve headache². The antimicrobial activity of bioactive constituents from the leaves of *Naravelia zeylanica* was studied³. The micropropagation of the plant, isolation and characterization of berberine⁴ from leaves of *Naravelia zeylanica* were done systematically. Antilucer activity of the leaf extract of *Naravelia zeylanica* was reported⁵. The quantitative estimation antioxidant studies on aerial parts of *Naravelia zeylanica* were reported⁶. The proximate analysis of *Dendrobium macrostachyum lindl* was reported⁷. The present study aims at the quantitative analysis of the primary and secondary metabolites of *Naravelia zeylanica*.

MATERIALS AND METHODS

The fresh plant samples were obtained locally from the Kolli Hills, Trichy. The plant species was verified with authentic specimen at Rapinat Herbarium, Trichy, Tamilnadu, India. The leaves were washed in tap water; shade dried, crushed and was taken for various phytochemical analyses.

The moisture content was determined by taking the fresh plant samples in petridish and kept overnight in an air oven at 100-110°C till they attain a constant weight. The loss in weight was regarded as a measure of moisture content⁸.

$$\text{Percentage moisture} = \frac{\text{Loss in weight of sample}}{\text{Weight of sample taken}} \times 100$$

The plant sample was hydrolyzed with dilute hydrochloric acid and subjected for spectrophotometric determination of total carbohydrate content using anthrone reagent at 630nm⁹. The nitrogen content was estimated by Kjeldahl method¹⁰. The protein content was estimated spectrophotometrically by Lowry's method¹¹ using Folin - Ciocalteu reagent at 660nm. The lipid was determined by direct weight method¹². For determination of crude fibre, the plant sample (free from moisture and fat) was treated with 1.25% NaOH, filtered, washed with hot water and then treated with 1.25% H₂SO₄. The dry residue obtained at 130°C, was ashed at 600°C. Crude fiber¹³ was calculated from the masses of dry residue and the ash content.

$$\text{Crude fiber \%} = \frac{\text{Weight of dry residue} - \text{weight of ash}}{\text{Weight of sample}} \times 100$$

The nutritive value was determined by the formulae:

$$\text{Nutritive value} = 4 (\% \text{protein}) + 9 (\% \text{fat}) + 4 (\% \text{Carbohydrate})$$

The total tannin content was estimated by treating the sample with Folin - Denis reagent¹⁴. Plant ash was used for quantitative

estimation of minerals. The amounts of phosphorus, potassium, sodium, zinc, copper, iron and manganese were estimated by flame photometric method¹⁵. Calcium and magnesium were estimated by EDTA method. The minerals - sulfur, boron and molybdenum were analyzed by using spectrophotometer. Alkaloid was determined by the method of Harborne¹⁶, plant sample was treated with acetic acid, ethanol and concentrated ammonium hydroxide was added for complete precipitation. The residue obtained was alkaloid, which was dried and weighed. Flavonoid was estimated by the method of Bohm and Kocipai-Abyazan¹⁷. Saponin was estimated following the method of Harborne. Phenol was estimated spectrophotometrically at 505 nm¹⁸. Estimation of vitamin A was done by spectrophotometer at 620nm using trichloroacetic acid and vitamin B was estimated using spectrophotometer at 540 nm using ferric sulphate and potassium thiocyanate solution. Estimation of vitamin C was done by Iodometric titration method.

RESULTS AND DISCUSSION

The moisture content in the leaves of *Naravelia zeylanica* is 65%. The leaves contained higher percentage of protein 16.6%. High ash content indicated that the leaves were good source of inorganic minerals. A low fat value (2.2%) was obtained in study confirms that the leaf is not a good source of oil.

Potassium was 4.16% in the leaves of *Naravelia zeylanica*. It is necessary for the formation of sugars, starch, carbohydrates, protein synthesis and cell division in all parts of the plant¹⁹. Calcium in the leaves of *Naravelia zeylanica* was about 4.56%. Calcium forms the structural component of cell walls, activates enzymes, and influences water movement in cell. Calcium is necessary for cell growth and cell division, bone formation and blood coagulation. Magnesium was the highest in percentage (5.19%) of all the macro elements. It is critical structural component of chlorophyll molecule and is necessary for the functioning of plant enzymes in the production of carbohydrates, sugars and fat²⁰.

The results of various chemical constituents are presented in Table 1.

Phosphorus was 0.87%, it is necessary for photosynthesis. It plays a vital role in energy metabolism in the formation of sugar phosphates and adenosine di- and tri-phosphates. Iron is an essential element for synthesis of hemoglobin. The quantity of iron is the highest (68.54ppm) of all the micronutrients. Next to iron, manganese is present in about 26.36ppm. Manganese is necessary for the functioning of the pituitary gland and the brain. Zinc and manganese are considered as antioxidant micro nutrients and their presence could therefore boost the immune system²¹. All the other micronutrients such as sodium, sulphur, copper, boron and molybdenum were present in trace amount. The nutritive value of the plant was found to be 121.4 cal/100g.

Flavonoid was present in considerable amount that is 0.87 ppm. Tannin and alkaloid were present in moderate amounts of 0.54 and 0.41ppm respectively. Presence of tannins possesses astringent properties. Alkaloids are reported to have analgesic and anti-inflammatory activities which help to alleviate pains, develop resistance against diseases and endurance against stress. Saponin, phenol and terpenoid were present in trace amount. Saponin is very

useful in the management of upper respiratory tract inflammation and cardio tonic in nature. *Naravelia zeylanica* leaves contain higher amount of vitamin B (0.648 ppm) compared to vitamin A (0.125 ppm) and vitamin C (0.263ppm). Vitamin C protects the vitamin B from oxidation²². The present study revealed the presence of various primary and secondary metabolites in the leaves of *Naravelia zeylanica*.

Table 1: Chemical constituents of *Naravelia zeylanica* leaves

S. No.	Phytoconstituents	Quantity	
		%	ppm
	Primary metabolites		
1	Moisture	65	
2	Protein	16.6	
3	Fat	2.2	
4	Fiber	24.4	
5	Carbohydrate	8.8	
6	Vitamin A		0.125
7	Vitamin B		0.648
8	Vitamin C		0.263
	Macroelements		
9	Nitrogen	1.26	
10	Phosphorus	0.87	
11	Potassium	4.16	
12	Calcium	4.56	
13	Magnesium	5.19	
	Microelements		
14	Sodium		0.12
15	Sulphur		0.52
16	Zinc		2.89
17	Copper		0.23
18	Iron		68.54
19	Manganese		26.36
20	Boron		0.12
21	Molybdenum		0.06
	Secondary metabolites		
22	alkaloid		0.41
23	flavonoid		0.87
24	tannin		0.54
25	glycoside		0.39
26	serpentine		0.12
27	terpenoid		0.06
28	saponin		0.02
29	phenol		0.13

REFERENCES

1. Arun Vijayan VB, Liju JV, Reena John, Parthipan B and Renuka C. Traditional remedies of Kani tribes of Kothoor reserve forest, Agasthayavanam, Thiruvananthapuram, Kerala. Indian Journal of Traditional Knowledge. 2007.6(4): 589-594.
2. The Wealth of India. A Dictionary of Indian Raw Materials and Industrial products. Vol VII (N-Pe), Publication and Information Directorate, Council of Scientific and Industrial Research, New Delhi. 1992.
3. Raja Naika H, Krishna V, Khadeer Ahamed BM and Mahadevan KM. Antimicrobial activity of bioactive constituents isolated from the leaves of *Naravelia zeylanica*. International journal of biomedical and pharmaceutical sciences.2007. 1(2):153-159.
4. Raja Naika H and Krishna V. Micro propagation, Isolation and Characterization of Berberine from leaves of *Naravelia zeylanica*. Research Journal of Medicinal Plant. 2008. 2(1):1-9.
5. Ashoka Shenoy M, Shastry CS, Sridevi and Gopakumar P. Antiulcer activity of *Naravelia zeylanica* leaves extract. Journal of Pharmacy Research. 2009 2(7): 1218- 1220.
6. Sutharsingh R, Kavimani S, Jayakar B, Uvarani M and Thangathirupathi A. Quantitative phytochemical estimation and antioxidant studies on aerial parts of *Naravelia zeylanica*. International Journal of Pharmaceutical Studies and research. 2011. 2(2): 52-56.
7. Nimisha PS and Hiranmai Yadav R. Proximate and physicochemical analysis of *Dendrobium macrostachyum lindl*. International Journal of pharmacy and pharmaceutical sciences.2012.4 (1) : 385-386.
8. Indrayan AK, Sharma S, Durgapal D, Kumar N and Kumar M . Determination of nutritive value and analysis of mineral elements for some medicinally valued plants from Uttaranchal. Current Science. 2005. 89(7): 1252-1255.
9. Sadasivam S and Manickam A. Biochemical methods for agricultural Sciences. New Age International (P) Ltd, New Delhi. 1996 :1-97.
10. AOAC Official method of analysis. Association of Official Agricultural Chemicals, 10th Edn., Washington, DC. 1985.
11. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ. Protein measurement with folin phenol reagent. J Biol Chem. 1951.193: 265-275.
12. Aparna Buzarbarua. A Textbook of Practical Plant Chemistry. S. Chand and Company Ltd. 2000.
13. Unni BG, Kakoty Ajit C, Khanikor D, Bhattacharya PR, Pathak MG, Pillai K, Pillai R, Choudhury A, Saikia PC and Ghosh AC. Lipid and fatty acid composition of muga silkworm, *Antheraea assama*, host plants in relation to silkworm growth. J.Lipid Mediators and Cell signaling.1996. 13(3): 295-300.
14. Schanderl SH. Method in Food Analysis. Academic Press, New York.1970: 709.

15. Thimmaiah SK. Standard method of Biochemical analysis, Kalyani Publishers 1999:449-446.
16. Harborne JB. Phytochemical methods, London, Chapman and Hall Ltd. 1973 : 49-188.
17. Bohm BA and Kocipai-Abyazan R. Flavonoids and condensed Tannins from leaves of Hawaiian *Vaccinium vaticulatum* and *V. calycinium*. Pacific Science. 1974.48: 458-463.
18. Krishnaiah D, Devi T, Bano A and Sarbatly R. Studies on phytochemical constituents of six Malaysian medicinal plants. J. Medicinal Pl. Research. 2009. 3(2): 67-72.
19. Watts D. Trace elements and other essential nutrients, clinical application of tissue mineral analysis. Writer's B-L-O-C-K Edn., United States of America. 1997.
20. Kitahara N, Shibata S and Nishidha T. Fao electronic conference: Mulberry for animal production. 2000.
21. De AC. Fundamentals of Biochemistry. A Sen new central book agency, Calcutta, India 1990:431 - 437.
22. Jimoh FO and Oladiji AT. Preliminary Studies on *Piliostigma thonningii* seeds: Proximate analysis, mineral composition and phytochemical screening. African J. Biotechnology. 2005. 4:1439-1442.