

A STUDY OF THE BEHAVIOUR OF L - GLUTAMIC ACID IN THE COURSE OF AND AFTER γ - RAY TREATMENT

DANKA OBRESHKOVA¹, DOBRINA TSVETKOVA¹, IVANKA PENCHEVA¹, LUCIANO SASO², KALIN IVANOV³

¹Medical University Sofia, Faculty of Pharmacy, Department of Pharmaceutical Chemistry 2 Dunav str., Sofia 1000, Bulgaria, ²Sapienza University Rome, Department of Physiology and Pharmacology "Vittorio Erspamer", Italy, ³Medical University Plovdiv, Faculty of Pharmacy. Email: dobrinkadoncheva30@yahoo.com

Received: 13 Feb 2012, Revised and Accepted: 21 Mar 2012

ABSTRACT

The aim of current study is quality control of L - Glutamic acid in supplement mixture before and after treatment with γ - ray. Microbiological methods, included in European Pharmacopoeia were used for examination of microbial purity of substance L - Glutamic acid. Abnormal content ($1.5 \cdot 10^{-4}$ g) of bacteria and contaminants were identified mostly as non pathogenic bacilli of Subtilis group. Pathogenic contaminants as Enterobacteriaceae and Staphylococcus aureus were not found. Resistency factors show moderate ray sensitivity of the microorganisms.

HPLC method was developed and applied and analytical parameters repeatability, limit of detection (LOD), limit of quantitation (LOQ) and linearity were studied and determined in accordance with ICH and European Pharmacopoeia requirements. For repeatability SD = 1.43, RDS = ± 0.44 . The obtained LOD is 10 μ g and LOQ is 40 μ g. The correlation coefficient is found to be 0.99746 at SD = ± 3914.60 AU. There are no significant difference between content of L - Glutamic acid in supplement mixtures before (RDS = ± 0.44 %) and after γ - ray treatment (RDS = ± 0.082 %).

Keywords: L-Glutamic acid.

INTRODUCTION

Proteins and peptides are polymers of α - aminoacids. Aminoacids can be classified on lots of different features. According to the fact, whether or not human can acquire them through the diet, are recognized 3 types: nonessential, essential and conditionally essential aminoacids. Nonessential are produced by the human body either out of the essential or from normal proteins breakdown. Nonessential aminoacids include L - Alanine, L - Arginine, L - Aspartic acid, Asparagine, L - Cysteine, L - Glutamic acid, Glutamine, L - Glycine, L - Proline, L - Serine, L - Tyrosine. L - Glutamic acid (2 - Aminopentanedioic acid) (Fig. 1) is one of the most common nonessential aminoacids and is a major excitatory neurotransmitter in the human brain and in the spinal cord. L - Glutamic acid is necessary for proper cell functioning, but is considered as a nonessential aminoacid, because human body is able to produce it. Being one of the few nutrients able to pass through the blood - brain barrier, L - Glutamic acid supports brain function. L - Glutamic acid has the ability to detoxify brain and muscle cells by transforming all excess ammonia into the aminoacid Glutamine, which has antioxidant properties. As a chemical messenger in human brain, L - Glutamic acid is able to enhance a clarity of thinking, mental alertness, mood and intelligence and is applied to help for treatment of Parkinson's, fatigue, mental retardation, schizophrenia, muscular dystrophy and alcoholism. L - Glutamic acid is acting as an intermediary in the Krebs' cycle 1.




Fig. 1: Structure of L Glutamic acid.

There are different types of radiation - UV, γ . UV - radiation can be a health risk on the population 2 because causes sunburns, ageing of the skin and skin cancer. Sunscreens and sunblocks, included in skin care products, reduce UV - B - generated ROS 3. The effect γ - rays on survivor, morphological variation and chlorophyll mutation of some plants (*Abelmoschus Moschatus*) is also studied 4.

A crucial step in pharmaceutical production is sterilization. For sterilization (S) have been developed the following methods: dry heat S, pressured vapor S, ethylene oxide (EtO) S, formaldehyde S,

gas plasma (H_2O_2) S, peracetic acid S, γ - radiation S and E - beam S. Each technique has aspects that make it suitable or unsuitable for the sterilization of a particular product 5.

Radiation sterilization of medical products is regulated by the following standards: EN 552 [6]; ISO 11137 [7]; ISO 11737 [8]; ISO 14937 [9].

In comparison with other methods for producing sterile products, the advantages of γ - radiation sterilization are:

- 1) better assurance of product sterility than filtration and aseptic processing
- 2) low - temperature process - preserving properties of materials
- 3) no residues (like EtO) or no radioactivity remain in the products
- 4) high penetrating power than E - beam.
- 5) simple validation process - only one process variable (exposure time or dose) needs to be controlled. Sterilization by EtO needs seven variables (temperature, time, pressure, vacuum, gas concentration, packaging and humidity) and steam sterilization needs six variables (temperature, time, pressure, vacuum, packaging and humidity) to be controlled 10.

One of the most important advantage of γ - radiation in sterilization of thermolabile drugs is the high penetrating power 11. On the other hand, γ - radiation can cause degradation and changing of physicochemical properties and therapeutic effect of drugs 12.

The first aspect to consider when sterilizing with γ - radiation is product tolerance to the radiation. During use of this type of radiation, high - energy photons bombard the product, causing electron displacement within. These reactions generate free radicals, which aid in breaking chemical bonds. Disrupting microbial DNA renders any organisms that survive the process nonviable or unable to reproduce. These high - energy reactions also have the potential to disrupt bonds within the pharmaceutical formulation, to weaken the strength of packaging materials and to cause changes in color or odor in some materials. For these reasons, drug manufacturers should perform prequalification Dmax (maximum dose) testing, whereby the drug and it's packaging are subjected to a high dose of γ - radiation and then evaluated for stability and functionality 10.

The radiation resistance of a microorganism is measured by the decimal reduction dose (D_{10} value), which is defined as the radiation

dose (kGy), required to kill 90 % of the total number of microorganisms 13. The study of microbiological purity is important not only for drugs, but also for medicinal plants [14] and cosmetic products 15.

Parameters to characterize include potency, efficacy, stability, biocompatibility and chemical acceptability. Per guidelines under the International Conference on Harmonization (ICH), known as Technical Requirements for Registration of Pharmaceuticals for Human Use, it is recommended to use high – performance liquid chromatography (HPLC), mass spectrometry or gas chromatography to characterize and compare different analytical aspects of irradiated product versus nonirradiated product 16.

The use of food additives is not a modern – day invention, but their excessive and uncontrolled use increases each year. Food additives consumed in excessive amounts will be toxic. The process called risk assessment is connected with determination of the safe levels of chemicals in food by review of all information for the types of harmful effect of chemicals. The risk assessment is conducted by independent scientific advisory committees 14.

The contemporary requirements for microbial purity of aminoacids substances, when been produced in non – sterile environment, set the problems of the microbial determination of the product, as well as of the initial materials. The results of the routine tests show that very often the materials contain impurities over the required limits. γ – ray treatment may usefully reduce level of impurities in aminoacids used for the production of food supplements.

The sensitivity of different drugs to radiation has been studied: antibiotics [17]: Cefoperazone 18, Cefotaxime 19, Cefotaxidime [20]; glycosides [21]; steroids: Hydrocortisone 22, Prednisolone [23]; anticancer drugs: Doxorubicin [24, 25]; Cyclophosphamide [26]; alkaloids [17]; antiemetic drugs [26]; Ketoprofen [27]; Metoprolol tartrate [28]; Metro – nidazole [29]; Salbutamol [30]; Sulfacetamide sodium [31]; Clinopodium vulgare L. 32.

For determination of L – Glutamic acid are described the following methods: I) HPLC with: 1) post – column derivatization with o – phthalaldehyde [33]; 2) ion – pairing reversed – phase with UV and evaporative light scattering detection [34]; 3) isocratic reversed – phase with fluorimetric detection [35, 36]; 3) mass spectrometry MS [37]; II) gas chromatography [38]; III) polarography [39]; IV) amperometry 40.

The aim of current study is the development and application of HPLC method for quality control of L – Glutamic acid in supplement mixture before and after treatment with γ – ray and the examination of microbial purity of substance L – Glutamic acid.

MATERIALS AND METHODS

Materials

Materials for microbiological methods

1. L – Glutamic acid substance.
2. Microorganisms: *Bacillus subtilis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*.
3. Buffered sodium chloride – peptone solution with pH = 7.

Materials for HPLC method

L – Glutamic acid substance, methanol, water

Methods

Microbiological methods

The following included in European Pharmacopoeia methods for determination of microbial contamination are used: 1) Method 2.6.12. Total viable aerobic count (for detection of bacteria: *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Candida albicans*, *Aspergillus niger*); 2) Method 2.6.13. Tests for specified microorganism (for detection of bacteria, belonging to the family of Enterobacteriaceae) 41.

Preparation of test strain of microorganisms

Plates with casein soya bean digest agar were inoculated with 100 colony forming units respectively with the following microorganisms: *Bacillus subtilis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*. Plates were incubated for 18 – 24 h at 30 – 35 °C.

Sample preparation

10 g of L – Glutamic acid substance were dissolved in 10 ml buffered sodium chloride – peptone solution with pH = 7. Sample was filtered through membrane filter with pore size 0.45 μ m. Membrane filter was transferred to the surface of casein soya bean digest agar. Plates were incubated for 18 – 24 h at 30 – 35 °C and sample were filtered. Filter was washed with 3 quantities of 100 ml buffered sodium chloride – peptone solution with pH = 7. The number of colony forming units were count.

HPLC method

Chromatographic system

The chromatographic procedure was carried out using: Liquid chromatograph Shimadzu LC – 10 Advp equipped with 4.6 x 250 mm column RP – 18, ODS Spherisorb with particle size 5 μ m; detector SPD 10 AVvp – UV – VIS with fixed analytical wavelengths.

Chromatographic conditions

Isocratic mobile phase, prepared by mixing of filtered and degassed methanol : water = 50 : 50 v/v; column temperature 25 °C; flow rate: 1.0 ml/min; 210 nm analytical wavelength.

Reagents

L – Glutamic acid CRS, supplement mixture containing 100 mg L – Glutamic acid , methanol HPLC grade, water R.

Standard preparation

100.00 mg L – Glutamic acid CRS were dissolved in water R and were diluted to 100.0 ml with a solvent mixture – mobile phase.

Test preparation

100.00 mg from the tested substance were dissolved in water and were diluted to 100.0 ml with mobile phase. 1 ml from the obtained solution was dissolved with the same solvent mixture to obtain solutions with different concentrations.

RESULTS AND DISCUSSION

The results from the tests for determining the existence of microbial impurities show abnormal content ($1.5 \cdot 10^{-4}$ g) of bacteria and contaminants were identified mostly as non pathogenic bacilli of *Subtilis* group. Impermissible (pathogenic) contaminants as Enterobacteriaceae, *Staphylococcus aureus* were not found. Resistency factors show moderate ray sensitivity of the microorganisms, which are comparatively low sterilizing doses. Efficiency of the integral (total) dose of 10 Kgy was confirmed by the microorganism impurity tests after treatment and this dose guarantees a stable sterilizing coefficient.

Validation of HPLC procedure

HPLC method for quality control of L – Glutamic acid in supplement mixtures was developed and validated in accordance with ICH and European Pharmacopoeia criteria. Analytical parameters specificity, repeatability, limit of detection, limit of quantitation and linearity were studied and determined.

Specificity in respect of reagents

"Placebo" solution containing all reagents without active substances was prepared. There are no peaks in the chromatogram obtained from this solution with Rt of L – Glutamic acid.

Repeatability

Six (6) equal solutions from homogenous samples containing L – Glutamic acid were analyzed by HPLC method. Standard deviation (SD) and relative SD (RSD) were found. The results are presented on Table 1.

Table 1: Repeatability of samples from supplement mixture containing L - Glutamic acid 100 mg.

N	Obtained amount of L - Glutamic acid (100 mg)	\bar{X}	SD	RSD (\pm %)
1.	102.20	101.5	1.4321	0.44
2.	100.35			
3.	103.30			
4.	102.80			
5.	100.20			
6.	100.15			

Limit of detection

10 μ g for L - Glutamic acid, established on the base of ratio noise/signal - 1 : 3.

Limit of quantitation

40 μ g for L - Glutamic acid, established on the base of ratio noise/signal - 1: 10.

Linearity

The analytical parameter linearity was studied in concentration ratio 10 μ g - 150 mg. The accordance between the area of peaks,

measured in absorption units (AU) and concentrations in g/ml is proportional in the mentioned interval. The correlation coefficients is found to be 0.99746 at SD = \pm 3914.60 AU.

The identification and assay of L - Glutamic acid in supplement mixtures before and after γ - ray treatment were determined and compared. The results are shown on Table 2. There are no a significant difference between the putted quantity and obtained values for L - Glutamic acid concentration in the supplement mixture. At applied chromatographic conditions related substances and other impurities are not observed.

Table 2: Results from HPLC assay test for supplement mixture containing L - Glutamic acid 100 mg before and after γ - ray treatment

Sample N	Putted amount L - Glutamic acid (mg)	Obtained amount L - Glutamic acid before γ - ray treatment (mg)	Obtained amount L - Glutamic acid after γ - ray treatment (mg)
1.	100.0	102.20	102.40
2.	100.0	100.35	100.35
3.	100.0	103.30	102.70
4.	100.0	102.80	102.05
5.	100.0	100.20	101.55
6.	100.0	100.15	100.15
		RSD = \pm 0.44 %	RSD = \pm 0.082 %

CONCLUSION

Abnormal content ($1.5 \cdot 10^{-4}$ g) of bacteria is determined, without pathogenic microorganisms. There are no significant difference between content of L - Glutamic acid in supplement mixtures before and after γ - ray treatment.

REFERENCES

- Balch PA. Prescription for nutritional healing: a practical A - to - Z reference to drug - free remedies using vitamins, minerals, herbs and food supplements. 4th ed. New York: Avery Publishing Group; 2006;35-6.
- Uzunova S, Vassileva N, Dancheva E, Bozhilova D, Teolova E, Tachev A et al. Study of health risk from sun UV radiation on the population from the Plevan region. J Biomed Clin Res 2010;3(1):Suppl.1:44.
- Kale S, Sonawane A, Ansari A, Ghoge P, Waje A. Formulation and invitro determination of sun protection factor of Ocimum basilicum, Linn. leaf oils sunscreen cream. International Journal of Pharmacy and Pharmaceutical Sciences, 2010; 2, Suppl 4, 147-49.
- Warghat AR, Rampure NH, Wagh P. Effect of sodium azide and gamma rays treatments on percentage germination, survival, morphological variation and chlorophyll mutation in Musk Okra (Abelmoschus Moschatus L.). International Journal of Pharmacy and Pharmaceutical Sciences 2011;3(5):483-86.
- Silindir M, Özer AY. Sterilization Methods and the Comparison of E - Beam Sterilization with Gamma Radiation Sterilization. FABAD J Pharm Sci 2009;34:43-53.
- European committee for Standardization, Sterilization of Medical Devices - Validation and Routine Control of Sterilization by Irradiation, EN 552:1994, CEN, Brussels, 1994.
- International Organization for Standardization, Sterilization of Health Care Products - Radiation - Part 1: Requirements for the Development, Validation and Routine Control of a Sterilization Process for Medical Products (ISO 11137 - 1), Part 2: Establishing the Sterilization Dose (ISO 11137 - 2); Part 3: Guidance on Dosimetric Aspects, (ISO 11137 - 3), ISO, Geneva, 2006.
- International Organization for Standardization, Sterilization of Medical Devices - Microbiological Methods - Part 1: Determination of a Population of Microorganisms on Products (ISO 11737 - 1); Part 2: Tests of Sterility Performed in the Validation of a Sterilization Process (ISO 11737 - 2), ISO, Geneva, 2006.
- International Organization for Standardization. Sterilization of Medical Devices - General Requirements for Characterization of a Sterilizing Agent and the Development, Validation and Routine Control of a Sterilization Process for Medical Devices, ISO - 14937 : 2000 (E), Geneva.
- Hammad AA. Microbiological aspects of radiation sterilization. Trends in radiation sterilization of health care products. International Atomic Energy Agency, Vienna, Austria, 2008.
- Kucera J. Radiation sterilization of pharmaceuticals. Radioisotopy 1988;29:478.
- Slegers C, Tilquin B. Theoretical approach to the destruction or sterilization of drugs in aqueous solution. Radiat Phys Chem 2005;72:363-65.
- Ingram M, Roberts TA. Microbial Ecology of Food, Vol. I: Factors Affecting Life and Death of Microorganisms (ICMSF, Ed.). New York : Academic Press, 1994.
- Verma S, Singh V, Tanwar S. Pharmacognostic validation of whole plant of Convolvulus Pluricaulis Choisy (Convolvulaceae). Int J Pharm Pharm Sci 2012;4(1):241-46.
- Uzunova S, Bainova A, Chipilska L, Mechkueva L, Ivanov T, Tachev. Indicator and admissible levels for microbiological and chemical purity of cosmetic products. Journal of Environmental Protection and Ecology 2006;7(11):186-95.
- Garcia R, Howard B, La Rue R, Parton G, Walker J. Strategies for Gamma Sterilization of Pharmaceuticals. Pharmaceutical & Medical Packaging News, Canon Communications LLC, 2004.

17. Boess C, Bogel KW. Influence of radiation treatment on pharmaceuticals: a review: alkaloids, morphine derivatives, and antibiotics. *Drug Dev Ind Pharm* 1996;22:495-529.
18. Basly J - P, Basly I, Bernard M. ESR spectroscopy applied to the study of pharmaceuticals radiosterilization: cefoperazone. *J Pharm Biomed Anal* 1998;17:871-75.
19. Zegota H, Koprowski M, Zegota A. Effect of gamma radiation on cefotaxime in the solid state. *Rad Phys Chem* 1994;45(2):223-29.
20. Miyazaki T., Kaneko T, Yoshimura T, Crucq A - S, Tilquin B. ESR study of radiosterilization of antibiotics: ceftazidime. *J Pharm Sci* 1994;83:68-71.
21. Rongyao Y, Jilan W. Gamma radiolysis of aqueous solutions of glycosides. *Int J Radiat Appl Instrum Part C* 1990;35:488-92.
22. Allen AE, Gupta VD. Stability of hydrocortisone in polyethylene glycol ointment base. *J Pharm Sci* 1974;63:107-09.
23. Bosela AA, Salamah KK, Alsarra IA, El - Bagory IM. Reactivity of prednisolone to gamma radiation in aqueous and organic solutions. *J Drug Del Sci Tech* 2010;20(3):225-29.
24. Lee J - W, Sung N - Y, Kim J - K, Kim J - H, Raghavendran HRB, Yoo Y - C et al. Effect of gamma irradiation on spleen cell function and cytotoxicity of doxorubicin. *Chem Biol Interactions* 2008;173(3):205-14.
25. Varshney L, Dodke PB. Radiation effect studies on anticancer drugs, cyclophosphamide and doxorubicin for radiation sterilization. *Radiat Phys Chem* 2004;71:1103-11.
26. Damian G. EPR investigation of γ - irradiated antiemetic drugs. *Talanta* 2003;60:923-27.
27. Katusin - Razem B, Hamitouche K, Maltar - Strmecki N, Kos K, Pucic I, Britvic - Budicin S et al. Radiation sterilization of ketoprofen. *Radiat Phys Chem* 2005;73:111-66.
28. Slegers C, Baldacchino G, Le Pare D, Hickel B, Tilquin B. Radical mechanisms in the radiosterilization of metoprolol tartrate solutions. *Pharm Res* 2003;20:1977-83.
29. Basly JP, Duroux JL, Bernard M. Gamma radiation induced effects on metronidazole. *Int J Pharmaceutics* 1996;139(1-2):219-21.
30. Basly JP, Duroux JL, Bernard M. The effect of gamma radiation on the degradation of Salbutamol. *J Pharm Biomed Anal* 1997;15(8):1137-41.
31. Colak S, Korkmaz M. Investigation of radiosterilization and dosimetric features of sulfacetamide sodium. *J Pharm Biomed Anal* 2000;19:791-98.
32. Obrechikova D, Grigorova P, Tashkov V. A Study of the Behaviour of *Clinopodium vulgare* L. in the course of and after gamma - ray Treatment. *Compt. Rend Acad Bulg Sci* 1994;47(9):101-4.
33. Gardner WS, Miller WH. Reverse - phase liquid chromatographic analysis of amino acids after reaction with o - phthalaldehyde. *Anal Biochem* 1980;101:60-5.
34. Respaud R, Tournamille J - F, Croix C, Laborie H, Elfakir C, Viaud - Massuard M - C. Development of an with ion - pairing reversed - phase liquid chromatography method using a double detection analysis (UV and evaporative light scattering detection) to monitor the stability of Alimta[®] - pemetrexed preparations: Identification and quantification of L - Glu - tamic acid as a potential degradation product. *J Pharm Biomed Anal* 2011;54(2):411-16.
35. Tcherkas YV, Denisenko AD. Simultaneous determination of several amino acids, including homocysteine, cysteine and glutamic acid, in human plasma by isocratic reversed - phase high - performance liquid chromatography with fluorimetric detection. *J Chromatogr A* 2001;913(1-2):309-13.
36. Swanepoel E, de Villiers MM, du Preez JL. Fluorimetric method of analysis for D - norpseudoephedrine hydrochloride, glycine and L - glutamic acid by reversed - phase high - performance liquid chromatography. *J Chromatogr A* 1996;729(1-2):287-91.
37. Ma D, Zhang J, Sugahara K, Ageta T, Nakayama K, Kodama H. Simultaneous deter - mination of γ - aminobutyric acid and glutamic acid in the brain of 3 - mercaptopropionic acid - treated rats using liquid chromatography - atmospheric pressure chemical ionization mass spectrometry. *J Chromatogr B: Biomed Sci Appl* 1999;726(1-2):285-90.
38. Collins FS, George KS. Determination of glutamine and glutamic acid in biological fluids by gas chromatography. *J Chromatogr B: Biomed Sci Appl* 1978;145(3):456-63.
39. Pérez AS, Conde FL, Méndez JH. Polarographic determination of phenylalanine, tyrosine, methionine, glutamic acid and histidine with a dropping copper amalgam electrode. *Journal of Electroanalytical Chemistry and Interfacial Electrochemistry* 1976;74(3):339-46.
40. Almeida NF, Mulchandani AK. A mediated amperometric enzyme electrode using tetrathiafulvalene and l - glutamate oxidase for the determination of l - glutamic acid. *Anal Chim Acta* 1993;282(2):353-61.
41. Method 2.6.12. Microbiological examination of non - sterile products: total viable aerobic count and tests for specified microorganisms. *European Pharmacopoeia* 5.6, 01/2007:20612;4398-4404.