

UTILITIES OF CRATAEVA NURVALA

VANDANA KHATTAR*, ANKITA WAL

Institute of Pharmacy, Pranveer Singh Institute of Technology, Kanpur, U.P, India. Email: vandanapsit@gmail.com

Received: 30 April 2012, Revised and Accepted: 03 Jun 2012

ABSTRACT

The discovery of a novel chemical component from a medicinal plant may form the basis of development of various therapeutic agents with better activity. More than 500 medicinal plants have been reported to possess medicinal properties. *Crataeva nurvala* Buch Ham, (family: Cappariaceae) is one of the most common species among them. The whole plant possesses high medicinal value and is traditionally used in treating various ailments for human beings. The plant is used internally as well as externally. Externally, the paste or its leaves or skin of bark is applied in cervical adenitis, abscess and edematous wounds. The same paste is salutary in rheumatic joint for relief of pain. The pulp of leaves is applied on abdomen in splenic enlargement, with great benefit. Internally, *varuna* is used in vast range of diseases. The decoction of leaves given along with ghee relieves flatulence and abdominal pain. It also works well as a laxative, cholagogue, appetizer and vermicide, hence useful in anorexia, tumors, liver disorders, flatulent dyspepsia and helminthiasis. Phytochemically the plant has been investigated for flavonoids, glucosinolates, plant sterols, including lupeol, saponins, tannins, cardenolides, alkaloids, triterpenes and saponins. The plant has been demonstrated to possess multiple pharmacological activities such as anti-inflammatory, urolithiatic, antidiabetic, antibacterial, analgesic, anti-fertility, anti-diarrhoeal, antinociceptive and cardioprotective activity. This review highlights on the existing information particularly on the phytochemistry and various pharmacological properties of *crataeva nurvala* which may provide incentive for proper evaluation of the plant as a medicinal agent.

Keywords: *Crataeva nurvala*, Lupeol, Anti-inflammatory, Urolithiatic, Antibacterial activity.

INTRODUCTION

In ancient times nature has been an important source of medicinal agents and a large number of natural products have been identified and developed from natural sources based on their use in traditional medicine. Numerous medicinal plants are of global interest today because of their therapeutic and economic significance. According to the World Health Organization, approximately 80% of the world's population currently uses herbal medicines directly as teas, decocts or extracts with easily accessible liquids such as water, milk, or alcohol.

Crataeva nurvala is commonly known as *Varuna*. *Varuna* is one of the best litholytic herbs and has been used throughout the ages for the treatment of urolithiasis and crystalluria. *Varuna* is mentioned in vedic literature, its therapeutic use being known to ancient Ayurvedic physicians, especially as a blood purifier, to maintain homeostasis. The plant has various synonyms in Ayurvedic scriptures delineating its peculiarities viz. *triparna-trifoliata*, *bilvapatra*- leaves resemble to those of *bilva* (*Aegle marmelos*). *Vrttaphala* - fruits, ovoid berries, *asmari-ghna*- litholytic, *tiktakitter* etc. *Maharsi Susruta* has mentioned *varuna* as a litholytic agent in treating *kapha* and *vata* varieties of *asmari* (*calculi*)^{1,2}.

It is an appetizer, febrifuge, diuretic and litholytic in properties. It is used in diseases like urinary disorders, urinary calculi, blood disorders, worms and tumors. The bark of the tree is an important drug for problem affecting the kidneys and bladder. In Ayurveda, the bark of the *Crataeva* has been traditionally used to heal kidney stones for more than 3,000 years. Findings of several studies undertaken by contemporary scientists have authenticated that the herb neutralizes the enzyme called glycolate oxidase and this particular effect of the herb lessens the production of oxalates by the body³.

The purpose of the present study is to gather together the available published information on the different vernacular names, phytoconstituents and pharmacological activity of the plant.

Taxonomic description

The plant is classified as shown in Table.1.

Table 1: Taxonomic classification of *crataeva nurvala*

Kingdom	Plantae
Division	Magnoliophyta
Class	Magnoliopsida
Family	Cappariaceae
Genus	<i>Crataeva</i>
Species	<i>Nurvala</i>

Vernacular names⁴

Sanskrit: *Varun*, *Tiktshaak*

Hindi: *Baruna*, *Barna*

Bengali: *Varne*, *Borun*

English: Three-leaved caper

Kannada: *Bitusi*, *Holenekki*, *Holethumbe*, *Maavilanga*, *Mata maavu*, *Naaram bele*, *Vitasi*, *Neervaala mara*, *Sethu bandhana*, *Vaayu varuna*, *Nervaala*

Malyalam: *Nirmatalam*, *Nirval*

Marathi: *Haravarna*, *Karvan*, *Kumla*, *Nirvala*, *Ramala*, *Varun*, *Vaayuvarna*

Tamil: *Mavilingam*, *Narvala*, *Varanam*, *Maavilangam*, *Maralingam*

Telgu: *Ulimidi*, *Bilvaram*, *Chinnavulimidi*, *Maagalingam*, *Maaredu*, *Peddamaagalingam*, *Peddavulimidi*, *Thellavulimidi*.

Habitat

Varuna is a small tree, often cultivated throughout India, especially along the streams and riverbanks. It is distributed in sub-Himalayan tracts and is indigenous to Tamil Nadu, Kerala and Karnataka. It is found in abundance, in Kerala, Madhya Pradesh, Bengal and Assam. The plant flowers mostly grow in March and fruits in June.

Botanical description

Macroscopy

Crataeva nurvala is a moderate sized deciduous tree. The mature bark is typically 6-15 cm long and 3-10 cm wide with a thickness varying from 5-15 mm. The outer surface of the bark is gray to grayish-brown and rough, due to the presence of several small and rounded lenticels. The inner surface is smooth and whitish-brown to buff colored. Leaves are trifoliolate. Flowers are white or cream colored. Fruits have multiple seeds and ovoid berries, 2.5 cm in diameter and seeds are embedded in the yellow, fleshy pulp of the fruits⁵.

Microscopic

Transverse section of mature stem bark shows, an outer cork composed of thin walled, rectangular and tangentially elongated cells, phellogen single layered, thin walled, tangentially elongated cells followed by a wide secondary cortex, consisting of thin-walled,

polygonal to tangentially elongated cells with a number of starch grains, starch grains mostly simple, occasionally compound with 2-3 components also present, large number of stone cells in groups of two or more, found scattered in secondary cortex, single stone cells not very common, stone cells vary in size and shape, being circular to rectangular or elongated with pits and striations on their walls, stone cells distributed somewhat in concentric bands in phloem region except in inner region of phloem which is devoid of stone cells, secondary phloem comparatively a wide zone, consisting of sieve tubes, companion cells, parenchyma and groups of stone cells, alternating with medullary rays, sieve elements found compressed forming ceratenchyma in outer phloem region, whereas in inner region of phloem, intact, medullary rays mostly multiseriate composed of thin-walled, radially elongated cells, tangentially elongated towards outer periphery, a number of starch grains similar to secondary cortex also present in phloem and ray cells, few rhomboidal crystals of calcium oxalate also found in this region, inner most layer is cambium².

Phytoconstituents of *Crataeva nurvala*: Flavonoids, glucosinolates, plant sterols, including lupeol, saponins, and tannins. The largest group of plant secondary metabolites is the terpenes. Triterpenes are a major subgroup of the terpene superfamily⁶. These compounds are low-molecular-weight metabolites that are synthesized from mevalonate via a 30-carbon intermediate, 2, 3-oxidosqualene. The cyclization of 2, 3-oxidosqualene by oxidosqualene cyclases (OSCs) to either tetracyclic sterols, through the activity of cycloartenol synthase or pentacyclic triterpenes, through the activity of enzymes such as β -amyrin synthase, α -amyrin synthase and lupeol synthase represents a critical branch point between primary and secondary metabolism⁷. Sterols have been studied extensively and function as structural components of membranes and as precursors of steroidal hormones in both plants and animals, and also have important signaling functions⁸.

Part used: Stem bark, Root bark, Leaves, Fruits, Flowers⁹

Plant extract: Extraction, as the term is used pharmaceutically, involves the separation of medicinally active portions of plant or animal tissues from the inactive or inert components by using selective solvents in standard extraction procedures. The purposes of standardized extraction procedures for crude drugs are to attain the therapeutically desired portion and to eliminate the inert material by treatment with a selective solvent known as menstrum.

The extract thus obtained may be ready for use as a medicinal agent in the form of tinctures and fluid extracts, it may be further processed to be incorporated in any dosage form such as tablets or capsules, or it may be fractionated to isolate individual chemical entities such as ajmalicine, hyoscine and vincristine, which are modern drugs. Thus, standardization of extraction procedures contributes significantly to the final quality of the herbal drug. General Methods of Extraction of Medicinal Plants- Maceration, Infusion, Digestion, Decoction, Percolation, Hot Continuous Extraction (Soxhlet), Aqueous Alcoholic Extraction by Fermentation, Counter-current Extraction, Ultrasound Extraction (Sonication), Supercritical Fluid Extraction and Phytonics Process¹⁰.

Generally used plant extract are as- *C. nurvala* aqueous extract (CNAE), *C. nurvala* ethanolic extract (CNEE), *C. nurvala* chloroform extract (CNCE), *C. nurvala* petroleum ether extract (CNPEE).

PHYTOCHEMISTRY

Phytochemical studies showed that stem bark of the plant, fruits, leaves and root bark contain different constituents as-

Stem bark of the plant contains saponins, flavonoids, sterols and glucosinolates and ceryl alcohol, friedelin, cadabacine diacetate, lupeol, betulinic acid and diosgenin¹¹.

Fruits contain glucocapparin, beta-sitosterol, triacontane, triacontanol, cetyl and ceryl alcohol, pentadecane, octanamide, 12-tricosanone and friedelin^{12, 13, 14, 15}.

Leaves contain L-stachydrine, dodecanoic anhydride, methyl pentacosanoate, kaemferol-0- α -D-glucoside and quercetin-3-0- α -D-glucoside¹⁶.

Root bark contains rutin, quercetin, lupeol, varunol and β -sitosterol. Presence of alkaloids has been reported in bark and stems¹⁷.

ETHNOMEDICINAL USES


The stem, roots and leaves of varuna have great medicinal value. The plant is used internally as well as externally. Externally, the paste or its leaves or skin of bark is applied in cervical adenitis, abscess and edematous wounds. The same paste is salutary in rheumatic joint for relief of pain. The pulp of leaves is applied on abdomen in spleen enlargement, with great benefit.






Internally, varuna is used in vast range of diseases. The decoction of leaves given along with ghee relieves flatulence and abdominal pain. It also works well as a laxative, cholegogue, appetizer and vermicide, hence useful in anorexia, tumors, liver disorders, flatulent dyspepsia and helminthiasis. The decoction of skin of varuna is given along with honey in abscesses for the potent anti inflammatory action. The decoction of skin of varuna alone is used as blood purifier in gout, internal abscess and adenitis and to reduce body fats in obesity. The leaves cooked as vegetables are also benevolent in obesity. Varuna has the cardinal properties as litholytic, diuretic and urinary antiseptic.

The decoction of bark skin or roots is beneficial in urinary calculi, dysuria and cystitis. The decoction of leaves effectively alleviates the fever and associated delirium. The fresh juice of its leaves is useful as a bitter tonic. Varuna is used as a cholegogue, anthelmintic and anti-amoebic in both intestinal and hepatic infestations¹⁸.

Thus, the bark of *C. nurvala* is contraceptive and cytotoxic and is especially useful in urinary disorders, kidney bladder stones, fever, vomiting and gastric irritation. Root and bark are laxative and lithontripic and increase appetite and biliary secretion¹⁹. Leaves are externally rubefacient and used in rheumatism; internally they are given as febrifuge and tonic^{20, 21}. Although, a large number of compounds have been isolated from various parts of *C. nurvala*, a few of them have been studied for biological activity and the structure of some of these bioactive compounds has been presented in Table 2.

Table 2: Structures and activities of some active compounds from *crataeva nurvala*.

Name	Structure	Uses
Lupeol		Responsible for antihepatotoxicity, antitumor, anti-inflammatory ²² , chemoprotective agents ²³ , antimicrobial ²⁴ , antiarthritic ²⁵ , antihyperglycemic, antioxidant, cytotoxic, hypotensive, antiedemic, and antiperoxidant activities ²⁶ .

β -sitosterol		Responsible for antidiabetic, antioxidant ²⁷ , atherosclerosis, prostate enlargement, antihypercholesterolemia, androgenic, anticancer, antimutagenic, antifeedent, antigonadotrophic, and antiprostatic ²⁸ .
Betulinic Acid		Responsible for inhibition of HIV ^{29,30,31} , antibacterial, antimalarial, anti-inflammatory, anthelmintic, antioxidant properties ³² and antitumor-activity selective against human melanoma cells ³³ .
Quercetin		Responsible for anti-inflammatory ³⁴ , antihistamine, antioxidant ³⁵ , anticancer ^{36,37,38,39,40} , cataracts, preprostatin ⁴¹ , antiulcer effect ⁴² , in heart diseases ⁴³ and respiratory diseases as bronchitis and asthma ⁴⁴ .
Kaempferol		Responsible for Antiinflammatory, antioxidant, spasm, antiulcer, in diuretic, cough, inhibit fertility, anticancer ⁴⁵ and antidepressants ⁴⁶ .
Friedelin		Responsible for antiulcerogenic activity ⁴⁷ , significance cytotoxic activity against T-lymphoblastic leukemia (CEM-SS) cell line with the LC50 of 5, 8 μ g/ml. It has the ability to decrease the affinity of histamine on histamine H-1 receptor ⁴⁸ .

PHARMACOLOGICAL STUDIES

Pharmacological studies have confirmed that *Crataeva nurvala* exhibit a broad range of biological effects. However, the crude extract of the plant have been used as a traditional medicine for the treatment of various diseases, some of which are very interesting for possible future development.

Anti-fertility activity

The ethanol and aqueous extracts of the dried stem bark of the plant *Crataeva nurvala* Buch-Hum have been found to possess significant anti-fertility effects in rats. Both ethanol and aqueous extracts exhibited partial and complete resorption of implants at 300 and 600 mg/kg dose levels, respectively. The ethanol extract is found to be more active than the aqueous extract. However in estrogenic activity study, both the extracts increased uterine weight and caused opening and cornification of vagina in immature rats. This work justifies its effectiveness in preventing pregnancy in all rats at dose levels⁴⁹.

Analgesic and antidiarrhoeal activity

The leaves of medicinal plant '*Crataeva nurvala* Buch. Ham' was extracted in ethanol to evaluate the peripherally acting analgesic potential using acetic acid induced writhing and antidiarrhoeal activity using intestinal motility test both in mice. The crude extract showed significant ($P < 0.01$) analgesic activity at oral doses of 200 and 400mg/kg body weight with an inhibition of writhing 68.4% and 76.3% compared to 67% for the positive control. In the motility test, the crude extract at same oral doses showed 31.16% and 35.31% inhibition of intestinal propulsion of charcoal marker where as positive control group exhibited 36.25% inhibition of propulsion of charcoal through the intestine⁵⁰.

The antidiarrhoeal activity of ethanol extracts of *Crataeva nurvala* stem bark was evaluated using castor oil-induced diarrhoea model in rats. The gastrointestinal transit rate was expressed as the percentage of the longest distance traversed by the charcoal divided by the total length of the small intestine. The weight and volume of intestinal content induced by castor oil were studied by

enteropooling method. Like atropine (3mg/kg, i.p.) there were significant reductions in fecal output and frequency of droppings when the plant extracts 500 mg/kg doses were administered intraperitoneally compared with castor oil treated rats. This dose of the plant extracts significantly retarded the castor-oil induced enteropooling and intestinal transit. It significantly inhibited ($P < 0.001$) weight and volume of intestinal content⁵¹.

Antiarthritic activity

Lupeol is a naturally occurring triterpene isolated from *Crataeva nurvala* stem bark, and its ester lupeol linoleate was synthesized. The effects of lupeol and lupeol linoleate on the development of complement in adjuvant arthritis in rats were studied and compared with those of indomethacin. The effect of lupeol linoleate in reducing the foot-pad thickness and complement activity in arthritic rats was even greater than that of unesterified lupeol and indomethacin. Because complement is highly involved in inflammation, the results suggest that the anti-inflammatory activity of triterpenes may be due to their anticomplementary activity⁵².

Cardioprotective activity

Cyclophosphamide (CP), an alkylating agent widely used in cancer chemotherapy, causes fatal cardiotoxicity. In the present study, lupeol, a pentacyclic triterpene, isolated from *Crataeva nurvala* stem bark and its ester, lupeol linoleate were investigated for their possible cardioprotective effects against CP-induced toxicity. Male albino rats of Wistar strain were injected with a single dose of CP (200 mg/kg body weight, ip). In CP-administered rats, activities of lactate dehydrogenase and creatine phosphokinase were elevated in serum with a concomitant decline in their activities in the cardiac tissue. Significant increases ($P < 0.001$) in the levels of lipid peroxides and a decrease ($P < 0.001$) in the levels of enzymic (superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase, glucose-6-phosphate dehydrogenase and glutathione-s-transferase) and nonenzymic (reduced glutathione, vitamin C and vitamin E) antioxidants in the heart were also observed. The cardioprotective effects of lupeol (50 mg/kg body weight for 10 days orally) and its ester, lupeol linoleate (50 mg/kg body weight for 10 days orally) were evident from the significant reversal of the above alterations induced by CP. These observations highlight the antioxidant property of triterpenes and their cytoprotective action against CP induced cardiotoxicity⁵³.

Urolithic property

Despite modern techniques, the recurrence rate of Urolithiasis is approximately 50% within 5 years. Thus, there must be some drug that corrects the metabolic errors and prevents the formation of stone. In Ayurveda, a detailed description of urolithiasis is mentioned under the heading of Ashmari. This work was designed to study the effect of Varuna (*Crataeva nurvala*) on the experimental model of urolithiasis (albino rats). The study was categorized into two groups: Group I, treated and Group II, control. In all albino rats, stone was surgically implanted into the urinary bladder. Estimation of the urinary and serum electrolyte done periodically and x-rays were exposed at a regular interval. This study suggests the decoction of *Crataeva nurvala* is effective in the management of urolithiasis⁵⁴.

Antinociceptive activity

The antinociceptive effect of crude ethanolic extracts was evaluated by 'acetic acid' analgesic test method in mice. Crude ethanolic extracts of *Crataeva nurvala* (250–500 mg/kg PO) produced dose-dependent, significant ($p < 0.05-0.001$) antinociceptive effect against chemically induced nociceptive pain stimuli in mice. The results obtained in this study suggest that the antinociceptive effect of the extracts of *Crataeva nurvala* are peripherally and centrally-mediated. The findings of this experimental animal study indicate that crude ethanolic extracts of *Crataeva nurvala* possesses antinociceptive properties and thus lend pharmacological support to folkloric, anecdotal uses of 'borun' in the treatment and/or management of painful, arthritic inflammatory conditions⁵⁵.

Antidiabetic activity

Crataeva nurvala stem bark extracts have activity against in alloxan-induced diabetic albino rats. A comparison was made between the action of different extracts of *C. nurvala* and a known antidiabetic drug glibenclamide (600 µg/kg b. wt.). An oral glucose tolerance test (OGTT) was performed in diabetic rats: *C. nurvala* petroleum ether extract (CNPEE) and ethanolic extract (CNEE) showed significant ($P < 0.001$) antidiabetic activities. In alloxan-induced model, blood glucose level of these extracts on seventh day of study were CNPEE (126.33±13.703 mg/dl) and CNEE (126.66±13.012 mg/dl) when compared with diabetic control (413.50±4.752 mg/dl) and chloroform extract (320.83±13.516 mg/dl). In OGGT model (glucose loaded rats), CNPEE showed a glucose level of 178.83±3.070 mg/dl after 30 min and 131.66±2.486 mg/dl after 90 min, whereas CNEE showed 173.66±4.224 mg/dl after 30 min and 115.50±3.394 mg/dl after 90 min. These extracts also prevented body weight loss in diabetic rats⁵⁶.

Antiinflammatory activity

Lupeol has been extensively studied for its inhibitory effects on inflammation under *in vitro* and in animal models of inflammation. A comprehensive study showed that topical application of Lupeol (0.5 and 1 mg/ear) alleviated 12-*O*-tetradecanoylphorbol acetate (TPA)-induced inflammation in an ear mouse model. This study showed that topical application of Lupeol decreases myeloperoxidase levels [neutrophil specific marker] thus causing reduction in cell infiltration into inflamed tissues in mice. The anti-inflammatory potential of Lupeol could be assessed from the observation that Lupeol pretreatment significantly reduced prostaglandin E2 (PGE2) production in A23187-stimulated macrophages⁵⁷. Thus, lupeol treatment (5–9.37 mg/kg) was reported to exhibit anti-inflammatory activity with a maximum inhibition of 57.14%^{58, 59}. Lupeol is also reported to treat or reduce inflammation in a mouse model of arthritis, which is an inflammation associated disease⁵².

Anticancer activity

Recent studies have shown that diets rich in phytochemicals can significantly reduce cancer risk by as much as 20%. Epidemiological data suggest that the phytochemicals content of the diet is associated with a reduction in common cancers including cancers of the colon, breast, and prostate^{60, 61, 62}. A number of triterpenoids have shown promise as antineoplastic agents and exhibit anti-proliferative activity when tested against various cancer cell lines. These triterpenoids include members belong to the cycloartane, lupane, friedelane, dammarane, ursane, oleanane, limonoid and cucurbitacin family. Betulinic acid and its derivative also possess anticancer activity as have action against mouse leukemia^{63, 64, 65}.

Topical application of Lupeol [40 mg/kg/3 times a week] for 28 weeks was shown to significantly decrease tumor burden, tumor multiplicity and increase tumor latency period in the mouse model⁶⁶. Lupeol [2 mg/animal] was not only found to suppress the tumor growth, but also to impair head and neck cancer cell invasion by targeting NFκB signaling⁶⁷. The chemotherapeutic potential of Lupeol was also tested against the human hepatocellular carcinoma cell SMMC7721 cells. Lupeol treatment was shown to inhibit the growth and induce the apoptotic death of SMMC7721 cells. This study showed that Lupeol-induced growth inhibition and apoptosis is due to down-regulation of DR3 expression in SMMC7721 cells⁶⁸.

USES

The skin, roots and leaves of varuna have great medicinal value. The plant is used internally as well as externally. It is applied externally on wounds, reduces inflammation, loss of appetite, abdominal pain in liver disorders, worm infestation, dysuria, pain in urinary tract, urinary tract infection, fever and in general weakness.

Classical ayurvedic preparations

Varunadi quath- Useful in treatment of urinary calculi and urinary tract disorders, Burning Micturition, Pyelitis, Nephritis, U.T.I, Prostatitis, Dysurea, Prostate enlargement.

Varunadya ghrta- Used in Ayurveda for reducing phlegm, fat deposition, metabolic disorders, rheumatoid arthritis, migraine headaches and inflammatory conditions.

Varunadya taila- Used in treatment of renal calculus or stone.

Contraindications and Toxicology

No contraindications noted ever and no toxic effects seen.

CONCLUSION

In order to evaluate the overall implications of crataeva nurvala as having different activities as antidiabetic, anti-fertility activity, analgesic and antiarrhythmic activity, antiarthritic activity, cardioprotective activity, urolithic property, antinociceptive activity, Antiinflammatory activity and anticancer activity due to having different phytoconstituents. As this plant has no contraindication and no toxic effect, therefore it used as a valuable medicinal plant.

Thus, it is believed that detailed information presented in this review would help the researchers to get aware of this plant and extensive research should be undertaken on crataeva nurvala for establishing new therapeutic drugs for mankind.

REFERENCES

- Farnsworth NR. The Role of Ethnopharmacology in Drug Development. In: Bioactive Compounds from Plants (Ciba Foundation Symposium 154), Chadwick D.J. and J. Marsh (Eds.). Wiley, Chichester, UK. p. 2-21.
- <http://logayurveda.com/plant-profiles/204-crataeva-nurvala.pdf> (accessed jan3, 2012).
- <http://www.herbosophy.com.au/crataeva-crataeva-nurvala/> (accessed jan3, 2012).
- Remya M.B, Manu Somanath, Santhosh N, Reshmi Manayat, Dr. Sarala Samuel, Dr. Jolly. Crataeva nurvala, a valuable medicinal plant in the treatment of benign prostatic hyperplasia. Kerala Ayurveda Vaidyam. 2009; 2(3).
- Chopra RN, Chopra IC, Varma BS. Supplement to Glossary of Indian Medicinal Plants. New Delhi: C.S.I.R. Publication; 1969. P.20.
- Xu R, Fazio GC, Matsuda SP. On the origins of triterpenoid skeletal diversity. Photochemistry 2004; 65: 261-91.
- PhillipsDR, Rasbery JM, Bartel B, Matsuda SPT. Biosynthetic diversity in plant triterpene cyclization. Current Opinion in Plant Biology 2006; 9: 305-14.
- Lindsey K, Pullen ML, Topping JF. Importance of plant sterols in pattern formation and hormone signalling. Trends Plant Science. 2003; 8: 521-5.
- Anjaria Jayvir, Parabia Minoo, Bhatt Gauri and Khamar Ripal. A glossary of selected indigenous medicinal plants of India. 2nd ed. SRISTI Innovations 2002. p. 23.
- Handa Sukhdev Swami, Khanuja Singh Suman Preet, Longo Gennaro, Dev Dutt Rakesh Extraction Technologies for Medicinal and Aromatic Plants. ICSHP Trieste; 2008.
- Lakshmi V, Chauhan JS. Triterpenoids and related compounds from Crataeva nurvala. Planta Med. 1975; 27: 254-6.
- Buckingham J. Dictionary of natural products. London: Chapman and Hall; 1994.
- Heilbron I, Cook AH, Bunbury HM, Hey DH. Dictionary of organic compounds. London: Eyre and Spottiswoode; 1965.
- Sukhdev. Handbook of Terpenoids. Vol. II. Florida: CRC Press; 1989.
- Gagandeep, Meera, Kalidhar SB. Chemical investigation of Crataeva nurvala Buch. Ham. fruits. Indian J. Pharm. Sci. 2009; 71: 129-30.
- Gagandeep M, Kalidhar SB. Chemical constituents of Crataeva nurvala (Buch-ham) leaves. Indian J. Pharm. Sci. 2006; 68: 804-6.
- Ghani A. Medicinal plants of Bangladesh with chemical constituents and Uses. 2nd ed. Asiatic Society of Bangladesh, Dhaka, Ramna 2003. p.184.
- <http://www.herbalcureindia.com/herbs/crataeva-nurvala.htm> (accessed jan3, 2012).
- Malini MM, Baskar R, Varalakshmi P. Effect of lupeol, a pentacyclic triterpene, on urinary enzymes in hyperoxaluric rats. Jpn. J. Med. Sci. Biol. 1995; 48: 211-20.
- Walia N, Kaur A, Babbar SB. An efficient, *in vitro* cyclic production of shoots from adult trees of Crataeva nurvala. Buch. Ham. Plant Cell Rep. 2007; 26: 277-84.
- Sanayaima RK, Kaur A, Agrawal A, Babbar SB. Cryopreservation of *in vitro*-grown shoot tips of Crataeva nurvala, Buch. Ham, an important medicinal tree. Cryo Letters. 2006; 27: 375-86.
- Shirwaikar A, Manjunath Setty M, Bommu P, Krishnanand B. Effect of lupeol isolated from Crataeva nurvala Buch.-Ham. stem bark extract against free radical induced nephrotoxicity in rats. Ind. J. Exp. Biol. 2004; 42: 686-90.
- Surh YJ. Cancer chemoprevention with dietary phytochemicals. Nature Rev. Cancer. 2003; 3: 768-80.
- Shai LJ, McGaw MA, Aderogba LK, Mdee, Eloff JN. Four pentacyclic triterpenoids with antifungal and antibacterial activity from Curtisia dentata (Burm.f) C.A. Sm. leaves. J. Ethnopharmacol. 2008; 119: 238-44.
- Agarwal RB, Rangari VD. Antiinflammatory and antiarthritic activities of lupeol and 19a-h lupeol isolated from strobilanthus callosus and strobilanthus ixiocephala roots. Ind. J. Pharmacol. 2003; 35: 384-87.
- Sunitha S, Nagaraj M, Varalakshmi P. Hepatoprotective effect of lupeol and lupeol linoleate on tissue antioxidant defence system in cadmium-induced hepatotoxicity in rats. Fitoterapia. 2001; 72: 516-23.
- Li CR, Zhou Z, Lin RX, Zhu D, Sun YN et al. Beta-sitosterol decreases irradiation-induced thymocyte early damage by regulation of the intracellular redox balance and maintenance of mitochondrial membrane stability. J. Cell Biochem. 2007; 102: 748-58.
- Imanaka H, Koide H, Shimizu K, Asai T, Kinouchi N Shimizu et al. Chemoprevention of tumor metastasis by liposomal beta-sitosterol intake. Biol. Pharm. Bull. 2008; 31: 400-04.
- Yogeeswari Perumal, Sriram Dharmarajan. Betulinic Acid and Its Derivatives: A Review on their Biological Properties. Current Medicinal Chemistry. 2005; 12: 657-66.
- Fujioka T, Kashiwada Y, Kilkuskie RE, Cosentino LM, Ballas LM, Jiang JB, et al. Anti-AIDS agents, 11. Betulinic acid and platanic acid as anti-HIV principles from Syzygium claviflorum, and the anti-HIV activity of structurally related triterpenoids. J Nat Prod. 1994; 57(2): 243-7.
- Huang L, Ho P, Lee KH, Chen CH. Synthesis and anti-HIV activity of bi-functional BA derivatives. Bioorg. Med. Chem. 2006; 14: 2279-89.
- Chandramu C, Manohar RD, Krupadanam DGL, Dashavantha RV. Isolation, characterization and biological activity of betulinic acid and ursolic acid from Vitex negundo L. Phytother. Res. 2003; 17: 129-34.
- Selzer E, Pimentel E, Wacheck V, Schlegel W, Pehamberger H, Jansen B, et al. Effects of betulinic acid alone and in combination with irradiation in human melanoma cells. J Invest Dermatol. 2000; 114: 935-40.
- Stewart LK, Soileau L, Jeff, Ribnicky David, Wang Q, Zhong, Ilya R, Poulev Alexander, et al. Quercetin transiently increases energy expenditure but persistently decreases circulating markers of inflammation in C57BL/6J mice fed a high-fat diet. Metabolism - Clinical and Experimental. 2008; 57: 39-46.
- Davis W. Lamson, MS, ND, Matthew S. Brignall ND. Antioxidants and Cancer III: Quercetin Altern. Med Rev. 2000; 5(3): 196-208.
- Scambia G, Raneletti FO, Panici PB, et al. Quercetin induces type-II estrogen-binding sites in estrogenreceptor-negative (MDA-MB231) and estrogen-receptor-positive (MCF-7) human breast-cancer cell lines. Int J Cancer. 1993; 54: 462-6.
- Singhal RL, Yeh YA, Prajda N, et al. Quercetin down-regulates signal transduction in human breast carcinoma cells. Biochem Biophys Res Comm. 1995; 208: 425-31.
- Choi JA, Kim JY, Lee JY, et al. Induction of cell cycle arrest and apoptosis in human breast cancer cells by quercetin. Int J Oncol. 2001; 19: 837-844.
- Scambia G, Raneletti FO, Panici PB, et al. Quercetin inhibits the growth of a multidrug-resistant estrogenreceptor-negative MCF-7 human breast-cancer cell line expressing type II estrogen-binding sites. Cancer Chemother Pharmacol. 1991; 28: 255-8.

40. Scambia G, Raneletti FO, Panici PB, et al. Synergistic antiproliferative activity of quercetin and cisplatin on ovarian cancer cell growth. *Anticancer Drugs* 1990; 1: 45-8.
41. Shoskes D et al. Use of bioflavonoid quercetin in patients with long standing chronic prostatitis. *J. Am. Nutraceut. Assoc.* 1999; 2: 18.
42. Beil W, Birkholz C, Sewing KF. Effects of flavonoids on parietal cell acid secretion, gastric mucosal prostaglandin production and *Helicobacter pylori* growth. *Arzneimittel-forschung.* 1995; 45: 697-700.
43. Pace-Asciak CR, Hahn S, Diamandis EP, et al. The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: implications for protection against coronary heart disease. *Clin Chim Acta.* 1995; 235: 207-19.
44. Butland B. Diet, lung function, and lung function decline in a cohort of 2512 middle aged men. *Thorax* 2000; 55: 102.
45. Nguyen, T.T., E. Tran, C.K. Ong, S.K. Lee and P.T. Do et al. Kaempferol-induced growth inhibition and apoptosis in A549 lung cancer cells is mediated by activation of MEK-MAPK. *J. Cell Physiol.* 2003; 197: 110-21.
46. Hadizadeh, F., N. Khalili, H. Hosseinzadeh and R. Khair-Aldine. Kaempferol from saffron petals. *Iran. J. Pharma. Res.* 2003; 2: 251-2.
47. Queiroga CL, Silva GF, Dias PC, Possenti A, de Carvalho JE. Evaluation of the antiulcerogenic activity of friedelan-3beta-ol and friedelin isolated from *Maytenus ilicifolia* (Celastraceae). *J Ethnopharmacol.* 2000; 72(3): 465-8.
48. Agung Endro Nugroho, Ratna Asmah Susidarti, Puji Astuti. Effects of Friedelin from *Eugenia Chlorantha* Duthie on Physiological Receptors-operated Guinea-Pig Trachea Contraction. *JPRCP* 2011; 1(2): 71-8.
49. BhaskarV. H., Profulla Kumar M.2, Balakrishnan B. R. Balakrishnan N. and Sangameswaran B. Evaluation of the anti-fertility activity of stem bark of *Crataeva nurvala* buch-hum. *AJB* 2009; 8(22): 6453-56.
50. Khatun Farjana, Md. Howlader Amran, Apu Sarker Apurba, Bachar Sitiesh C, Qais Nazmul Evaluation of analgesic and anti-diarrhoeal properties of the ethanolic extract of *crataeva nurvala* buch. ham (capparidaceae) leaves. *IJPSRR.* 2012; 12(2): 5-8.
51. Inayathulla, Shariff W R, Karigar Asif, Sikarwar Mukesh. Ethanolic extract of *crataeva nurvala* root bark in experimental animals. *IJPPS* 2010; 2: 158-61.
52. Geetha T, Varalakshmi P. Anticomplement activity of triterpenes from *Crataeva nurvala* stems bark in adjuvant arthritis in rats. *Gen Pharmacol.* 1999; 32(4): 495-7.
53. Sudharsan PT, Mythili Y, Selvakumar E, Varalakshmi P. Cardioprotective effect of pentacyclic triterpene, lupeol and its ester on cyclophosphamide-induced oxidative stress. *Hum Exp Toxicol.* 2005; 24(6): 313-18.
54. Agarwal S, Gupta SJ, Saxena AK, Gupta N, Agarwal S. Urolithic property of *Varuna* (*Crataeva nurvala*): An experimental study. *Ayu.* 2010; 31(3): 361-6.
55. Alam MA, Haque ME, Shilpi JA, Daulla KA. Antinociceptive effect of the crude ethanolic extract of *crataeva nurvala*. buch. on mice. *Bangl. J. Vet. Med.* 2006; 4(1):65-8.
56. Sikarwar MS, Patil MB. Antidiabetic activity of *Crataeva nurvala* stem bark extracts in alloxan-induced diabetic rats. *J. Pharm. Bioall. Sci.* 2010; 2: 18-21.
57. Fernández MA, de las Heras B, García MD, Sáenz MT, Villar A. New insights into the mechanism of action of the anti-inflammatory triterpene lupeol. *J. Pharm. Pharmacol.* 2001; 53: 1533-9.
58. Nguemfo EL, Dimo T, Dongmo AB, Azebaze AG, Alaoui K, Asongalem AE, et al. Anti-oxidative and anti-inflammatory activities of some isolated constituents from the stem bark of *Allanblackia monticola* Staner L.C (Guttiferae) *Inflammopharmacology.* 2009; 17:37-41.
59. Saleem Mohammad. Lupeol, A Novel Anti-inflammatory and Anti-cancer Dietary Triterpene. *Cancer Lett.* 2009 November 28; 285(2): 109-115.
60. Liby KT, Yore MM, Sporn MB. Triterpenoids and rexinoids as multifunctional agents for the prevention and treatment of cancer. *Nat. Rev. Cancer.* 2007; 7: 357-69.
61. Setzer WN, Setzer MC. Plant-derived triterpenoids as potential antineoplastic agents. *Mini Rev Med Chem.* 2003; 3: 540-56.
62. Bradford PG, Awad AB. Phytosterols as anticancer compounds. *Mol. Nutr. Food Res.* 2007; 51: 161-70.
63. Min BS, Gao JJ, Nakamura N, Hattori M. Triterpenes from the spores of *Ganoderma lucidum* and their cytotoxicity against meth-A and LLC tumor cells. *Chem. Pharm. Bull. (Tokyo)* 2001; 48: 1026-33.
64. Nagumo A, Takanashi K, Hojo H, Suzuki Y. Cytotoxicity of Betulinic acid derivative hopane-32-ol against mouse leukemia L1210 and P388 cells in vitro. *Toxicol. Lett.* 2001; 58: 309-13.
65. Molnár J, Gyémánt N, Tanaka M, Hohmann J, Bergmann-Leitner E, Molnar P, et al. Inhibition of multidrug resistance of cancer cells by natural diterpenes, triterpenes and carotenoids. *Curr. Pharm. Des.* 2006; 12: 287-311.
66. Saleem M, Afaq F, Adhami VM, Mukhtar H. Lupeol modulates NF-kappa B and PI3K/Akt pathways and inhibits skin cancer in CD-1 mice. *Oncogene* 2004; 23: 5203-14.
67. Lee TK, Poon RT, Wo JY, Ma S, Guan XY, Myers JN, et al. Lupeol suppresses cisplatin-induced nuclear factor-kappa B activation in head and neck squamous cell carcinoma and inhibits local invasion and nodal metastasis in an orthotopic nude mouse model. *Cancer Res.* 2007; 67: 8800-09.
68. Zhang L, Zhang Y, Zhang L, Yang X, Lv Z. Lupeol, a dietary triterpene inhibited growth, and induced apoptosis through down-regulation of DR3 in SMMC7721 cells. *Cancer Invest.* 2009; 27: 163-70.