

ANTISPERMATOGENIC EFFECT OF CARICA PAPAYA SEED EXTRACT ON STEROIDOGENESIS IN ALBINO RATS

J. LAKSHMAN AND C. CHANGAMMA*

Department of Zoology, S.V. University, Tirupati, A.P., India. Email: challa_changamma@yahoo.co.in

Received: 05 Sep 2012, Revised and Accepted: 16 Oct 2012

ABSTRACT

Steroidogenesis plays a key role in the development and maintenance of male reproductive function and fertility. The objective of the present study was to investigate the effect of Carica papaya seed extract on steroidogenesis. The cholesterol levels in testes were significantly decreased by the Carica papaya seed extraction indicating decreased mobilization towards androgenesis which leads to decreased steroidogenesis and thereby inhibition of spermatogenesis in testes. It is noticed that the liver cholesterol was significantly enhanced with reduced blood cholesterol. The lowering of the 3 β -HSD and 17 β -HSD activity levels in the testes suggest the antifertility agents interfere with steroid hormone biosynthesis, which ultimately result in impaired spermatogenesis and infertility.

Keywords: 3 β -HSD, 17 β -HSD, Carica papaya, Cholesterol, Spermatogenesis.

INTRODUCTION

Medicinal plants have successfully been used to induce sterility in laboratory animals [1-3]. Pawpaw seed (*Carica papaya*) [4,5] reported high success in using. Pawpaw (*Carica papaya*) seeds had been used as fertility control agents in some animal models and even on human beings [6,7] respectively. Chloroform extract of papaya seeds tested in langan monkeys for one year, caused a steady decrease in sperm production with no sign of toxicity [8,9]. Crude extract fed to male rats deteriorated quantity and quality of the sperm [10,11]. At higher dose, it provided 100% contraception, but resulted in weight loss, possibly due to toxicity [7,12,9]. Suppression of spermatogenesis was observed in rats followed the administration of papaya seed extract [13]. The Oral administrations of extract induced reversible male infertility [14,17]. The biochemical studies on carbohydrate metabolism reveals the decreased oxidative metabolism [18]. Male reproduction is a multifaceted process that involves the testes, epididymis, accessory sex glands and associated hormones. Testes perform two highly organized and intricate events, called spermatogenesis and steroidogenesis, which are vital for the perpetuation of life. Spermatogenesis, a highly dynamic and synchronized process, takes place within the seminiferous tubules of the testis with the support of somatic Sertoli cells, leading to the formation of mature spermatozoa from undifferentiated stem cells [19]. Pawpaw (*Carica papaya*) seeds contain antifertility properties, particularly of the seeds [20]. A complete loss of fertility has been reported in male rabbits, rats and monkeys fed an extract of papaya seeds [20,8,21]. Thus the steroidogenesis is important for spermatogenesis. Hence in the present study it is important to know how the steroid enzymes are modulating during spermatogenesis and antispermatogenesis.

MATERIALS AND METHODS

Healthy adult male Wistar strain albino rats (90days old, weight 160 \pm 10g) were administered with 100mg/kg body wt/day of alcoholic extract of papaya seed orally for 15days. The alcoholic extract was prepared according to WHO 1983 [22] protocol CG-04. Seeds were shed-dried, powdered and extracted with 95%ethanol (v/v) at 55-60 $^{\circ}$ C for 3h. The solvent was distilled off under reduced pressure; the resulting mass was dried under vacuum and kept at 24 $^{\circ}$ C until use. The control animals were given normal saline or sterile distilled water. Both control and experimental rats were maintained in standard air conditioned animal house at a temperature of 25 \pm 2 $^{\circ}$ C, exposed to 12-14h day light and fed on standard rat feed obtained from Hindustan Lever Ltd., Bombay, India. The usage of animals was approved by the Institutional Animal Ethics committee (Regd.No. 438/01/a/CPCSEA/dt.17/02/2001) in its resolution number 9/IAEC/SVU/Zool/dt.4-3-2002.

Twenty four hours after the last dose, the animals were autopsied. The tissues like testes, epididymis, seminal vesicle, prostate gland

and liver were isolated, chilled immediately and blood was collected, used for biochemical analysis. The cholesterol estimated by Natelson 1971 [23]. The enzymes like 3 β -HSD&17 β -hydroxy steroid dehydrogenase by Bergmeyer 1974 [24] were estimated in control and experimental rat tissues.

RESULTS AND DISCUSSION

The data represented in tables 1-3 shows the effect of Carica papaya seed extraction on cholesterol in reproductive and non reproductive tissues and 3 β -HSD & 17 β -HSD in testes of albino rats.

The cholesterol levels were significantly decreased in testes as it is necessary for steroidogenesis (table-1). This result indicates the decreased steroidogenesis which leads to decreased spermatogenesis. Cholesterol is the precursor of the steroid hormones [25], providing the backbone of the steroid molecule. The biosynthesis of testosterone directly from cholesterol can only occur in the Leydig cells [26].

Inter- group comparison

- a- P < 0.001.statistically significant when compared control with experimental,
- b- P < 0.01.statistically significant when compared control with experimental,
- e- Non significant changes.

The impact of treatment on cholesterol is more in liver (+59.39), the central organ in cholesterol metabolism than in blood (table-2). In sex accessories there were no significant changes in epididymis while in seminal vesicle and prostate cholesterol levels were slightly(P<0.01) increased.

Cholesterol is one of the most important sterols and is a structural component of membranes as well as the precursor for bile acids and steroid hormones. Cholesterol is a sterol with special functions in various tissues and organs. First of all, it is a structural component of all cell membranes. Furthermore, it is the precursor molecule of steroid hormones, such as progesterone, testosterone and cortisol. The unsaponifiable fraction of the neutral lipid fraction of the rat testis represents primarily cholesterol and steroids [27]. Cholesterol is an important precursor for the steroid hormones. The testis and its metabolism are dependent on the plasma and endogenously synthesized cholesterol. Hence the cholesterol levels were estimated and found to be significantly decreased in the experimental rat testis. This observation indicates either its decreased uptake from the plasma or increased synthesis or decreased mobilization towards androgenesis or decreased catabolism. This was supported by enhanced blood cholesterol levels (Table-2) by the treatment in the present study.

The spermatogenesis is a complex process which is strictly regulated by the hypothalamo-pituitary-testicular axis, which involves the pituitary gonadotropins, luteinizing hormone (LH) and follicle-stimulating hormone (FSH). Apart from LH, FSH and androgens, various growth factors, hormones and estrogens are involved in regulating the testicular functions [28]. Thus the reduced levels of cholesterol in testes by the treatment in the present study indicate the antispermatic effect of Carica papaya seed extraction through decreased steroidogenesis [29].

The activity levels of 3 β -HSD and 17 β -HSD the key enzymes of androgenesis were decreased significantly in the treated rat testis (table-3, fig-3). The decreased cholesterol content reveals that there is no dearth of substrate for steroidogenesis. Since the 3 β -HSD and 17 β -HSD activity levels were decreased. Carica papaya seed extraction inhibits testicular steroidogenesis. Suggesting the impaired steroidogenesis [29,30]. There is decreased activity levels of 3 β -HSD and 17 β -HSD in the testes suggest the antifertility agents interfere with steroid hormone biosynthesis, which ultimately result in impaired spermatogenesis and infertility. The kinetic characteristics of 17 β -HSD were determined in the cell system which would reflect the native kinetic properties of the enzyme under the influence of native intracellular milieu [30].

Two different pathways of androgen biosynthesis have been reported in testis, (1) progesterone pathway or Δ^4 -pathway and (2) dehydroepiandrosterone pathway or Δ^5 -Pathway. 3 β -HSD acts

on C-19 and C-21 steroids by specifically acting on 3 β -hydroxy groups. This enzyme has NAD⁺ as the preferred cofactor and is localized in the microsomes. It converts the pregnenolone to progesterone. This enzyme is almost irreversible reaching equilibrium towards the progesterone formation.

The 17 β -hydroxysteroid dehydrogenases (17 β -HSD enzymes) are a group of alcohol oxidoreductases which catalyse the dehydrogenation of 17-hydroxysteroids in steroidogenesis. The major reactions catalysed by 17 β -HSD the conversion of androstenedione to testosterone are in fact hydrogenation (reduction) rather than dehydrogenation (oxidation) reactions that can affect the primary and/or secondary sex characteristics of both males and females. In sex steroid metabolism 17 β -hydroxysteroid dehydrogenases (17 β -HSDs) catalyze the final steps in androgen and estrogen biosynthesis, thus playing a crucial role in the biosynthesis and inactivation of sex steroid hormones [31].

The Carica papaya seeds contain active ingredients such as caricacin, an enzyme carpasemine, a plant growth inhibitor, and oleanolic glycoside [32], which caused sterility in male rats [33]. Pawpaw (*Carica papaya*) seeds yield 660-760mg (bactericidal a glycone of glucotropaeolin benzyl isothiocyanate), a glycoside, sinigrin, the enzyme myrosin, and carpasemine. Hence oleanolic glycoside, sinigrin is acting on spermatogenesis by inhibiting the steroidogenesis which leads to antispermaticogenesis in the present study.

Table 1: The Levels of Cholesterol in reproductive tissues of control and papaya seed extraction treated rats.

S. No.	Name of the tissue	Control	Experimental	%Change
1.	Testis (mg/g wet wt.)	0.685±0.054	0.548±0.039	-20.01 ^a
2.	Epididymis (mg/g wet wt.)	1.345±0.101	1.307±0.099	-3.47 ^e
3.	Seminal Vesicle (mg/g wet wt.)	0.265±0.014	0.291±0.017	+9.81 ^b
4.	Prostate gland (mg/g wet wt.)	0.279±0.015	0.319±0.019	+14.34 ^b

Values are mean ± S.E.M (n = 6)

Table 2: The Levels of Cholesterol in Liver and Blood of control and papaya seed extraction treated rats.

S. No.	Name of the Tissue	Control	Experimental	% Change
1.	Liver (mg/g wet wt.)	5.32±0.421	8.48±0.736	+59.39 ^a
2.	Blood (mg/100ml)	194.85±10.42	160.29±12.04	-17.74 ^a

Values are mean ± S.E.M (n = 6)

Table 3: The levels of 3 β -HSD&17 β -HSD in control and Papaya seed extraction treated rat testes.

S. No.	Parameter	Control	Experimental	% Change significance
1.	3 β -HSD (μ mol NAD ⁺ reduced/mg protein/min)	0.475±0.011	0.330±0.012	-30.53 ^a
2.	17 β -HSD (μ mol NADPH oxidized /mg protein/min)	0.592±0.023	0.492±0.031	-16.89 ^a

Values are mean ± S.E.M (n = 6)

CONCLUSION

It is concluded that the Carica papaya seed extraction decreases the testicular cholesterol levels and steroidogenic enzyme levels which leads to impaired spermatogenesis and infertility.

ACKNOWLEDGEMENTS

The authors were grateful to UGC, New Delhi for financial assistance.

REFERENCES

- Gary S K & Gary, Antifertility screening VII. Effect of five indigenous plant parts on early pregnancy in albino rats. Ind J Med Res., 1971; 56: 302-306.
- Bodharkar S L, Gray S K & Mathus V S, Antifertility screening parts 1X. Effect of five indigenous plants on early pregnancy in female albino rats Ind J of Med Res., 1974; 62: 831-837.
- Das R P, Effect of Papaya seeds on the genital organs and fertility of male rats. Ind J Expt Bio, 1980; 18: 408-409.
- Udoh P & Kehinde A, Studies on antifertility effects of pawpaw seeds (*Carica papaya*) on the gonads of male albino rats. Phytotherapeutic Research, 1999; 13 (3): 226-228.
- Ekanem S B & Okoronkwo T E, Pawpaw seed as a fertility control agent on male Nile tilapia. NAGA,World Fish Center Quarterly, 2003; 26(2): 8-10.
- Lohiya N K, Manivannan B, Mishra P K, Pathak N, Sriram S, Bhande S S and Panneerdoss S, Ultra-structural changes in the testis and epididymis of rats following treatment with the benzene chromatographic fraction of the chloroform extract of the seeds of Carica papaya. Phytotherapeutic Research, 2004; 18 (4): 285-289.
- Lohiya N K, Manivannan B, Mishra P K, Pathak N, Sriram S, Bhande S S and Panneerdoss S, Chloroform extract of *Carica papaya* seeds induces long-term reversible azoospermia in langur monkey. Asian Journal of Andrology, 2002; 4: 17-26.
- Lohiya N K, Mishra P K, Pathak N, Manivannan B, Bhande S S, Panneerdoss S and Sriram S, Efficacy trial on the purified compounds of the seeds of Carica papaya for male contraception in albino rat. Reproductive Toxicology, 2005; 20(1): 135-48.
- Mojica-Henshaw M P, Francisco A D, De Guzman F, Tigno X T, Possible immunomodulatory actions of Carica papaya seed extract. Clinical Hemorheology and Microcirculation, 2003; 29: 219-229.

10. Kamal R, Gupta R S and Lohiya N K, "Plants for male fertility regulation." *Phytotherapy Research*, 2003; 17(6): 579-90,
11. Kusemiju O, Noronha C, Okanlawon A, The effect of crude extract of the bark of *Carica papaya* on the seminiferous tubules of male Sprague-Dawley rats. *Nigerian Postgraduate Medical Journal*, 2002; 9: 205-209.
12. Uche-Nwachi E O, Ezeokoli D C, Adogwa A O and Offiah V N, Effect of water extract of *Carica papaya* seed on the germinal epithelium of the seminiferous tubules of Sprague Dawley rats. *Kaibogaku Zasshi*, 2001; 76(6): 517-521.
13. Udoh F V, Uodh P B & Umoh E E, Activity of Alkaloid extract of *Carica papaya* seeds on Reproductive functions in male Wistar rats. *Pharmaceutical Biology*, 2005a; 4 (3): 563-567.
14. Udoh P, Essien I, Udoh F, Effects of *Carica papaya* (paw) seeds extract on the morphology of pituitary-gonadal axis of male Wistar rats. *Phytotherapy Research*, 2005b; 19: 1065-1068.
15. Lohiya N K, Goyal R B, Antifertility investigations on the crude chloroform extract of *Carica papaya* Linn. Seeds in male albino rats. *Ind J Exp Biol*, 1992; 30: 1051-5.
16. Lohiya N K, Pathak N, Mishra P K, Manivannan B, Reversible contraception with chloroform extract of *Carica papaya* Linn. seeds in male rabbits. *Reprod Toxicol*, 1999a; 13: 59-66.
17. Changamma C, Biochemical studies on the antispermatogenic effect of *Carica papaya* Linn. Seed extraction, *Bulletin of Pure and Applied Sciences*, 2008; 27 (2): 215-219.
18. Hess R A, Renato de Franca L, Spermatogenesis and cycle of the seminiferous epithelium, *Adv Exp Med Biol*, 2008; 636: 1-15.
19. Lohiya N K, Mishra P K, Pathak N, Manivannan B, Jain S C, Reversible azoospermia by oral administration of the benzene chromatographic fraction of the chloroform extract of the seeds of *Carica papaya* in rabbits. *Advances in contraception*, 1999b; 15: 141-161.
20. Pathak N, Mishra P K, Manivannan B, Lohiya N K, Sterility due to inhibition of sperm motility by oral administration of benzene chromatographic fraction of the chloroform extract of the seeds of *Carica papaya* in rats. *Phytomedicine*, 2000; 7: 325-333.
21. WHO: Protocol CG-40, Preparation of Alcoholic Extract for Bioassay and Phytochemical Studies. (APJF/IP, 1001 A). Geneva, World Health Organization; 1983.
22. Natelson S, Total cholesterol procedure Liebermann-Burchard reagent. In: *Techniques of clinicalchemistry*. Charles, C. Thomas publishers, Springfield, Illinois, USA, 3rd edn., 1971; p.263.
23. Bergmeyer H U, *Methods of enzymatic analysis*, H U Bergmeyer (ed) Verlag chemie Weinheim, Academic press, Inc., New York, 1974; vol.1.
24. Kraemer F B, Shen W J, Harada K, Patel S, Osuga J, Ishibashi S, Azhar S, Hormone-sensitivelipase is required for highdensity lipoprotein cholesteryl ester-supported adrenal steroidogenesis, *Mol Endocrinol*, 2004; 18:549-557.
25. Payne A H, YoungbloodGL, Regulation of expression of steroidogenic enzymes in Leydig cells. *Biol Reprod*, 1995; 52:217-225.
26. Jurgens M H, Blunn C T and Peo E R, JR. Vitamin D2 and Cholesterolemia in the Growing Rat *J. Nutr.*, 1971; 101: 153-160.
27. Litscher E S, Williams Z, Wassarman P M, Zona pellucida glycoprotein ZP3 and fertilization in mammals. *Mol Reprod Dev*, 2009; 76: 933-41.
28. Changamma C, Reddanna P, Effect of antifertility agents on steroidogenesis of the albino rat testis *Adv. Contracept. Deliv. Syst.* 1985; 2: 236.
29. Changamma C, Reddanna P, Effect of Antispermatic agents on the kinetic studies of the enzyme 3 β - hydroxy steroid dehydrogenase. *Ind J Expt Biol*, 1987; 25: 506-509.
30. Stoffel-Wagner B, Watzka M, Steckelbroeck S, Schramm J, Bidlingmaier F and Klingmuller D, Expression of 17 β - hydroxysteroid dehydrogenase types 1, 2, 3 and 4 in the human temporal lobe *Journal of Endocrinology*, 1999; 160: 119-126.
31. Emeruwa A C, Antibacterial substance from *Carica papaya* fruit extract. *Journal of Natural Products*, 1982; 45: 123-127.
32. Das R P, Effect of papaya seeds on the genital organs and fertility of male rats. *Ind J Expt Biol*, 1980; 18: 408-409.