

## EVALUATION OF PLANT AND CALLUS EXTRACTS OF *JUSTICIA GENDARUSSA* BURM. F. FOR PHYTOCHEMICALS AND ANTIOXIDANT ACTIVITY

BHAGYA N, K. R. CHANDRASHEKAR\*

Department of Applied Botany, Center of Excellence, Mangalore University, Mangalagangothri, Mangalore 574199, Karnataka, India  
Email: konambi@yahoo.com

Received: 16 Oct 2012, Revised and Accepted: 15 Jan 2013

### ABSTRACT

The *in vitro* grown stem and leaf derived calli of *Justicia gendarussa* Burm. f. obtained on solid MS medium supplemented with NAA+BAP (1+0.1mg/l) were cultured on solid and liquid MS medium for 30-35days. The, stem, leaf and their respective calli samples were extracted in ethanol, methanol and ether. A significantly highest on par concentrations of 183.33 ± 57.73mg/g of phenolics was observed in the solid grown stem callus extracted in methanol and solid grown leaf callus extracted in ethanol. Ethanol found to be a better solvent for the extraction of flavonoids with a highest on par concentrations of 17.00 ± 7.31mg/g, 17.39 ± 1.50mg/g and 14.23 ± 0.28mg/g in liquid grown leaf callus, solid grown leaf callus and liquid grown stem callus respectively. Highest DPPH scavenging activity of methanol extracts of stem derived callus cultured on solid medium was observed at the concentrations of 145.00 ± 5.00µg/ml. The highest reductive capacity was observed for ethanol extract of stem derived callus cultured on solid medium (136.66 ± 2.88mg AAE/g of dried extract).

**Keywords:** *Justicia gendarussa*, *In vitro* culture, Phenolics, Flavonoids, Antioxidant activity

### INTRODUCTION

The over production of reactive oxygen (ROS) and nitrogen species (RNS) in the system damage to DNA, lipids and proteins which is associated with an increased risk of cardiovascular disease, cancer and other chronic diseases[1]. The natural antioxidants of late, considered to be important due to their fewer side effects are the compounds that can neutralize free radicals by donating hydrogen thus, reduce the risk of vascular diseases, some forms of cancer and oxidative stress responsible for DNA, protein and membrane damage[2]. Antioxidant properties of sesame (*Sesamum indicum*) cake extract used in the stabilization of sunflower and soybean oils[3], antioxidant and free radical scavenging activity of Korean medicinal plants[4] due to the presence of phenolics and flavonoids were reported. The phytochemicals produced in the *in vitro* cultured cells were also reported to show the antioxidant properties in *Tecoma stans* (L.) Juss. ex Kunth.[5], *Brassica nigra* L.[6], *Gardenia jasminoides* Ellis[7].

*Justicia gendarussa* Burm. f. (Acanthaceae) is well known for its medicinal properties such as larvicidal, antioxidant, immunosuppressive, antinociceptive, ametic, anti-arthritis and anti-inflammatory activities[8-15]. The plant was also reported to contain phenolics, alkaloids and sterols[16-20]. The present work is aimed at the maintenance of stem and leaf derived callus on solid and liquid nutrient media, extraction of phytochemicals in different solvents and determination of phenolics, flavonoids and antioxidant activity followed by comparison with the plant samples.

### MATERIALS AND METHODS

#### Plant material

*Justicia gendarussa* was collected from natural forests of Dakshina Kannada District, Karnataka, India, and identified following the Flora of Udipi and Dakshin Kannada[21] and the voucher specimen (MU/AB/BN-02) were deposited at the herbarium of Department of Applied Botany.

The healthy stem and leaf samples were collected from the plant, cleaned, dried at 40°C and powdered using Cyclotech lab mill for the extraction of phytochemicals.

#### Callus induction and harvesting

The surface sterilization of the stem and leaf samples were carried out as reported by Bhagya and Chandrashekar[22]. The stem and leaf explants were washed thoroughly under running tap water for

20-30min to remove the surface debris and treated with a systemic fungicide Bavistine for 30-45min. The explants were then sterilized with 70% alcohol (2min) and 0.1% HgCl<sub>2</sub> (8 min). After each step, the explants were washed using sterile distilled water. The callus was induced on MS medium with NAA + BAP (1+0.1mg/l) under 25±2°C with 16 hrs of photoperiod and 40.0±3.0µmol m<sup>-2</sup>s<sup>-1</sup> light intensity. The cultures maintained for 320-360d were inoculated to both solid and liquid MS medium. The cells were harvested at the stationary phase i.e. between 30-35<sup>th</sup> day based on the growth analysis (Unpublished data), washed in distilled water and freeze dried.

#### Chemicals and reagents

All the chemicals used were of analytical grade, purchased from Merck, Himedia and SRL. DPPH was purchased from Sigma Aldrich, USA.

#### Extraction and spectrophotometric estimation of phytochemicals

The phenolics and flavonoids found in the stem, leaf, stem derived callus and leaf derived callus was extracted in methanol, ethanol and ether as per Harborne[23]. Total phenolic and total flavonoid contents in the crude extract was determined as per Malic and Singh[24] and Jia *et al*[25] respectively. Free radical scavenging activities of the extracts were determined using 1, 1-diphenyl-2-picryl hydrazyl radical (DPPH) as per Mensor *et al*[26] and the reducing power of the samples by the method of Oyaizu[27].

#### Statistical analysis

The data presented are the average of three replicates and expressed as Mean±SD. The statistical analysis of all the data were carried out using SAS 9.0 version and the treatment means were compared using DMRT at a level of 5% significance.

### RESULTS AND DISCUSSION

#### Spectrophotometric estimation of phenolics

The spectrophotometric estimation of phenolics in the test samples showed a significantly highest on par concentrations of 183.33 ± 57.73mg/g in the solid grown stem callus extracted in methanol and solid grown leaf callus extracted in ethanol (alcohol) followed by 133.33 ± 14.43mg/g and 100.00 ± 0.00mg/g in solid grown stem callus in ethanol (alcohol) and liquid grown stem callus in methanol respectively (Fig. 1).




Fig. 1: Concentration of phenolics in different extracts of *J. gendarussa*

S.E/plot = 21.21, C.V. (%) = 36.26

Values represent the Mean $\pm$ SD. Means with different letters indicate significant differences at 5% level.

SLC = stem callus in liquid medium, SSC = stem callus on solid medium, S = stem, LLC = leaf callus in liquid medium, LSC = leaf callus on solid medium, L = leaf

The concentration of total phenolics accumulated in the cultured tissue was higher in the callus of *Tecoma stans* (L.) Juss. ex Kunth.[5] and in the hypocotyl explants derived callus of *Brassica nigra*[6]. Even in the present study, the spectrophotometric estimations of phenolics in *J. gendarussa* showed the highest concentration in solid grown callus of stem explant extracted in methanol and solid grown callus of leaf extracted in ethanol. Amid *et al.*[28]estimated the phenolic content in the suspension culture from the leaf derived callus of *J. gendarussa* grown in MS medium supplemented with NAA + BAP (1 + 0.5mg/l). The phenolic content of 88mg/g observed by them was very low compared to the total phenolic content of 183.33  $\pm$  57.73mg in the solid grown callus of stem and leaf explants extracted in methanol and ethanol respectively in the present study.

#### Spectrophotometric estimation of flavonoids

Ethanol (alcohol) was a better solvent for the extraction of flavonoids with a highest on par concentrations of 17.00  $\pm$  7.31mg/g, 17.39  $\pm$  1.50mg/g and 14.23  $\pm$  0.28mg/g in liquid grown leaf callus, solid grown leaf callus and liquid grown stem callus respectively (Fig. 2).

Flavonoids were reported from the leaf derived callus of *Saussurea medusa* Maxim.[29] and callus cultures of *Saussurea involucrata*[30]. The cell suspension cultures from leaf, fruit and root explants of Indian Mulberry (*Morinda citrifolia*) in MS medium supplemented with NAA, BAP and Kn led to the increased production of flavonoids[31]. Similarly, in the present study the suspension grown leaf callus, solid grown leaf callus and suspension grown stem callus showed the highest on par concentrations of flavonoids in *J. gendarussa*.




Fig. 2: Concentration of flavonoids in different extracts of *J. gendarussa*

S.E/plot = 2.04, C.V. (%) = 37.29

Values represent the Mean $\pm$ SD. Means with different letters indicate significant differences at 5% level.

SLC = stem callus in liquid medium, SSC = stem callus on solid medium, S = stem,

LLC = leaf callus in liquid medium, LSC = leaf callus on solid medium, L = leaf




Fig. 3: DPPH activity of different extracts of *J. gendarussa*

S.E/plot = 7.01, C.V (%) = 0.45

Values represent the Mean±SD of three experiments. Means with different letters indicate significant differences at 5% level.

1 - Liquid callus of stem; 2 - Solid callus of stem; 3 - Stem; 4 - Liquid callus of leaf; 5 - Solid callus of leaf; 6 - Leaf; A - Ethanol extract; E - Ether extract; M - Methanol extract




Fig. 4: Reducing power assay of different extracts of *J. gendarussa*

S.E/plot = 5.72, C.V (%) = 17.32

Values represent the Mean±SD of three experiments. Means with different letters indicate significant differences at 5% level.

1 - Liquid callus of stem; 2 - Solid callus of stem; 3 - Stem; 4 - Liquid callus of leaf; 5 - Solid callus of leaf; 6 - Leaf; A - Ethanol extract; E - Ether extract; M - Methanol extract

#### Determination of antioxidant activity

Highest DPPH scavenging activity of methanol extracts of stem derived callus cultured on solid medium was observed at the concentration of  $145.00 \pm 5.00 \mu\text{g/ml}$  followed by methanol extract of stem derived callus cultured in liquid medium at the concentration of  $185.00 \pm 8.66 \mu\text{g/ml}$  (Fig. 3).

However, the scavenging activity was lower than the gallic acid and quercetin standards used in the present study. The reducing power assay showed a significant reducing capacity of the extracts and the absorbance of the sample increased with increase in the concentrations. The highest reductive capacity was observed for ethanol extract of stem derived callus cultured on solid medium ( $136.66 \pm 2.88 \text{ mg AAE/g}$  of dried extract) followed by methanol extracts of stem derived callus cultured on solid medium ( $96.66 \pm 5.77 \text{ mg AAE/g}$  of dried extract) (Fig. 4).

The higher antioxidant activity of the solid grown callus samples of *J. gendarussa* observed in the present study may be due to the type and concentrations of chemical components produced *in vitro*. In solid medium the cultures will grow under stress compared to suspension grown callus in terms of nutrient and oxygen supply. The present study also confirmed the presence of phenolics and flavonoids in the test samples. All these compounds are responsible for the antioxidant properties as reported by Dimitrios[2], Jayaraj and Punja[32]. Amid *et al*[28] reported the highest antioxidant activity of  $8.46 \times 10^{-3} \text{ mg/ml}$  ( $8460 \mu\text{g}$ ) in the leaf derived callus of *J. gendarussa* cultured in liquid medium supplemented with NAA + BAP ( $1 + 0.5 \text{ mg/l}$ ). However, in the present study, the antioxidant activity of *J. gendarussa* illustrated a higher scavenging activity at  $145.00 \pm 5.00 \mu\text{g/ml}$  in methanol extract of stem derived callus cultured on solid medium. Uddin *et al*[20] reported the antioxidant activity ( $\text{IC}_{50} = 18.8 \mu\text{g/ml}$ ) in chloroform fraction of whole plant

extract of *J. gendarussa*. The enhanced production of phenolics content and free radical scavenging activity was reported in gamma irradiated leaves of *Justicia adhatoda*[33]. Methanol extract of stem and leaf samples of 2 months old *in vitro* grown seedlings of *Lycium barbarum* (goji) showed the highest antioxidant activity[34]. The antioxidant activity of *Justicia spicigera* and also the presence of phenolics and flavonoids in the aqueous and methanolic extracts of the leaf, stem and flowers were reported by Sepulveda-Jimenez *et al*[35]. However, in the present study, the scavenging activity was higher in ethanol and methanol extracts of solid grown stem callus (350.00±86.60µg/ml and 446.67±50.33µg/ml).

The *in vitro* culturing of stem and leaf derived calli of *J. gendarussa* on the solid and liquid medium for the production of bioactive compounds and a comparative study of the concentrations produced *in vitro* with their respective plant parts is the first report. The phytochemicals detected in the present study may have their role as antidiabetic, anti-inflammatory, hepatoprotective, anticarcinogenic agents as these properties of *J. gendarussa* were already reported from various laboratories as mentioned before. The present study forms an alternative method for the extraction of bioactive compounds from the *in vitro* grown cultures and also to isolate and characterize many more bioactive compounds from the cultures.

#### ACKNOWLEDGMENTS

The authors are grateful to VGST, DBT, Govt. of Karnataka for financial assistance. Senior author (BN) is grateful to UGC, New Delhi for financial assistance under RFSMS scheme.

#### REFERENCES

- Dimitrios B. Sources of natural phenolic antioxidants. Trends Food Sci Tech 2006; 17: 505-512
- Ferreira ICFR, Paula B, Miguel V-B, Lillian B. Free-radical scavenging capacity and reducing power of wild edible mushrooms from northeast Portugal: Individual cap and stipe activity, Food Chem 2007; 100: 1511-1516
- Mohdaly AAA, Smetanska I, Ramadan MF, Sarhan MA, Mahmoud A. Antioxidant potential of sesame (*Sesamum indicum*) cake extract in stabilization of sunflower and soybean oils, Ind Crops Prod 2011; 34: 952-959
- Choi CW, Kim SC, Hwang SS, Choi BK, Ahn HJ, Lee MY, Park SH, Kim SK. Antioxidant activity and free radical scavenging capacity between Korean medicinal plants and flavonoids by assay-guided comparison. Plant Sci 2002; 163: 1161-1168
- Lopez-Laredo AR, Ramirez-Flores FD, Sepulveda-Jimenez G, Trejo-Tapia G. Comparison of metabolite levels in callus of *Tecoma stans* (L.) Juss. ex Kunth. cultured in photoperiod and darkness. In Vitro Cell Dev Biol - Plant 2009; 45: 550-558
- Hussein EA, Taj-Eldeen AM, Al-Zubairi AS, Elhakimi AS, Al-Dubaie AR. Phytochemical screening, total phenolics and antioxidant and antibacterial activities of callus from *Brassica nigra* L. hypocotyl explants. Int J pharmacol 2010; 6: 464-471
- Sayd SS, Taie HAA, Taha LS. Micropropagation, antioxidant activity, total phenolics and flavonoids content of *Gardenia jasminoides* Ellis as affected by growth regulators. Int J Acad Res 2010; 2: 184-191
- Senthilkumar N, Varma P, Gurusubramanian G. Larvicidal and adulticidal activities of some medicinal plants against the malarial vector, *Anopheles stephensi* (Liston). Parasitol Res 2009; 104: 237-244.
- Mruthunjaya K, Hukkeri VI. Antioxidant and free radical scavenging potential of *Justicia gendarussa* Burm. Leaves *in vitro*. Nat Prod Rep 2007; 13: 199-206.
- Arokiyaraj S, Perinbam K, Agastian P, Balaraju K. Immunosuppressive effect of medicinal plants of Kolli hills on mitogen stimulated proliferation of the human peripheral blood mononuclear cells *in vitro*. Indian J Pharmacol 2007; 39:180-183.
- Ratnasooriya WD, Deraniyagala SA, Dehigaspitiya DC. Antinociceptive activity and toxicological study of aqueous leaf extract of *Justicia gendarussa* Burm.f. in rats. Phcog Mag 2007; 3: 145-155
- Prajogo BEW, Ayun S, Soedjak N. The influence of 60% ethanol fraction and water fraction of *Justicia gendarussa* Burm.f leaf on pegan (*Columba livia*) emetic activity, Vol: 1.Veterinaria Medica; 2008.
- Paval J, Kaitheri SK, Potu BK, Kumar RS, Narayanan SN, Moorkoth S. Anti-arthritis potential of the plant *Justicia gendarussa* Burm F. Clinics (Sao Paulo) 2009; 64: 357-362
- Shikha P, Latha PG, Suja SR, Anju GI, Shymal S, Shine VJ, Sini S, Kumar NMK, Rajasekharan S. Anti-inflammatory and antinociceptive activity of *Justicia gendarussa* Burm. F. leaves. Indian J Nat Prod Resources 2010; 1: 456-461
- Jothimanivannan C, Kumar RS, Subramanian N. Anti-inflammatory and analgesic activities of ethanol extract of aerial parts of *Justicia gendarussa*. Int J Pharmacol 2010; 1-6
- Wahi SP, Wahi AK, Kapoor R. Chemical study of the leaf of *Justicia gendarussa* Burm. JRIM 1974; 9: 65-66
- Oliveira AFM, Xavier HS, Silva NH, Andrade LHC. Chromatographic screening of medicinal Acanthaceae: *Justicia pectoralis* Jacq. and *J. gendarussa* Burm. Revista Brasileira de Plantas Mediciniais. 2000; 3: 37-41
- Gulfraz M, Arshad M, Nayyer N, Kanwal N, Nisar U. Investigation for bioactive compounds of *Berberis lyceum* Royle and *Justicia adhatoda* L. Ethnobotanical leaflets 2004; opensiuc.lib.siu.edu/cgi/viewcontent.cgi?article=1268 & context.
- Bachheti RK, Pandey DP, Joshi A, Rana V. Chemical analysis of aerial parts of *Justicia gendarussa*. Int J Chem Tech Res CODEN( USA) 2011; 3: 244-247
- Uddin MdR, Sinha S, Hossain MdA, Kaiser MdA, Hossain MdK, Rashid MA. Chemical and biological investigations of *Justicia gendarussa* (Burm. F), Dhaka Univ J Pharm Sci 2011; 10: 53-57
- Bhat KG. Flora of Udipi, Indian Naturalist (Regd.), Udipi, Karnataka, India. 2003
- Bhagya N, Chandrashekar KR. Effect of auxin concentration on callus induction from *Justicia gendarussa* L. stem and leaf explants. IJBST 2010; 3: 27-35
- Harborne JB. Phytochemical methods, A guide to modern techniques to plant analysis. Chapman and Hall (Pub), USA; 1984
- Malic CP, Singh MB. In: *Plant enzymology*, Kalyani Publishers, New Delhi;1980
- Jia Z, Tang M, Wu J. The determination of flavonoid content in mulberry and scavenging effect on superoxide radicals. Food Chem 1999; 64: 555-559
- Mensor LI, Menezes FS, Leitao GG, Reis AS, Dos ST, Coube CS, Leitao SG. Screening of Brazilian Plant extracts for antioxidants activity by the use of DPPH free radical method. Phytother Res 2001; 15: 127-130
- Oyaizu M. Studies on product of browning reaction prepared from glucose amine. Jpn J Nutr 1986; 44: 307-315.
- Amid A, Johan NN, Jamal P, Zain WNW Mohd. Observation of antioxidant activity of leaves, callus and suspension culture of *Justicia gendarussa*. Afr J Biotechnol 2011; 10: 18653-18656
- Liu CZ, Saxena PK. *Saussurea medusa* cell suspension cultures for flavonoid production. Methods Mol Biol 2009; 547: 53-59.
- Wu LQ, Guo SX, Xiao PG. Cell suspension culture and flavonoids production in *Saussurea involucrate*. Zhongguo Zhong Yao Za Zhi 2005; 30: 965-8.
- Deshmukh SR, Wadegaonkar VP, Bhagat RP, Wadegaonkar PA. Tissue specific expression of Anthraquinones, flavonoids and phenolics in leaf, fruit and root suspension cultures of Indian Mulberry (*Morinda citrifolia* L.). POJ 2011; 4: 6-13
- Jayaraj, Punja ZK. Transgenic carrot plants accumulating ketocarotenoids show tolerance to UV and oxidative stresses. Plant Physiol Biochem 2008; 46: 875-883.
- Nilima S, Rajurkar, Gaikwad KN, Razavi MS. Evaluation of free radical scavenging activity of *Justicia adhatoda*: a gamma radiation study. IJPPS 2012; 4 (Suppl 4): 93-96
- Osman NI, AWALA and SidikNJ, AbdullahS. Antioxidant activities of *in vitro* seedlings of *Lycium barbarum* (goji) by diphenyl picrylhydrazyl (DPPH) assay. IJPPS 2012; 4 (Suppl 4): 137-141
- Sepulveda-Jimenez G, Reyna-Aquino C, Chaires-Martnez L, Bermudez-Torres K, Rodriguez-Monroy M. Antioxidant activity and content of phenolic compounds and flavonoids from *Justicia spicigera*. J Biol Sci 2009; 9: 629-632.