

CYTOTOXIC INFLUENCING ACTIVITY OF LATEX OF *EUPHORBIA ANTIQUORAM* LINN

HAMSA D., \*SUMATHI S., SOWMINI C.M. AND PADMA P.R.

Department of Biochemistry, Biotechnology and Bioinformatics, Avinashilingam Institute of Home science and Higher Education for Women University, Coimbatore. Email: sumathi\_vnktsh@yahoo.co.in

Received: 12 Dec 2012, Revised and Accepted: 23 Jan 2013

## ABSTRACT

Cytotoxicity measurements were originally designed for rapid and inexpensive analysis of soluble pharmaceuticals. The brine shrimp lethality assay (BSLA) has been used routinely in the primary screening of the crude extracts as well as isolated compounds to assess the toxicity towards brine shrimp, which could also provide an indication of possible cytotoxic properties of the test materials. Mouse spleen cells have been successfully used to analyze mechanisms of cytotoxicity for a variety of anticancer drugs. The results in BSLA showed that the percentage of lethality rate of the shrimps was found to be less which indicates that at low concentration the survival rate was increased and protected from the effects of test sample. The calculated LC<sub>50</sub> value for the tested sample was 20µg. The percentage of viable cells was assessed by MTT assay in a time and dose dependent manner in mouse spleenocytes and LC<sub>50</sub> value of methanolic extract of latex milk was found to be 20µg at different time intervals. SRB and Neutral red assay showed that latex milk is safe to normal spleen cells.

**Keywords:** Apoptosis, *Euphorbia antiqoram*, Cytotoxicity, Medicinal plants.

## INTRODUCTION

Cancer has become a leading cause of death in the world because changed disease spectrum by social and environmental risk factors. Cancer is a series of malignant disease with multiple pathological stages (e.g. cancer initiation, promotion and progression) and involved in multiple factors (genetic, environmental, biological, chemical, physical and psychological factors). Cancer is characterized by a rapid and uncontrolled formation of abnormal cells, which may mass together to form a growth of tumor, proliferate throughout the body and initiate abnormal growth at other sites[1]. Chemotherapy is one of the important treatment methods, but the side effects are difficult to tolerate, and as a result, people are paying more attention in searching for new antitumor agents from natural products[2]. The use of alternative medicines, especially when used in conjunction with conventional cancer treatments, can serve to mitigate the side effects, enhance the uptake of conventional medicines, and, bolster the immune system to fight the cancer. Since these medicines are primarily extracts of naturally occurring flora, their bioavailability is less likely to induce severe immune responses[3]. *Euphorbia antiquorum* is one of the largest armed trees in Euphorbias, with an average height of 5-7 m, it has been known to attain gigantic proportions if left undisturbed. The odour of its latex is pungent and lingering[4]. The absorbent prepared from *Euphorbia antiquorum* L is an effective absorbent for the removal of coloring matter present in the dyeing industry effluent[5]. Previous studies in our laboratory has shown that, the latex milk of *Euphorbia antiquorum* can be exploited for its antimicrobial activity and found to be safe to normal cells namely *S. cerevisiae* and chick embryo fibroblast cells[6] The present study was designed to study the cytotoxic effect of methanolic extract of latex milk of *Euphorbia antiquorum*.

## MATERIALS AND METHODS

## Plant material

The plant specimen was identified and authenticated (Specimen No 365) by Dr.G.V.S Murthy, Scientist E, Director, Botanical survey of India, Tamilnadu. The latex milk was collected from plant by breaking up the stem of *Euphorbia antiquorum*. The latex milk was collected in the morning hours between 7 to 8 a.m in a glass container and maintained in an ice - cold condition till the use of latex for extraction. The collected latex milk was extracted with methanol. The methanolic extract was prepared by dissolving 1.0ml of latex in 5.0 ml of methanol and allowed to evaporate at 60°C in a water bath. The residue was dissolved in minimal volume of DMSO, stored and used for the assays.

## Cytotoxicity Screening

Two model organisms were used to screen the cytotoxic activity of *Euphorbia antiquorum*.

## 1. Brine shrimp lethality bioassay

*Artemia salina* the brine shrimp is an invertebrate component of the fauna of saline aquatic and marine ecosystem[7]. It plays an important role in the energy flow of the food chain. And it can be used in a laboratory bioassay in order to determine the toxicity. Different concentrations of the latex milk were added and after 24 hours exposure, the live nauplii were counted and LC<sub>50</sub> value was calculated using probit analysis.

## 2. Primary cell culture

Spleenocytes were used as a source of primary cells. Spleenocytes were isolated from three to four weeks old mice and cultured in DMEM with 10% fetal calf serum.

## Parameters analyzed using spleenocytes

The cell proliferation capacity was assessed by MTT assay. The viability of the cells in the presence and absence of etoposide, with or without latex milk at different concentrations and for different time periods was examined by MTT assay as proposed by Igarashi and Miyazawa (2001)[8] The SRB assay provides rapid and sensitive methods for measuring the cytotoxicity and metabolic activity of the cells. The assay was done in the presence and absence of standard chemotherapeutic drug etoposide with or without latex milk proposed by Skehan *et al* [9]. The extent of viability in the treatment group based on the uptake of neutral red dye by viable cells which was then measured photometrically by neutral red assay[10].

## RESULTS AND DISCUSSION

## Brine shrimp lethality test

The results of the brine shrimp lethality after 24 hrs of exposure to the sample were determined. A chart was plotted using the % of lethality in Y-axis and concentrations of the latex milk extract in X-axis. At lower concentration, the percentage of lethality rate of the shrimps was found to be less which indicates that at low concentration the survival rate increased and protected the effects of test sample. The calculated LC<sub>50</sub> value for the tested sample was 20µg.

The *n*-hexane fraction of *Myrsine africana* L. showed good cytotoxicity (66.66%) at 1000 µg/ml, while low activity (33.33 and 13.33%) was observed at 100 and 10 µg/ml, respectively. The LD<sub>50</sub> value calculated was 30.344 µg/ml for *n*-hexane fraction. The chloroform (CHCl<sub>3</sub>) extract showed low activity at all concentrations 30.0, 16.66 and 13.33% at 1000, 100 and 10 µg/ml respectively. The ethyl acetate (EtAc) fraction showed low brine shrimp lethality at respective test concentrations that was: 26.66, 20.00 and 13.33% at 1000, 100 and 10 µg/ml, respectively[11]. These findings are in agreement with our results and the LC<sub>50</sub> was found to be 20µg/ml




Fig. 1: Brine shrimp lethality test (BSLT)

**Cytotoxicity Testing**

The percentage of viability of splenocytes was assessed using different concentrations of plant extracts and also at different time intervals namely 12, 24, 36 and 48 hrs. We checked the cytotoxicity in a time and dose dependent manner. The 24 hours incubation time showed maximum percentage of cell survival and LC<sub>50</sub> value of methanolic extract of latex milk was found to be 20 µg at different time intervals using probit analysis in SPSS statistical package.

Studies reported by other workers have shown that the potency of methanolic extract of *Euphorbia hirta* calculated in terms of percent decrease in viable Hep-2 cells as compared to the control value. The extract showed dose dependent antitumor activity. The MTT assay showed an anti proliferative activity (IC<sub>50</sub>) at 625µg/ml of crude extract[12]. These studies support our findings. Cytotoxicity was also tested in the presence of standard chemotherapeutic drug etoposide.




Fig. 2: Dose - Dependent effect of *Euphorbia antiquorum* latex milk on proliferation of splenocytes by MTT assay




Fig. 3: Dose - Dependent effect of *Euphorbia antiquorum* latex milk on proliferation of splenocytes co treated with etoposide by MTT assay. E = Etoposide

### SRB Assay

The results obtained for the effect of etoposide on cytotoxicity and co treatment with latex milk, as determined by the SRB assay, showed that the viability of the splenocytes decreased in the etoposide treated group. Only a slight reduction in the viability was observed in the group treated with latex milk extract alone, which showed that the latex milk extract by itself was not toxic to normal cells. However an improvement in viability was observed in the spleen cells treated with etoposide as

well as the latex milk. This improvement must be due to the protective effect of latex milk extract against etoposide. Several authors have reported their findings in tune with our studies.

The percentage growth inhibition was found to be increasing with increasing concentration of test compounds of *Solanum Nigrum* on HeLa cell line up to 0.0196 mg/ml and the  $IC_{50}$  value of *Solanum Nigrum* on HeLa cell and Vero cell were 847.8 and 0.8724, 0.9088 and 0.1017 respectively by SRB assay[13].




Fig. 4: Effect of latex milk of *Euphorbia antiquorum* linn. on the cell survival of splenocytes treated with etoposide by SRB assay

The values are mean of duplicates. The control was fixed as having 100% cell survival and the percent survival in the other groups was calculated.

### Neutral Red Assay

Neutral red assay, which is based on the uptake and accumulation of the supravital dye neutral red by viable cells. The result obtained in the neutral red assay showed that etoposide treatment caused a marked decrease in the viability of the cells and the co-administration of latex milk extract increases the cell survival. The extent of cell survival in the latex milk extract alone treated group was comparable to that of control cells suggesting that the latex milk of *Euphorbia antiquorum* was not toxic to the spleen cells. Similar

results have been reported that the cytotoxic effects of the crude methanol and fractionated extracts (hexane, ethyl acetate) of *C. mangga* against six human cancer cell lines, namely the hormone-dependent breast cell line (MCF-7), nasopharyngeal epidermoid cell line (KB), lung cell line (A549), cervical cell line (Ca Ski), colon cell lines (HCT 116 and HT-29), and one non-cancer human fibroblast cell line (MRC-5) by neutral red cytotoxicity assay. The crude methanol and fractionated extracts (hexane and ethyl acetate) displayed good cytotoxic effects against MCF-7, KB, A549, Ca Ski and HT-29 cell lines, but exerted no damage on the MRC-5 line[14].




Fig. 5: Effect of latex milk of *Euphorbia antiquorum* on the viability of splenocytes treated with etoposide as determined by the Neutral red assay

The values are mean of duplicates. The control was fixed as having 100% cell survival and the percent survival in the other groups was calculated relative to this.

## CONCLUSION

Methanol extract of latex milk of *Euphorbia antiquorum* was tested for the cytotoxic and apoptotic activity using brine shrimp and mice spleenocytes and it showed latex milk is safe to brine shrimp and normal spleen cells in lower concentration. The need of the present study is to develop drugs that can potentially target cancer cells by means of their inherent difference to normal cells. The development of such drugs with differential action will be very valuable in cancer chemotherapy without the observed side effects. These findings potentiate the use of latex milk of *Euphorbia antiquorum* in combination therapy when this extract can overcome the toxic side effect of chemotherapeutic drug, because it is safe to normal cells. If its anticancer activity is probed further using cancer cell lines we can see if it exhibits a differential response to normal cells and cancer cells. The potential use of *Euphorbia antiquorum* as therapeutic agent holds great promise as the isolation of one or more cytotoxic chemicals from crude latex milk extract and the judicious use of such chemicals can control the progression of cancer and also can prevent the formation of tumor in individuals who are highly susceptible to developing a tumor.

## REFERENCES

1. Feng Y, Zhu W, Feng L and Tsao S. *Recent progress on anticancer candidates in patents of herbal medicinal products*, Recent Patents on Food, Nutrition and Agriculture 2011, 3:30-48.
2. Dai Y, Theodore S, Lawrence and Xu L. *Overcoming cancer therapy resistance by targeting inhibitors of apoptosis proteins and nuclear factor kappa B*, Am J. Trans. Res 2010; 1:1-15.
3. Chan LL, George S, Ahmad I, Gosangari, SL, Abbasi A, Cunningham BT and Watkin, KL. *Cytotoxicity effects of Amoora rohataka and chittagonga on breast and pancreatic cancer cells*, Evidence-Based Complementary and Alternative Medicine 2011; 860605:1- 8.
4. Sivakumar P and Palanisamy PN. *Adsorption studies of basic red 29 by a nonconventional activated carbon prepared from Euphorbia antiquorum*, Int.J. Chem Tech Res 2009; 1:502-510.
5. Sivakumar P, Palanisamy PN, Hameed BH and Radha K. *Novel non-conventional activated carbon for the remediation of dyeing industry effluent*, International Journal of Civil and Environmental Engineering 2011 ;3:45-50.
6. Sumathi S, Malathy N, Dharani B, Sivaprabha J, Hamsa D, Radha P and Padma PR. *Cytotoxic studies of latex of Euphorbia antiquorum in in vitro models*, Journal of medicinal plant research 2011; 5, 4715-4720
7. Ramachandran S, Vamsikrishna M, Gowthami KV, Heere B and Dhanaraju MD. *Assessment of cytotoxic activity of Agave cantula using brine shrimp (Artemia salina) lethality bioassay*, Asian J. Scientific Research 2011; 4:90-94.
8. Igarashi M. and Miyazawa T. *The growth inhibitory effect of conjugated linolenic acid on a human hepatoma cell line HepG 2 is induced by a change in fatty acid metabolism but not the facilitation of lipid peroxidation in cells*, Biochem. Biophys. Acta/Mol.Cell Biol. Lipids 2011; 1530:162-171.
9. Skehan P, Storeng R, Scudiero D, Monks A, Mahan J, Vistica D, Warren JT, Bokesch, H, Kenney S and Boyol M.R. *New colorimetric cytotoxicity assay for anticancer drug screening*, J. Natl. Can. Inst, 1990; 82: 1107-1112.
10. Borenfreund E, Babich H, Martin-alguail N. *Rapid chemo sensitivity assay with normal and tumor cells in vitro*. In vitro cell. Dev. Biol, 2011; 26:1030-1034.
11. Ahmad B, Azam S, Bashir S, Hussain F and Chaudhary MI. *Insecticidal, brine shrimp cytotoxicity, antifungal and nitric oxide free radical scavenging activities of the aerial parts of Myrsine africana L.*, African Journal of Biotechnology 2011; 10:1448-1453.
12. Sidambaram, R., Dinesh, MG and Jayalakshmi, ET. *An in vitro study of cytotoxic activity of Euphorbia hirta on Hep2 cells of human epithelioma of larynx*, International Journal of Pharmacy and Pharmaceutical Sciences 2011; 3, 100-103.
13. Patel S, Gheewala N, Suthar A and Shah A. *In-vitro cytotoxicity activity of Solanum nigrum extract against HeLa cell line and vero cell line*, International Journal of Pharmacy and Pharmaceutical Sciences 2009; 1: 38-46.
14. Malek SNA, Lee GS, Hong SL, Yaacob HL, Wahab NA, Faizal J, Weber F and Shah SA. *Phytochemical and cytotoxic investigations of Curcuma mangga rhizomes*, Molecules 2011; 16: 4539-4548.