

## AMELIORATING EFFECT OF LAWSONE AND ALPHA-PINENE ON L-ARGININE INDUCED ACUTE PANCREATITIS IN RATS

SANDEEP BIRADAR\* & B VEERESH

Department of Pharmacology, G. Pulla Reddy College of Pharmacy, Mehdipatnam, Hyderabad 500 028, India.  
Email: sandeepbiradar84@gmail.com

Received: 02 Mar 2013, Revised and Accepted: 20 Apr 2013

### ABSTRACT

**Objective:** This study is intended to study the protective effect of Lawsone and Alpha-pinene on L-Arginine induced acute pancreatitis in rats.

**Method:** In this study lawsone and alpha-pinene against L-arginine induced acute pancreatitis was determined at 24 h by determination of serum levels of amylase, lipase and proinflammatory cytokines [tumor necrosis factor (TNF)- $\alpha$ , C-reactive proteins and interleukin (IL)], pancreatic myeloperoxidase (MPO) activity, lipid peroxidation (thiobarbituric acid reactive substances (TBARS)), nitrate/nitrite levels, and the wet weight/body weight ratio.

**Result:** Lawsone, alpha-pinene and methylprednisolone treatments significantly attenuated the L-arginine-induced increases in pancreatic wet weight/body weight ratio, and decreased the serum levels of amylase and lipase, and TNF- $\alpha$  and IL-6 and significantly lowered pancreatic levels of MPO, TBARS, and nitrate/nitrite. The histopathological findings further proved the amelioration of pancreatic injury by lawsone and alpha-pinene. Further it also proved anti-inflammatory and antioxidant agent property of lawsone and Alpha-pinene.

**Conclusion:** This study had proved ameliorating effect of Lawsone and alpha-pinene on L-Arginine induced acute pancreatitis in rats.

**Keywords:** Acute pancreatitis, Anti-inflammatory, Cytokines, L-arginine, Lawsone, Pinene

### INTRODUCTION

Inflammation of pancreatic gland called pancreatitis (AP) may lead to severe complication and high mortality without treatment. The pathogenesis is not fully understood, however the leukocyte activation, microcirculatory disturbances and oxidative stress are the major constituents of AP. This is characterized by activation of widespread inflammatory cell infiltration, leukocyte and digestive proteases. Reactive oxygen, nitrogen species and various kinds of inflammatory mediators are released in inflammatory process. Previously it was reported that several factors are responsible for the AP, like alcohol, gallstones, hereditary pancreatitis, hypercalcemia, hyperlipidemia, malnutrition, abdominal trauma, penetrating ulcers, malignancy, drugs like steroids, sulfonamides, furosemide, thiazides, infections like mumps, Coxsackie Virus, Mycoplasma Pneumoniae, Ascaris, Clonorchis, and structural abnormalities like choledochocoele and pancreas divisum [1,2]. Repeated attacks of acute pancreatitis have the potential to develop into chronic pancreatitis or pancreatic cancer characterized by fibrosis and loss of acinar cell function [1,2]. No specific treatment is available to treat AP. Many therapies and medical management is aimed to control the sign and symptoms of AP, using steroids, analgesics and anti-inflammatory agents. The use of the synthetic and semi-synthetic treatment has various kinds of drawbacks like photosensitivity skin reactions, intolerance and addiction. Apart from this, these compounds are very expensive and not reliable. Hence, there is need to explore potential antioxidant and anti-inflammatory agents available from natural sources, which are cost effective and have several advantages than the synthetic and semisynthetic compounds. Anti-oxidant, anti-inflammatory and anti-carcinogenic activities [3,4] of alpha-pinene and anti-oxidant, anti-inflammatory and anti-carcinogenic activities [5,6] of lawsone have been reported. The alpha-pinene has shown to inhibit nuclear translocation of Nuclear Factor Kappa B [4] and lawsone has haematotoxic properties, leading to stimulation of cell proliferation; representing a sufficient explanation for a weak induction of 'late micronuclei' [6]. Hence in this study the non-toxic dose of alpha-pinene and lawsone (100 and 200 mg/kg) were used to evaluate the potential effect to ameliorate pancreatic injury induced by L-arginine.

### MATERIALS AND METHODS

**Animals:** Male Wistar rats (42) weighing 180-200 g obtained from Mahaveer Enterprises, Hyderabad were maintained at a constant

room temperature ( $23 \pm 2$  °C) with 12:12 h light-dark cycles and free access to water and standard laboratory chow. The rats were randomly divided into 7 groups of 6 in each and experiments were performed after 12 h of fasting. All the experimental procedures were carried out in accordance with Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) guidelines. The study was reviewed and approved by the Institutional Animal Ethics Committee (320/CPCSEA dated 03-01-2001), G. Pulla Reddy College of Pharmacy, Hyderabad, India.

**Chemicals:** L-arginine (Sigma Aldrich Co Pvt Ltd, USA), lawsone (Sigma Aldrich Co Pvt Ltd, Japan), alpha-Pinene (Destilaciones Bordas Chinchurreta SA), hexa-decyl-tri-methylammonium bromide (HETAB) (Sigma Aldrich Co Pvt Ltd, Switzerland), o-dianisidinedihydrochloride, thiobarbituric acid (TBA) (Sigma Aldrich Co Pvt Ltd, Germany), Griess reagent (Sigma Aldrich Co Pvt Ltd, Germany) and vanadium trichloride (Sigma Aldrich Co Pvt Ltd, USA) were procured from Sigma Aldrich Chemical Co. All other chemicals and reagents were of highest commercial grade available locally.

L-arginine, powder: prepared as a solution by dissolving in 0.9% saline to a final concentration of 500 mg/mL and the pH was adjusted to 7 with 5 N HCl.

Lawsone was prepared as a solution by dissolving in 3% tween 80 and 0.9% saline to a final concentration of 100 and 200 mg/mL and the pH was adjusted to 7 with 0.1 N NaOH.

Pinene was prepared as a solution by dissolving in 3% tween 80 and 0.9% saline to a final concentration of 100 and 200 mg/mL and the pH was adjusted to 7 with 0.1 N NaOH.

L-arginine-induced pancreatitis model—Acute pancreatitis was induced in five groups of rats by two intraperitoneal (ip) injections of L-arginine (2.5 g/kg, 1 h apart). One hour following the last injection of L-arginine, the rats were treated orally as follows: Gr. 1 received the vehicle (3% Tween 80) of lawsone (vehicle control); Grs 2 and 3 were treated with lawsone (100 and 200 mg/kg, respectively), Gr 4 and 5 were treated with pinene (100 and 200 mg/kg, respectively), Gr. 6 acted as positive control and received methylprednisolone (30 mg/kg), all in a volume of 10 mL/kg and Gr. 7, received saline (0.9%, NaCl, ip) in place of L-arginine and served as a normal control. After 24 h of the last

injection of L-arginine or saline, a midline laparotomy was performed in rats under ether anaesthesia and blood samples were collected from the inferior vena cava, the rats were then exsanguinated, the whole pancreas was quickly removed and stored at -70 °C until use. The pancreatic weight/body weight ratio was evaluated as an estimate of the degree of pancreatic edema (mg/g).

#### Macroscopic evaluation

**Pancreas weight/body weight ratio**—The pancreas was removed immediately after the blood collection, trimmed free of fat and weighed. The pancreatic weight/body weight ratio (mg/g) was calculated for each animal, to estimate the level of pancreatic edema [7].

**Serum analysis**—For serum analysis, blood samples were centrifuged at 3000 g at 4 °C for 10 min. The serum amylase and lipase were determined by routine colorimetric methods using the commercial kits for amylase (Rapid diagnostics), lipase (Accurex diagnostics), C-reactive protein and interleukin- $\alpha$  and expressed as U/dl [7].

**Biochemical estimations**—Pancreatic total protein content [8] was determined. Myeloperoxidase activity [7], nitrate/nitrite level [9,10], TBARS level [11], catalase activity [12,13], SOD level [14] and reduced GSH level [14] were measured.

**Histopathological evaluation**—Pancreas was removed immediately and part of it was fixed in 10% neutral buffered formalin and embedded in paraffin by standard methods. Paraffin sections of 5  $\mu$ m thicknesses were cut and stained with haematoxylin and eosin, assessed under dark field microscope and examined blind by a morphologist for grading histopathological changes. Pancreatic damage was assessed and scored by grading acinar cell degeneration, interstitial inflammation, edema, and haemorrhage as described by Schmidt's standards [15,16] with modification as follows: Grading for edema was scaled as 0: absent or rare; 1: edema in the interlobular space; 2: edema in the intralobular space; 3: isolated island shape of pancreatic acinus. Inflammation was scored as 0: absent; 1: mild; 2: moderate; 3: severe. Acinar cell necrosis was scored as 0: absent; 1: mild; 2: moderate; 3: severe. Parenchyma haemorrhage was scored as 0: absent; 1: mild; 2: moderate; 3: severe. The maximum score for acinar cell damage was 12.

**Statistical analysis**—Statistical analysis was performed by one way ANOVA followed by Newman Keuls as post-hoc test using Graph pad

Prism 5. Values were presented as mean  $\pm$  SE. The difference was considered to be statistically significant when  $P < 0.05$ .

#### RESULTS

**Serum biochemical parameters and pancreatic edema**—Induction of pancreatitis resulted in significant raise in the serum amylase, lipase and pancreatic edema. Treatment with lawsone and pinene (100 and 200 mg/kg) dose dependently decreased the serum amylase, lipase and pancreatic edema (Table 1).

**Pancreatic, lung, liver and kidney MPO and total protein**—Induction of pancreatitis resulted in significantly increased the pancreatic MPO and decreased the pancreatic total protein levels. Treatment with lawsone and pinene (100 and 200 mg/kg) dose dependently reversed the change in pancreatic MPO and total protein levels (Table 1).

**Pancreatic, lung, liver and kidney MDA, nitrate/nitrite, GSH and antioxidant enzymes catalase and SOD**—Induction of pancreatitis resulted in a significant raise in MDA, nitrate/nitrite, catalase and SOD and decline in GSH levels. Treatment with lawsone and pinene (100 and 200 mg/kg) dose dependently reversed the change in MDA, nitrate/nitrite, catalase, SOD and GSH levels (Table 1).

**Assessment of interleukins, TNF-  $\alpha$  and C- reactive protein**—Induction of pancreatitis resulted in a significant raise in interleukins, TNF-  $\alpha$  and C- reactive protein. Treatment with lawsone and pinene (100 and 200 mg/kg) dose dependently decreased the interleukins and C- reactive protein (Table 2).

**Pancreatic histology**—Histological examination of normal control group (saline treated) showed normal architecture and absence of edema, neutrophil infiltration, hemorrhage and necrosis (Fig 1). Whereas, pancreatic sections of disease control group showed extensive tissue damage characterized by acinar cell degeneration, necrosis, edema, mononuclear cell infiltration, hemorrhage and thus received significantly higher scores. Treatment with lawsone (100 and 200 mg/kg), pinene (100 and 200 mg/kg) and methyl prednisolone (30 mg/kg) ameliorated the inflammation, edema and more significantly acinar cell degeneration and necrosis and protected the pancreas from L-arginine induced damage. Treatment with lawsone dose dependently decreased the total pathological scores compared to disease control group.

**Table 1: Effect of alpha-pinene and Lawsone on pancreas weight, total body weight, serum amylase, serum lipase, total nitrate, total protein, MDA, MPO and SOD after L-arginine induced acute pancreatitis.**

Parameter/Groups	N.C	D.C	STD	LW 100	LW 200	PIN 100	PIN 200
Pancreas weight	870.3 $\pm$ 15.36	1015 $\pm$ 19.0*	911.3 $\pm$ 18.83 $\alpha$	843.3 $\pm$ 46.33 $\alpha$	788.3 $\pm$ 21.37 $\alpha$	862 $\pm$ 83.67 $\alpha$	869 $\pm$ 80.17 $\alpha$
Total body wt	187.5 $\pm$ 4.889	192 $\pm$ 3.578*	191.2 $\pm$ 3.60	188.7 $\pm$ 5.57	189 $\pm$ 3.68	188 $\pm$ 18.85	174 $\pm$ 18.7
Pancreatic ind (x10-3)	4.652 $\pm$ 1.26	5.286 $\pm$ 1.57	4.769 $\pm$ 1.32	4.48 $\pm$ 1.28	4.169 $\pm$ 1.22	4.58 $\pm$ 1.28	4.99 $\pm$ 1.24
Serum Amylase	2000 $\pm$ 85.63	7667 $\pm$ 349*	3317 $\pm$ 110.8 $\alpha$	2733 $\pm$ 140.6 $\alpha$	3167 $\pm$ 187.4 $\alpha$	2933 $\pm$ 527.9 $\alpha$	3200 $\pm$ 46 $\alpha$
Serum Lipase	191.7 $\pm$ 4.014	566.7 $\pm$ 30.84*	346.7 $\pm$ 39.21 $\alpha$	236.7 $\pm$ 14.06 $\alpha$	260 $\pm$ 7.303 $\alpha$	233.3 $\pm$ 27.33 $\alpha$	250 $\pm$ 24.49 $\alpha$
Total Nitrate	11.87 $\pm$ 1.372	16.07 $\pm$ 1.462*	7.06 $\pm$ 0.39 $\alpha$	10.13 $\pm$ 1.31 $\alpha$	8.5 $\pm$ 0.77 $\alpha$	0.85 $\pm$ 0.796 $\alpha$	0.71 $\pm$ 0.853 $\alpha$
Total Protein	0.73 $\pm$ 0.032	0.355 $\pm$ 0.037*	0.91 $\pm$ 0.067 $\alpha$	0.73 $\pm$ 0.02 $\alpha$	0.82 $\pm$ 0.05 $\alpha$	0.73 $\pm$ 0.02 $\alpha$	0.82 $\pm$ 0.05 $\alpha$
Kidney GSH	0.47 $\pm$ 0.011	0.284 $\pm$ 0.04*	0.71 $\pm$ 0.014 $\alpha$	0.51 $\pm$ 0.017 $\beta$	0.64 $\pm$ 0.049 $\alpha$	0.10 $\pm$ 0.049	0.10 $\pm$ 0.061 $\alpha$
Liver GSH	0.48 $\pm$ 0.010	0.284 $\pm$ 0.04*	0.67 $\pm$ 0.03 $\alpha$	0.48 $\pm$ 0.03 $\beta$	0.59 $\pm$ 0.04 $\alpha$	0.10 $\pm$ 0.049	0.08 $\pm$ 0.060 $\gamma$
Lung GSH	0.50 $\pm$ 0.020	0.284 $\pm$ 0.04*	0.67 $\pm$ 0.03 $\alpha$	0.51 $\pm$ 0.02 $\gamma$	0.60 $\pm$ 0.04 $\beta$	0.08 $\pm$ 0.039 $\gamma$	0.10 $\pm$ 0.047 $\beta$
Pancreas GSH	0.48 $\pm$ 0.011	0.284 $\pm$ 0.04*	0.67 $\pm$ 0.03 $\alpha$	0.52 $\pm$ 0.02 $\beta$	0.57 $\pm$ 0.04 $\alpha$	0.10 $\pm$ 0.049	0.08 $\pm$ 0.060 $\alpha$
Kidney MDA	105.6 $\pm$ 13.67	170.9 $\pm$ 10.45*	135.2 $\pm$ 16.42 $\beta$	119.4 $\pm$ 12.77 $\alpha$	127.1 $\pm$ 12.16 $\alpha$	121.6 $\pm$ 17.5 $\alpha$	129.4 $\pm$ 13.55 $\alpha$
Liver MDA	105.6 $\pm$ 13.67	170.9 $\pm$ 10.45*	135.2 $\pm$ 16.42 $\gamma$	122.6 $\pm$ 17.42 $\alpha$	132 $\pm$ 22.09 $\alpha$	119.6 $\pm$ 17.54 $\alpha$	126.9 $\pm$ 16.51 $\beta$
Lung MDA	14.95 $\pm$ 0.66	17.09 $\pm$ 1.04	9.82 $\pm$ 1.04 $\beta$	13.18 $\pm$ 1.7 $\alpha$	11.5 $\pm$ 0.66 $\alpha$	13.23 $\pm$ 1.291 $\alpha$	14.24 $\pm$ 1.62 $\alpha$
Pancreases MDA	1.62 $\pm$ 0.6615	1.04 $\pm$ 0.4267*	2.39 $\pm$ 0.9756 $\alpha$	1.70 $\pm$ 0.69 $\beta$	1.21 $\pm$ 0.49 $\beta$	1.75 $\pm$ 0.71 $\alpha$	1.35 $\pm$ 0.55 $\beta$
Pancreas MPO	4.75 $\pm$ 2.1	32.8 $\pm$ 4.7*	6.2 $\pm$ 2.2 $\alpha$	17.2 $\pm$ 3.7 $\alpha$	11.2 $\pm$ 3.7 $\alpha$	19.2 $\pm$ 3.7 $\alpha$	14.2 $\pm$ 3.7 $\alpha$
Lung MPO	7.7 $\pm$ 0.9402	38.92 $\pm$ 2.259*	10.62 $\pm$ 1.512 $\alpha$	24.96 $\pm$ 3.03 $\alpha$	15.07 $\pm$ 1.824 $\alpha$	26.15 $\pm$ 3.48 $\alpha$	16.98 $\pm$ 2.043 $\alpha$
Liver MPO	7.883 $\pm$ 0.5419	31.73 $\pm$ 2.223*	11.6 $\pm$ 1.913 $\alpha$	26.1 $\pm$ 2.726 $\alpha$	16.98 $\pm$ 2.195 $\alpha$	28.2 $\pm$ 3.426 $\alpha$	17.38 $\pm$ 1.533 $\alpha$
Kidney MPO	7.883 $\pm$ 0.5419	32.52 $\pm$ 4.093*	6.953 $\pm$ 1.043 $\alpha$	19.3 $\pm$ 1.637 $\alpha$	11.58 $\pm$ 2.487 $\alpha$	21.65 $\pm$ 2.294 $\alpha$	15.1 $\pm$ 2.494 $\alpha$
Pancreas Catalase	0.33 $\pm$ 0.13	0.41 $\pm$ 0.1707*	0.19 $\pm$ 0.07	0.23 $\pm$ 0.0934 $\beta$	0.23 $\pm$ 0.0948 $\alpha$	0.18 $\pm$ 0.07 $\alpha$	0.12 $\pm$ 0.05 $\alpha$
kidney SOD	2.33 $\pm$ 0.95	1.50 $\pm$ 0.6146	2.61 $\pm$ 1.06 $\beta\alpha$	1.89 $\pm$ 0.774	1.891 $\pm$ 0.771 $\beta$	1.63 $\pm$ 0.66	1.171 $\pm$ 0.47
Liver SOD	0.83 $\pm$ 0.34	1.01 $\pm$ 0.4153	1.97 $\pm$ 0.80 $\alpha$	1.527 $\pm$ 0.62 $\beta$	1.78 $\pm$ 0.7266 $\beta$	1.323 $\pm$ 0.54 $\beta$	1.63 $\pm$ 0.66
Lung SOD	1.04 $\pm$ 0.42	1.03 $\pm$ 0.42*	3.44 $\pm$ 1.40 $\alpha$	2.06 $\pm$ 0.84	3.83 $\pm$ 1.563 $\alpha$	1.04 $\pm$ 0.42	2.25 $\pm$ 0.91 $\alpha$
Pancreases SOD	2.09 $\pm$ 0.85	1.51 $\pm$ 0.6191*	1.49 $\pm$ 0.61 $\alpha$	1.32 $\pm$ 0.54	1.81 $\pm$ 0.74	1.50 $\pm$ 0.614 $\gamma$	1.04 $\pm$ 0.42

\* $p < 0.0001$  when compared with normal control,  $\alpha p < 0.0001$ ,  $\beta p < 0.001$ ,  $\gamma p < 0.01$  when compared with disease control group. LW1: Lawsone 100, LW2: Lawsone 200, PI 100: pinene 100 and PI200: pinene 200 mg/kg. [Values are mean  $\pm$  SEM from 6 animals in each group]

**Table 2: Effect of alpha-pinene and Lawsone on IL-6, TNF- $\alpha$  and CRP after L-arginine induced acute pancreatitis.**


Parameter/Groups	NC	DC	LW1	LW2	PIN1	PIN2
IL-6	29.3 $\pm$ 1.569	90.48 $\pm$ 1.689*	62.42 $\pm$ 1.358 $\gamma$	33.57 $\pm$ 1.182 $\gamma$	66.9 $\pm$ 1.131	35.35 $\pm$ 2.464 $\gamma$
TNF-a	19.33 $\pm$ 1.541	26.32 $\pm$ 3.036	21.58 $\pm$ 0.9711	18.57 $\pm$ 1.316	21.9 $\pm$ 1.987	17.35 $\pm$ 1.383
CRP	415.3 $\pm$ 7.762	16403 $\pm$ 119*	8611 $\pm$ 101.7 $\gamma$	546.8 $\pm$ 9.874 $\gamma$	7436 $\pm$ 63	562.8 $\pm$ 17.44 $\gamma$

N.C- Normal Control, D.C- Disease Control, LW 100- Lawsone 100 mg/kg, LW 200- Lawsone 200 mg/kg, PI 100: pinene 100 and PI200: pinene 200 mg/kg. \* $P < 0.0001$  when compared with normal control,  $\gamma P < 0.01$  when compared with disease control group.


**Histopathology**

Fig. 1—Effect of lawsone on pancreatic histopathological changes after L-arginine induced acute pancreatitis [(a) normal control, (b)


standard control, (c) disease control 24 h hemorrhage, (d) disease control 24 h edema and necrosis, (e) lawsone 100 mg/ kg, ip, 24, (f) lawsone 200 mg/ kg, ip, 24 h, (g) pinene 100 mg/ kg, ip, 24, (h) pinene 200 mg/ kg, ip, 24 h] (H&E  $\times$ 200)




a. Normal Control




b. Standard Control




c. Disease control 24 hrshemorrhage



d. Disease control 24 hrsOedema& necrosis




e. Lawsone 100 mg/ kg, ip, 24 hrs




f. Lawsone 200 mg/ kg, ip, 24 hrs





G. Pinene 100 mg/ kg, ip, 24 hrs



H. Pinene 200 mg/ kg, ip, 24 hrs

## DISCUSSION

The present study demonstrated that treatment with lawsone (200 mg/kg) and pinene (200 mg/kg) efficiently reduced the severity of L-arginine induced acute pancreatitis in rats. In consistent with previous reports[2,19,20], in the present study administration of L-arginine significantly developed the acute pancreatitis characterized by raised levels of serum amylase, lipase and acinar cell necrosis. Serum amylase and lipase levels are the important diagnostic markers for acute pancreatitis. They usually rise within 4.8 hours of the initial attack, peaks at 24 hours[4,19,21]. Similarly, in accordance with previous reports, in the present study induction of pancreatitis significantly increased the serum amylase and lipase levels at 24 hours. Treatment with lawsone decreased the serum amylase and lipase levels, indicates protective effect of lawsone at early stage of the disease progression. In consistent with previous reports[3,4,19], in the present study induction of pancreatitis significantly increased the pancreatic MPO, MDA, nitrite, catalase and SOD and decreased the GSH levels. MPO, a marker of neutrophil infiltration is an enzyme found in neutrophils and its activity is linearly related to infiltration of neutrophils[22,23]. In agreement with previous reports[3,24], in the present study induction of pancreatitis with L-Arginine increased the pancreatic MPO levels. Inhibition of the neutrophils infiltration can attenuate the pancreatic injury[25]. Treatment with lawsone significantly decreased the pancreatic MPO levels probably due to its anti-inflammatory action. MDA, a marker of lipid peroxidation was elevated in Larginine treated rats. Lipid peroxidation is a process mediated by free radicals, which results in impairment of the membrane functional and structural integrity[4,20,26]as a consequence of oxidative deterioration of polyunsaturated fatty acids of cell membrane. It could be attributed to the accumulation of free radicals proposed to be generated by L-arginine. The change in levels of catalase and SOD remains controversial. A few studies[26] reported the fall in these enzyme levels at 24 hours and few studies[20] reported as raised levels of these enzymes. In consistent with previous reports[2], in the present study significant increase in SOD and catalase level was observed. It indicates that oxidative stress caused by L-arginine may up-regulate the activity of antioxidant enzymes to facilitate rapid removal of accumulated reactive oxygen and nitrogen species[20]. It is well known that GSH is found to be decreased in L-arginine treated rats indicating enhanced oxidative stress as the disease progresses[20]. The role of NO in the initiation and progression of acute pancreatitis remains controversial[27]. Some studies[28,29,30][31]reported that NO increase the pancreatic blood flow and/or secretion in response to endothelium derived NO and ameliorates the pancreatic dysfunction, whereas others[32,33,34]suggested that NO aggravates pancreatic oxidative stress and damage. In agreement with previous reports[34], in the present study significant increase in NO and pancreatic edema was observed in L-arginine received rats. Previous investigations[34] demonstrated that, administration of excess L-arginine could induce iNOS activity and increase the NO levels in pancreas. The raised

levels of NO can increase vascular/micro capillary permeability and may contribute to the pancreatic edema and acinar cell damage[34]. Treatment with lawsone significantly restored the pancreatic MDA, nitrite, edema, catalase, SOD and GSH in L-arginine received rats. Passaglia[35] stated that acinar cells are the protein factory of the body. In acute pancreatitis, catabolism of proteins could increase up to 80%. Consequently, a sharp decline in protein content was observed in pancreas. In consistent with previous reports[4,36,37,38]pancreatic total protein content, a marker of the tissue damage was found to decrease in L-arginine received rats, this was also corrected from previous results. Treatment with lawsone significantly increased the total protein content. It is well known that the extent of pancreatic tissue damage in acute pancreatitis correlates with the levels of inflammatory mediators and free radicals. In agreement with previous reports [4,24,21,39], in the present study, histopathological assessments revealed that, induction of pancreatitis resulted in pancreatic damage characterized by acinar cell necrosis, mono nuclear cell infiltration, edema and hemorrhage. Treatment with lawsone protected the pancreas from L-arginine induced injury. In conclusion, the present study suggests that treatment with lawsone significantly ameliorated the severity of L-arginine induced pancreatitis by reducing the neutrophil infiltration and oxidative stress markers and this effect may be due to antioxidant and anti-inflammatory properties of the lawsone.

## Disclaimer

The views and opinions expressed in this article are those of the authors, and they do not reflect in any way those of the institutions to which they are affiliated with.

## REFERENCE

1. Robbins K, Basic Pathology, Elsevier Publications, Chicago, 1997; 902-928.
2. Robbins K, Basic Pathology, Elsevier Publications, Chicago, 2002; 675-676.
3. Miguel MG, Antioxidant and Anti-Inflammatory Activities of Essential Oils: A Short Review. 2010; 15:9252-9287.
4. Zhou JY, Tang FD, Mao GG, Bian RL. Effect of alpha-Pinene on nuclear translocation of NF-kappa B in THP-1 cells, 2004; 25 (4): 480-4.
5. Sauriasari R, Wang DH, Takemura Y, Tsutsui K, Masuoka N, Sano K, Horita M, Wang BL & Ogino K, Cytotoxicity of lawsone and cytoprotective activity of antioxidants in catalase mutant *Escherichia coli*, Toxicology. 2007; 235(1-2):103-11.
6. Mikhaeil BR, Badria FA, Maatooq GT & Amer MMA, Antioxidant and Immunomodulatory Constituents of Henna Leaves, Z. Naturforsch. 2004; 59 c: 468-476.
7. Melo CM, Carvalho KMMB, Neves JCDS, Morais TC, Rao VS, Santos FA, Brito GADC & Chaves MH,  $\alpha$ , $\beta$ -amyrin, a natural triterpenoid ameliorates L-arginine induced acute pancreatitis in rats, World J Gastroenterol, 2010; 16(34):4272.

8. Sidhu S, Pandhi P, Malhotra S, Vaiphei K & Khanduja KL, Melatonin treatment is beneficial in pancreatic repair process after experimental acute pancreatitis, *Eur J Pharmacol.* 2010; 628:282.
9. Bradley PP, Priebat DA, Christensen RD & Rothstein G, Measurement of cutaneous inflammation: estimation of neutrophil content with an enzyme marker, *J Invest Dermatol.* 1982; 78:206
10. Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS & Tannenbaum SR, Analysis of nitrate and nitrite in biological fluids, *Anal Biochem,* 126 (1982) 131.
11. Gamal el-din AM, Mostafa AM, Al-Shabanah OA, Al-Bekairi AM & Nagi MN, Protective effect of arabic gum against acetaminophen-induced hepatotoxicity in mice, *Pharmacol Res.* 2003; 48:631.
12. Ohkawa H, Ohishi N & Yagi K, Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction, *Anal Biochem.* 1979; 95: 351.
13. Aebi H, Catalase, in *Methods in enzymology*, edited by L Packer, Academic Press, Orlando. 1984; 105:121.
14. Kakkar R, Mantha SV, Radhi J, Prasad K & Kalra J, Increased oxidative stress in rat liver and pancreas during progression of streptozotocin-induced diabetes, *Clin Sci.* 1998; 94: 623.
15. Misra HP & Fridovich I, The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase, *J Biol Chem,* 1972; 247:3170.
16. Sedlak J & Lindsay RH, Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent, *Anal Biochem,* 1968; 25:192.
17. Schmidt J, Rattner DW, Lewandrowski K, Compton CC, Mandavilli U, Knoefel WT & Warshaw AL, A better model of acute pancreatitis for evaluating therapy, *Ann Surg.* 1992; 215:44.
18. Eşrefoglu M, Gül M, Ates B, Batçioğlu K & Selimoğlu MA, Antioxidative effect of melatonin, ascorbic acid and N-acetylcysteine on caerulein-induced pancreatitis and associated liver injury in rats, *World J. Gastroenterol,* 2006; 12:259.
19. Szabolcs A, Reiter JR, Letoha T, et al. Effect of melatonin on the severity of L-arginine-induced experimental acute pancreatitis in rats. *World J. Gastroenterol.* 2006; 12: 251-258.
20. Hegyi P, Rankoncay Z, Sari R, et al. L-arginine induced experimental pancreatitis. *World J. Gastroenterol.* 2004; 10: 2003-2009.
21. Caroline MM, Karine MM, Julliana C, et al. a-amyrin, a natural triterpenoid ameliorates L-arginine induced acute pancreatitis in rats. *World J Gastroenterol.* 2010; 16(34): 4272-4280.
22. Poch B, Gansauge F, Rau B, et al. The role of polymorphonuclear leukocytes and oxygen-derived free radicals in experimental acute pancreatitis: mediators of local destruction and activators of inflammation. *FEBS Lett.* 1999; 461: 268-272.
23. Song AM, Bhagat L, Singh VP, et al. Inhibition of cyclooxygenase-2 ameliorates the severity of pancreatitis and associated lung injury. *Am. J. Physiol. Gastrointest. Liver Physiol.* 2002; 283: 1166-1174.
24. Abdin A, Abd El-Hamid A, Abou El-Seoud H, et al. Effect of pentoxifylline and/or alpha lipoic acid on experimentally induced acute pancreatitis. *Eur J Pharmacol.* 2010; 643: 289-296.
25. Shi C, Zhao X, Wang X, et al. Potential effects of PKC or protease inhibitors on acute pancreatitis induced tissue injury in rats. *Vascul. Pharmacol.* 2007; 46:406-411.
26. Czako L, Takacs T. Involvement of oxygen derived free radicals in L-arginine induced acute pancreatitis. *Dig. Dis. Sci.* 1998; 43(8): 1770-77.
27. Sweiry JH, Mann GE. Role of oxidative stress in the pathogenesis of acute pancreatitis. *Scand. J. Gastroenterol.* 1996; 31: 10-15.
28. Gukovskaya A, Pandol S. Nitric oxide production regulates cGMP formation and calcium influx in pancreatic acinar cells. *Am. J. Physiol.* 1994; 266:350-356.
29. Holst JL, Rasmussen TN, Schmidt P. Role of nitric oxide in neurally induced pancreatic exocrine secretion in pigs. *Am. J. Physiol.* 1994; 266: 206-213.
30. Satoh A, Shimosegawa T, Abe T, et al. Role of nitric oxide in the pancreatic blood flow response to caerulein. *Pancreas* 1994; 9: 574-579.
31. Patel AG, Toyama MT, Nguyen TN, et al. Role of nitric oxide in the relationship of pancreatic blood flow and exocrine secretion in cats. *Gastroenterology* 1995; 108: 1215-1220.
32. Tani S, Itoh H, Okabayashi Y. New model of acute necrotizing pancreatitis induced by excessive doses of arginine in rats. *Dig. Dis. Sci.* 1990; 35(3): 367-374.
33. Dabrowski A, Gabryelewicz A. Nitric oxide contributes to multi-organ oxidative stress in acute pancreatitis. *Scand. J. Gastroenterol.* 1994; 29: 943-948.
34. Takacs T, Czako L, Morschl E, et al. The role of nitric oxide in edema formation in L-arginine-induced acute pancreatitis. *Pancreas* 2002; 25: 277-282.
35. Passaglia C. Nutritional problems in acute pancreatitis. *Recent Prog. Med.* 2007; 98: 335-338.
36. Biradar S and Veeresh B, Screening of Natural Antioxidants by using L-Arginine induced acute Pancreatitis Model, *International Journal of Drug Development & Research,* 2012: 364 (4): 284-297.
37. Biradar S and Veeresh B, Protective effect of lawsone on L-arginine induced acute pancreatitis in rats, *Indian journal of experimental biology.* 2013; 51, 256-61.
38. Aruna R, Geetha A and Suguna P, Rutin modulates ethanol and high fat diet induced inflammatory changes in pancreas - a dose response study in rats. *Int J Pharm Pharm Sci,* 2012; 4 (4): 409-14
39. Priyadarsini RI, Namratha JR and Ravi sankarreddy D (2012), Glucokinase activators: a glucose sensor role in pancreatic islets and hepatocyte. *Int J Pharm Pharm Sci,* 4(2): 81-87.