

ANTIBACTERIAL ACTIVITY OF LEAVES AND ROOTS OF *Eclipta alba*

KARUNAMBIGAI, A. AND GAYATHRI DEVI, S.*

*Department of Biochemistry, Biotechnology and Bioinformatics, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore 641043, Tamil Nadu, India. Email: gayathridevi.adu@gmail.com.

Received: 18 Oct 2013, Revised and Accepted: 09 Nov 2013

ABSTRACT

Objective: The present study was aimed to investigate the antibacterial activity of leaves and roots of *Eclipta alba*.

Methods: The antibacterial activity of leaves and roots of *Eclipta alba* was evaluated by agar well diffusion method and minimum inhibition concentration against different bacterial species such as *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Staphylococcus aureus* and *Shigella flexneri*.

Results: 20 µl of chloroform extract of leaves and root powder of *Eclipta alba* was found to exhibit best inhibitory activity against *Klebsiella pneumoniae* such as 32 mm and 35 mm respectively when compared to other solvents and hot aqueous extract. Leaves of benzene extract exhibited good inhibitory zone of 26mm against *Shigella flexneri* and growth of *Bacillus subtilis* was inhibited up to 28mm by chloroform extract of root. All tested organisms exhibited lowest zone of inhibition between 8-10mm with acetone, methanol and aqueous extract of both leaves and roots of *Eclipta alba*. MIC was between concentration of 8-0.25mg/ 100 µl with all tested bacteria. From the results, it is clear that the extracts of leaves and root of *Eclipta alba* possess potential broad spectrum antibacterial activity.

Keywords: *Eclipta alba*, Micro organisms, Antibacterial activity, Leaves and Root extraction.

INTRODUCTION

The frequency of life threatening infections caused by pathogenic microorganisms has increased worldwide and is becoming an important cause of morbidity and mortality in immun compromised patients in developed countries [1]. The ready availability and economy of plants as direct therapeutic agents make plants more attractive when compared to modern medicine [2]. Medicinal plants have been used for centuries before the advent of orthodox medicine. Leaves, flowers, stems, roots, seeds, fruit and bark can also be constituents of herbal medicines [3]. More than 80 per cent of the world's population in the world still depends on traditional medicine for their primary health care. They continue to provide scientists with invaluable compounds of starting points for the development of new drugs [4]. Conversely, because information on the use of plant species for therapeutic purpose has been passed from one generation to the next through oral tradition, this knowledge of therapeutic plants has started to decline and become obsolete through the lack of recognition by younger generations as a result of a shift in attitude and ongoing socio-economic changes [5].

Historically medicinal plants have provided a good source of anti-infective agents; emetine, quinine and berberine remain highly effective instruments in the fight against microbial infections [6]. The author stated that plant derived medicines have been a part of traditional health care in most parts of the world of thousands of years and there is increasing interest in them as source of agents to fight microbial diseases [7]. *Eclipta alba* is commonly known as Bhringaraja or Maka belonging to the family Asteraceae/Compositae. The herb contain wedelolactone and demethylwedelolactone which possessing potent Antihepatotoxic property. Other prominent chemical constituents present are ecliptal, ecliptine, ecliptalbine, α-terthienylmethanol, β-amyrin and sigmasterol [8]. The main objective of this study is assessment of antibacterial activity *Eclipta alba*.

MATERIALS AND METHODS

Collection of plant samples

The plant *Eclipta alba* was freshly collected from local market at Coimbatore. The plant was duly authenticated by Botanical Survey of India, Tamil Nadu Agricultural University Coimbatore. A voucher specimen was deposited in the Department of Biochemistry, Avinashilingam University for Women, Coimbatore, Tamil Nadu, India (Plate I).

Plate I

*Eclipta alba*

Collection of microorganisms

The bacterial strains used for the experiment such as *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Staphylococcus aureus* and *Shigella flexneri* were collected from the Department of Biotechnology, PSG Institute of Medical Science, Coimbatore and stored at -70°C.

Preparation of extracts

The collected leaves and roots of the plant were washed and air dried in the shade at room temperature for complete drying. The dried sample was crushed with the aid of mechanical grinder to make powder form. The powder obtained was extracted with selected solvents based on polarity using soxhlet apparatus in the 6 hours intervals. The solvents used in sequential order were respectively petroleum ether, benzene, chloroform, acetone and methanol.

Preparation of hot aqueous extract

5 grams of dried leaves and root powder were boiled with 100ml of water until it becomes half the volumes. The extracts were kept as such in the room temperature for evaporating until it become semi solid and all of the dried extracts were dissolved in dimethylsulfoxide (DMSO), stored in the refrigeration until required for use.

Agar well disc diffusion method for antibacterial activity

Organic extract were screened for antibacterial activity using the agar disc diffusion method [9]. The test samples dissolved in DMSO were used. Muller - Hinton agar was used as a culture media for bacterial growth. Loaded 20 micro liter of the each extract in the corresponding 6mm diameter in Whatmann filter paper No1. Plates were incubated for 24 h at 37 ° C for bacterial growth. The diameter of inhibition zone around each well was measured and recorded. The antibiotic disc such as penicillin, ampicillin, gentamycin and streptomycin were used as control [10].

Determination of Minimum Inhibitory Concentration (MIC)

The MIC was defined as the lowest concentration of the sample that did not show any growth of the tested microorganism. In the present study 100µl of nutrient broth was taken in the wells of 96 well plate. To this 100µl of these diluents was added in the well making a twofold dilution. 100µl of this dilution was transferred and mixed with 100µl of the diluents in second row making a 4:1 dilution. This proceeds consecutively down the plate making two fold dilutions in each well. After the dilution of the sample is completed and aliquot of test organisms was added to all wells and the microtitre plates were incubated overnight. If the sample concentration was sufficient to kill the organism, no growth appeared and the wells will be clear. At this point there was an insufficient sample to kill the organism and well will be cloudy, indicating growth that is the minimum inhibitory concentration of the sample against that specific organism [11].

Statistical analysis

Values are given in the mean ± SD, and the differences between values were determined by the student's t-test. Values of P < 0.01 were considered significant.

RESULTS

The results of antibacterial activity of *Eclipta alba* were given in Table 1 and Table 2. Among all the tested extracts, the chloroform extracts of leaves and root were to be more effective than remaining other solvents and hot aqueous extract against all the tested organisms. In chloroform extract, all test organisms gave good susceptibility with zone of inhibition ranges from 08-35 mm than antibiotic disc. For benzene extract of leaves and root *Shigella flexneri* was most susceptible to the extract and shows 26mm and 12 mm of zone of inhibition respectively. *Bacillus subtilis*, *Pseudomonas fluorescens* and *Staphylococcus aureus* were exhibited similar activity for both leaves and root extract between 09 -11 mm zone of inhibition. No zone of inhibition was observed in the petroleum ether, acetone and aqueous extracts of leaves and root of *Eclipta alba* against the *Klebsiella pneumoniae*. Methanol and acetone extract did not show any inhibition against the microorganism *Escherichia coli* and *Bacillus subtilis* respectively. The hot aqueous extract gave better results against inhibition of *Shigella flexneri* than acetone and methanol. Results of different extracts of the leaves and root of *Eclipta alba* were compared with the control plates. Hence, the leaves and root extracts of the *Eclipta alba* was found to show a good inhibition against the selected bacterial strains.

Table 1: Antibacterial activity of *Eclipta alba*

Extracts		Zone of Inhibition (mm)					
		<i>E.coli</i>	<i>B.subtilis</i>	<i>P. fluorescens</i>	<i>S. flexneri</i>	<i>K. pneumoniae</i>	<i>S. aureus</i>
PE	Leaf	15	13	10	-	-	-
	Root	08	10	10	09	-	08
BEN	Leaf	13	09	09	26	08	09
	Root	08	10	11	12	12	11
CHL	Leaf	13	28	22	23	32	10
	Root	08	09	12	12	35	09
ACE	Leaf	09	-	-	09	-	08
	Root	08	-	09	09	-	-
MeOH	Leaf	-	08	09	10	-	08
	Root	-	09	09	-	12	-
Aqu	Leaf	08	08	-	15	-	-
	Root	08	-	08	11	-	09

PE = Petroleum ether, BEN = Benzene, CHL = Chloroform, ACE = Acetone, MeOH = Methanol, Aqu = Aqueous, - = No activity.

#Values are the average of triplicate; includes Whatmann filter paper No1 diameter (6mm)

Antibiotics discs such as penicillin, ampicillin, gentamycin and streptomycin were used as a control for all bacterial strains. The results were shown in the Table 2.

Table 2: Antibiotic sensitivity against control

Control	<i>E.coli</i>	<i>B.subtilis</i>	<i>P.fluorescens</i>	<i>S.flexneri</i>	<i>K.pneumoniae</i>	<i>S.aureus</i>
PCN	07	10	-	16	-	22
AMP	13	-	11	26	-	22
GEN	17	24	20	18	-	21
STREP	10	23	18	15	11	20

PCN - Pencilin, AMP - Ampicillin, GEN - Gentamycin , STREP - Streptomycin

Table 3: Antibacterial activity of *Eclipta alba* against *Escherichia coli* and *Bacillus subtilis* by minimum inhibitory concentration (MIC) method

Extracts		<i>Escherichia coli</i> (mg/100µl)						<i>Bacillus subtilis</i> (mg/100µl)						
		0	8	4	2	1	0.5	0	8	4	2	1	0.5	0.25
Benzene	Leaves	-	-	+	-	+	-	-	+	-	-	-	-	-
	Root	-	+	+	-	-	-	-	+	+	+	+	-	-
Chloroform	Leaves	-	+	+	+	-	-	-	+	+	+	+	+	+
	Root	-	-	+	+	-	-	-	+	+	+	+	-	-

+ indicates positive -indicates negative

Table 4: Antibacterial activity of *Eclipta alba* against *Pseudomonas fluorescens* and *Shigella flexneria* by minimum inhibitory concentration (MIC) method

Extracts		<i>Pseudomonas fluorescens</i> (mg/100µl)						<i>Shigella flexneria</i> (mg/100µl)							
		0	8	4	2	1	0.5	0.25	0	8	4	2	1	0.5	0.25
Benzene	Leaves	-	+	+	+	-	-	-	-	+	-	+	+	+	-
	Root	-	+	+	+	+	-	-	-	-	+	+	-	-	-
Chloroform	Leaves	-	+	+	+	+	-	-	-	+	+	+	-	-	-
	Root	-	+	+	-	-	-	-	-	+	-	-	+	-	-

+ indicate positive - indicate negative

Table 5: Antibacterial activity of *Eclipta alba* against *Klebsiella pneumoniae* and *Staphylococcus aureus* by minimum inhibitory concentration (MIC) method

Extracts		<i>Klebsiella pneumoniae</i> (mg/100µl)						<i>Staphylococcus aureus</i> (mg/100µl)							
		0	8	4	2	1	0.5	0.25	0	8	4	2	1	0.5	0.25
Benzene	Leaves	-	+	+	+	-	+	-	-	+	-	+	+	-	-
	Root	-	+	+	-	+	+	-	-	-	+	+	-	-	-
Chloroform	Leaves	-	-	+	+	-	-	-	-	+	+	-	-	-	-
	Root	-	+	+	-	+	+	-	-	-	+	+	+	-	-

+ indicate positive - indicate negative

Minimum inhibitory concentration (MIC)

Those extracts giving zone of inhibition for all the selected bacterial strains were chosen to assay the minimum inhibitory concentration with agar dilution method and it is given in Table 3-5.

Eclipta alba extracts were inoculated against *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Staphylococcus aureus* and *Shigella flexneri* in the concentration of 8-0.25 mg / 100µl. The growth of *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Staphylococcus aureus*, *Shigella flexneri* were inhibited by benzene, chloroform extracts of leaves and root of *Eclipta alba* of the concentration of 8-1.0 mg / 100µl. 8 -2 mg /100µl of benzene and chloroform extract of leaves and root were found to have significant killing effects.

DISCUSSION

Among the tested organisms *Klebsiella pneumoniae* showed higher percent of resistance against the different extracts of leaf and root. These observations have helped in identifying the active principle responsible for such activities and in the developing drugs for the therapeutic use in human beings[12]. With no antibacterial activity, extracts may be active against other bacterial species which were not tested [13]

CONCLUSION

In the present study *Eclipta alba* has shown significant antibacterial activity against *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescens*, *Staphylococcus aureus*, *Shigella flexneri*. Hence the antibacterial activity might be due to the presence of active compounds in *Eclipta alba* against the growth of bacteria.

REFERENCE

1. Tiwari P, Kumar K, Panik R, Pandey A, Pandey A, Sahu PK. Antimicrobial activity evaluation of the root of *Carica papaya* Linn. Int.J. Pharm Tech Res. 2011; 3 (3):1641- 1648.
2. Odey MO, Iwara IA, Udiba UU, Johnson JT, Inekwe UV, Asenye ME, Victor O. Preparation of Plant Extracts from Indigenous Medicinal Plants. IJST. 2012; 1 (12): 688-692.
3. Ujowundu CO, Kalu FN, Emejulu AA, Okafor, OE, Nkwonta CG, Nwosunjoku EC. Evaluation of the chemical composition of *Mucuna utilis* leaves used in herbal medicine in Southeastern Nigeria. Afr. J. Pharm. Pharmacol. 2010; 4 (11): 811-816.
4. Fakim AG. Traditional Roles and Future Prospects for Medicinal Plants in Health Care. Asian biotechnology and development review. 2011; 13 (3):77-83.
5. Tantiado RG. Survey on Ethnopharmacology of medicinal plants in Iloilo, Philippines. International Journal of Bio-Science and Bio-Technology. 2012; 4 (4): 11-26.
6. Zia-Ul- Haq M, Ahmad M, Mehjabeen, Jehan N, Ahmad S, Qayum M, Marwat IK. Antimicrobial screening of selected flora of Pakistan. Arch. Biol. Sci., Belgrade. 2011; 63 (3): 691-695.
7. Ajay AO, Alinfolo TA. Evaluation of antibacterial activity of some medicinal plants on common enteric food-borne pathogens. Afr. J. Microbiol. Res. 2010; 4 (4): 313- 316.
8. Prasad KV, Kavita YN, Vidya NS, Sumeet BK, Manohar PJ. *Eclipta alba*: A Phytopharmacognostic Study. J.Pharm. Phytopharmacol. Res. 2012; 1 (6): 350-353.
9. Rios JL, Recio MC, Villar A. Screening methods for natural products with antibacterial activity, a review of the literature. Journal of Enthopharmacol. 1988; 23:127-149.
10. NCCLS. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: Approved standard fifth edition, NCCLS dominant M17A5; NCCLS, Wayne 1997; PA.USA.
11. Eloff JN. A sensitive and quick microplate method to determine the minimal inhibitory concentration of plants extracts for bacteria. Planta Med. 1998; 64:711-713.
12. Shrikant, Sethi S, Lal GB. Antimicrobial activity of medicinal plants on urinary tract pathogens. Int J Pharm Pharm Sci. 2012; 4(2): 626-628.
13. Majji LR, Battu GR, Jangiti R, Talluri MR. Evaluation of *in vitro* antibacterial activity of *Cassia siamea* leaves. Int J Pharm Pharm Sci. 2013; 5(3): 263-265.