

Original Article

## SIMULTANEOUS DETERMINATION OF QUERCITIN AND AMENTOFLAVONE IN METHANOLIC LEAF EXTRACT OF SEMECARPUS ANACARDIUM (LINN.F.) BY REVERSE PHASE LIQUID CHROMATOGRAPHY

PARAG A. PEDNEKAR<sup>1</sup>, VANITA KULKARNI<sup>1</sup>, BHANU RAMAN<sup>1</sup>

<sup>1</sup>Department of Chemistry, K.J. Somaiya College of Science and Commerce, Vidyavihar, Mumbai 400077.  
Email: parapednekar81@gmail.com, vanitakulkarni2@gmail.com

Received: 18 Mar 2014 Revised and Accepted: 27 Apr 2014

### ABSTRACT

**Objective:** To develop a sensitive, simple, rapid and reliable RP-HPLC method that can simultaneously determine Quercetin and Amentoflavone markers in methanolic leaf extract of *Semecarpus anacardium* (Linn.f.). In this regard HPLC method has been developed and validated as per ICH guidelines.

**Methods:** The reverse phase chromatographic separation was achieved using a Surveyor Agilent 1100 series HPLC system equipped with U.V.detector. The column used for method development and validation was Kromasil 100-5C18 (250x 4.6, 5  $\mu$ m) as stationary phase with a mobile phase comprising of Water of pH 3 (adjusted with O-phosphoric acid): Acetonitrile (55:45 v/v) at a flow rate of 1.0 ml min<sup>-1</sup>. The injection volume of the sample was 20  $\mu$ l with U.V. detection at 280 nm. The run time of the sample was 10.0 min & the column oven temperature was 25°C.

**Results:** The proposed RP-HPLC method has been validated as per ICH guidelines for its system suitability, linearity, LOD & LOQ, precision (Intra-day & Inter-day), accuracy & robustness.

**Conclusion:** The developed RP-HPLC method holds potential for detection and quantification of Quercetin and Amentoflavone markers in methanolic leaf extract of *Semecarpus anacardium* (Linn.f.).

**Keywords:** RP-HPLC, Quercetin, Amentoflavone, *Semecarpus anacardium* (Linn.f.).

### INTRODUCTION

*Semecarpus anacardium* is a deciduous tree belonging to Anacardiaceae family, growing in tropical and temperate regions of south east asian countries. Anacardiaceae family contains 700 species distributed among 60 genera. The nuts are used for treatment of variety of disorders in Ayurveda. It has been used therapeutically in neurological disorders, ulcers [1], corns [2] leprosy, leucoderma [3]. The plant is rich in secondary metabolites and mineral compositions which can be a potential source of useful therapeutic drugs [4-7].

Amentoflavone is naturally occurring biflavonoid that is found in a number of plants. It has many pharmacological activities. Investigation studies on the anti-inflammatory activities of amentoflavone and its mechanism of action has been done recently [8]. It exerts antibacterial, anti-inflammatory and antioxidative activities [9-13]. It has exhibited strong anticancer activities against MCF-7 and HeLa cancer cells through activation of hPPAR $\gamma$ , which is regulated by PTEN activation [14]. Amentoflavone, the main biflavonoid obtained from the genus *Calophyllum*, has been shown to inhibit HIV-reverse transcriptase, prevent histamine release from mast cells and cause mice to enhance their production of interferon [15]. Amentoflavone from *Selaginella*, has various biological and pharmacological effects, including antioxidant [16-19], anticancer [20-22], anti-inflammatory [23-25], antimicrobial [26], antiviral such as influenza (A, B), hepatitis (B), human immunodeficiency virus (HIV-1), herpes (HSV-1, HSV-2), herpes zoster (VZV), measles [27-32], respiratory syncytial virus (RSV) [33], vasorelaxation [34], antiurcerogenic [10], antistomachic-ache [11], antidepressant [23], anxiolytic [35,36], analgesic [37] and antiangiogenesis agent [38].

Quercetin has an antioxidative capacity [39]. Hydroxyl radical scavenging activity was detected in quercetin [40]. A human study showed topical and oral administration of quercetin to reduce skin damage during radiotherapy in patients with head and neck cancers [41]. It has a history of use by nutritional physicians as an anti-inflammatory and antiallergy agent [42]. It is found in a broad range

of foods like grape skins, red onions, green tea and tomatoes. Quercetin is attracting intense scientific interest for its unique antiaging and immune boosting activities [43]. Laboratory studies demonstrate that quercetin traps developing cancer cells in the early phases of their replication cycle, effectively preventing further malignant development and promoting cancer cell death [44].

### MATERIALS AND METHODS

#### Collection and Authentication of the plant

The leaves of *Semecarpus anacardium* were collected from Mumbai, Maharashtra. The identification of the plant was done at the Blatter Herbarium, St. Xavier's College, Mumbai. The plant specimen matches with the Blatter Herbarium specimen no.T- 472 of S. C. Tavakari.

#### Chemicals and Standards

HPLC grade Acetonitrile from Qualigens, Mumbai were used. Ultra pure water, generated by use of a Milli-Q System (Millipore), was used for preparation of mobile phase. Standard compounds Quercetin and Amentoflavone were from Sigma Aldrich. O-phosphoric acid of Merck specialities Pvt. Ltd., Mumbai were used in study.

#### Sample preparation

The leaf powder of *Semecarpus anacardium* (20 gm) was extracted with 250 ml methanol by soxhlet extraction for 8 hrs. The extract was concentrated on water bath at 60°C. The obtained dark brown thick liquid was stored in a glass vial in refrigerator.

#### Selection of Mobile Phase

Sample and standard solubility, stability and suitability were considered while choosing from different mobile phase compositions and proportions to get good chromatographic separations. From the various mobile phases, Water of pH3: Acetonitrile (55:45 %v/v) was selected. Water of pH3 was adjusted with O-phosphoric acid. The detection wavelength of 280nm was used throughout.

### Selection of Diluent

The Distilled water was used as diluent.

### Preparation of Standard Stock Solution

An accurately weighed quantity of Quercitin and Amentoflavone was transferred to a volumetric flask, dissolved and diluted to the mark with distilled water to obtain standard stock solution of 1000 µg ml<sup>-1</sup>.

### Apparatus and Chromatographic Parameters

The optimized parameters which were used as a final method for the simultaneous estimation of Quercitin and Amentoflavone is represented in the Table 1.

### Method validation

The optimized chromatographic method was completely validated according to the procedures described in ICH guidelines Q2 (R1) for the validation of analytical methods [45].

### System Suitability

20 µl of mixed standard solution of Quercitin and Amentoflavone (5 µg/ml) was injected (n=4) under optimized chromatographic conditions to evaluate the suitability of system. Values with % RSD < 2 were accepted (Table 2).

### Specificity

Specificity of the HPLC method was demonstrated by the separation of the analytes from other potential components such as impurities, degradants or excipients. A volume of 20 µl of individual standards, mixed stock of standards and extract solution were injected and the chromatogram was recorded (n=3). Hence, the proposed method was specific for Quercitin and Amentoflavone (Figure 1).

### Linearity

The linearity of calibration curve of Quercitin and Amentoflavone over the concentration range of 0.25-20 µg/ml (n=3) through

proposed HPLC method was carried out. Correlation coefficient (r) was found to be 0.9949 for Quercitin & 0.9964 for Amentoflavone respectively (Table 3 & Figure 2 & 3).

### Limit of Detection and Limits of Quantitation

Limit of Detection (LOD) and Limit of Quantification (LOQ) were determined by using the formula based on the standard deviation of the response area of blank and the slope of calibration curve. LOD and LOQ were calculated by using equations,

$$\text{LOD} = 3.3 \times \sigma / \text{slope} \quad \& \quad \text{LOQ} = 10 \times \sigma / \text{slope}$$

Where,  $\sigma$  = Standard deviation of Blank area (n=3), Slope = Slope of the calibration curve.

By these equations LOD was found to be 5.690 x 10<sup>-3</sup> µg/ml for Quercitin & 6.729 x 10<sup>-5</sup> µg/ml for Amentoflavone and LOQ for Quercitin 0.017 µg/ml & 2.039 x 10<sup>-4</sup> µg/ml for Amentoflavone.

### Precision

#### Intra-day precision

Intra-day precision was determined by injecting 3 different concentrations (80%, 100%, 120%) of the target concentration (n=4) times in the same day. Peak area was measured and %RSD was calculated (Table 4).

#### Inter-day precision

Inter-day precision was determined by injecting 3 different concentrations (80%, 100%, 120%) of the target concentration (n=4) times for 3 consecutive days. Peak area was measured and %RSD was calculated (Table 5-7).

### Accuracy

For the accuracy of proposed method (n=3), recovery studies were performed by Standard addition method at 3 different levels 50%, 80% and 100% of the target concentration i.e. Quercitin (15 µg/ml) & Amentoflavone (9.49 µg/ml). Results of Recovery studies were found to be satisfactory (Table 8).

**Table 1: Optimized chromatographic conditions for Quercitin and Amentoflavone.**

<b>Equipment</b>	Surveyor Agilent 1100 series HPLC system with G1379A Degasser, G1311A Quaternary pump, G1329A Autosampler, G1330B Thermostat ALS Therm, G1316A Column oven (COLCOM), G1314A U.V. visible detector
<b>Column</b>	Kromasil 100-5C18 (250x 4.6, 5 µm)
<b>Mobile Phase</b>	Water of pH3: Acetonitrile (55:45 %v/v)
<b>pH of Mobile Phase</b>	pH 3.00 adjusted with OPA
<b>Flow rate</b>	1.0 ml min <sup>-1</sup>
<b>Injection volume</b>	20 µl
<b>Column oven</b>	25°C
<b>Wavelength</b>	280 nm
<b>Run time</b>	10.0 min
<b>Mode of Operation</b>	Isocratic elution

**Table 2: System Suitability of Quercitin and Amentoflavone (n=4).**

Compound	Parameter	Acceptance	Average	%RSD
<b>Quercitin 5 µg/ml</b>	Retention Time (min)	% RSD < 2	4.55	0.03
	Peak area	% RSD < 2	112.8	1.45
	Theoretical plates per meter	> 3500	11,252.64	1.69
	Symmetry factor / Tailing factor	< 2	0.63	1.62
	Resolution	>2	9.59	0.54
<b>Amentoflavone 5 µg/ml</b>	Retention Time (min)	% RSD < 2	6.29	0.04
	Peak area	% RSD < 2	175.05	0.48
	Theoretical plates per meter	> 3500	17,437.52	0.45
	Symmetry factor / Tailing factor	< 2	0.74	0.68
	Resolution	>2	9.59	0.54




Fig. 1: Chromatograms of A: Methanolic soxhlet leaf extract of *Semecarpus anacardium*, B: Quercetin 10 µg/ml, C: Amentoflavone 10µg/ml D: Quercetin & Amentoflavone 10 µg/ml.




Fig. 2: Linearity studies of Quercetin




Fig. 3: Linearity studies of Amentoflavone

Table 3: Linear regression data for Calibration curves (n= 3).

Parameters	Quercetin	Amentoflavone
Detection wavelength	280 nm	280 nm
Linearity range	0.25 µg/ml - 20 µg/ml	0.25 µg/ml - 20 µg/ml
Correlation coefficient (r)	0.9949	0.9964
Regression equation	$y = 17.05x + 0.5$	$y = 56.88x + 1.5$
LOD	$5.690 \times 10^{-3}$ µg/ml	$6.729 \times 10^{-5}$ µg/ml
LOQ	0.017 µg/ml	$2.039 \times 10^{-4}$ µg/ml

Table 4: Intra-day precision of Quercitin and Amentoflavone (n=4).

Quercitin	Intra-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
12 µg/ml	4.53	0.08	254.68	0.23
15 µg/ml	4.54	0.05	329.08	0.07
18 µg/ml	4.54	0.04	389.6	0.10
Amentoflavone	Intra-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
7.59 µg/ml	6.25	0.07	466.75	0.12
9.49 µg/ml	6.26	0.08	593.5	0.03
11.39 µg/ml	6.25	0.10	721.8	0.09

Table 5: 1<sup>st</sup> Inter-day precision of Quercitin and Amentoflavone (n=4).

Quercitin	1 <sup>st</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
12 µg/ml	4.55	0.18	263.38	0.13
15 µg/ml	4.54	0.12	331.58	0.08
18 µg/ml	4.56	0.07	373.63	0.06
Amentoflavone	1 <sup>st</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
7.59 µg/ml	6.26	0.17	412.25	0.03
9.49 µg/ml	6.25	0.15	547.40	0.13
11.39 µg/ml	6.29	0.14	590.53	0.05

Table 6: 2<sup>nd</sup> Inter-day precision of Quercitin and Amentoflavone (n=4).

Quercitin	2 <sup>nd</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
12 µg/ml	4.54	0.08	254.78	0.07
15 µg/ml	4.55	0.10	318.58	0.03
18 µg/ml	4.55	0.04	372.83	0.06
Amentoflavone	2 <sup>nd</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
7.59 µg/ml	6.23	0.18	384.45	0.02
9.49 µg/ml	6.26	0.16	504.40	0.07
11.39 µg/ml	6.27	0.06	506.43	0.04

Table 7: 3<sup>rd</sup> Inter-day precision of Quercitin and Amentoflavone (n=4).

Quercitin	3 <sup>rd</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
12 µg/ml	4.55	0.07	247.45	0.19
15 µg/ml	4.55	0.04	322.65	0.04
18 µg/ml	4.55	0.07	374.63	0.07
Amentoflavone	3 <sup>rd</sup> Inter-day			
	Retention time		Peak Area	
	Mean	%RSD	Mean	%RSD
7.59 µg/ml	6.26	0.19	363.45	0.07
9.49 µg/ml	6.26	0.04	489.10	0.02
11.39 µg/ml	6.27	0.12	471.75	0.03

### Robustness

The Robustness of the HPLC method (n=3) was evaluated by analysing the parameters after varying the flow rate ( $\pm 20\%$ ),

temperature ( $\pm 5$  °C) and wavelength ( $\pm 5$  nm). Changes in % RSD of peak area w.r.t Quercitin and Amentoflavone standards and *Semecarpus anacardium* methanolic leaf extract are given below in Table 9.

Table 8: Accuracy study (n=3).

Standard	Level	Prealysed sample ( $\mu\text{g ml}^{-1}$ )	Amount of standard added to preanalysed sample ( $\mu\text{g ml}^{-1}$ )	Total amount Recovered/ Found in sample ( $\mu\text{g ml}^{-1}$ )	SD	% RSD	% Recovery
Quercitin	0%	15.00	0	0	0	0	0
	50%	15.00	7.50	21.88	0.01	0.06	91.69
	80%	15.00	12.00	27.89	0.14	0.49	107.43
	100%	15.00	15.00	29.51	0.05	0.16	96.72
Amentoflavone	0%	9.49	0	0	0	0	0
	50%	9.49	4.75	13.56	0.01	0.08	85.79
	80%	9.49	7.59	16.96	0.01	0.06	98.35
	100%	9.49	9.49	18.07	0.01	0.07	90.43

Table 9: Robustness study (n=3).

Changing Parameter	Quercitin and Amentoflavone Standards				
	Actual Parameter	Modification	% RSD of Peak areas		
			Quercitin	Amentoflavone	
Flow rate	1 ml/min	0.8ml/min	0.68	0.05	
		1.2ml/min	0.60	0.32	
Wavelength	280nm	275nm	0.40	0.50	
		285nm	0.61	0.15	
Temperature	25°C	20°C	1.02	0.09	
		30°C	1.80	0.33	
<b>Semecarpus anacardium methanolic leaf extract</b>					
Flow rate	1 ml/min	0.8ml/min	0.06	0.06	
		1.2ml/min	0.22	0.11	
Wavelength	280nm	275nm	0.29	0.03	
		285nm	0.21	0.14	
Temperature	25°C	20°C	0.06	0.06	
		30°C	0.32	0.13	

## RESULTS

A satisfactory separation and good peak symmetry was obtained with Kromasil 100-5C18 (250x 4.6, 5  $\mu\text{m}$ ) column and mobile phase comprising of Water of pH 3: Acetonitrile (55:45 v/v).

The pH of water was adjusted at 3.00 using O-phosphoric acid. The flow rate was 1.0 ml/min to get better reproducibility and repeatability. Quantification was achieved with UV detection at 280nm based on peak area. The retention time was found to be 4.55 min for Quercitin and 6.29 min for Amentoflavone. Correlation coefficient (r) was found to be 0.9949 for Quercitin & 0.9964 for Amentoflavone respectively. The optimised method was validated as per ICH guidelines.

## CONCLUSION

A specific, precise, accurate, sensitive, rapid and reliable Reverse phase HPLC method has been developed and validated. It has runtime of 10 min and retention time 4.55 min for Quercitin and 6.29 min for Amentoflavone. Short run time allows analysis of large number of samples in a less period of time. So this RP-HPLC method can be used in the quality control department for estimation of Quercitin and Amentoflavone.

## REFERENCES

- Kurup PNV, Ramdas VNK, Joshi P. Handbook of Medicinal Plants. New Delhi India Indian Council of Agricultural research 1979.
- Raghunathan S, Mitra R. Pharmacognosy of Indigenous Drugs. Vol I- II. New Delhi, India: Oxford & IBH Publishing; 1982.
- Sivarajan VV, Indira B. Ayurvedic Drugs and their plant sources. New Delhi, India: Oxford & IBH Publishing; 1994.p.184.
- Pednekar PA, Raman B. Pharmacognostic and phytochemical studies of *Semecarpus anacardium* (Linn.f.) leaves. Int J Pharm Pharm Sci 2012; 4(3): 682-685.
- Pednekar PA, Raman B. Assessment of *Semecarpus anacardium* (Linn.f.) leaf methanolic extract for their antibacterial, antifungal and antioxidant activity. Int J Pharm Pharm Sci 2013;5,(1):170-74.
- Pednekar PA, Raman B. The FT-IR spectrometric studies of vibrational bands of *Semecarpus anacardium* Linn.f. leaf, stem powder and extracts. Asian J Pharm Clin Res 2013; 6 Suppl 1: 159-198.
- Pednekar PA, Raman B. Multielement determination in methanolic soxhlet leaf extract of *Semecarpus anacardium* (Linn.f.) by ICP-AES technique. Asian J Pharm Clin Res 2013; 6 Suppl 3: 132-137.
- Lee E, Shin S, Kim JK, Woo ER, Kim Y. Anti-inflammatory effects of Amentoflavone on modulation of signal pathways in LPS-stimulated RAW264.7 cells. Bull. Korean Chem. Soc. 2012;33(9):2878-82.
- Kaikabo AA, Eloff JN. Antibacterial activity of two biflavonoids from *Garcinia livingstonei* leaves against *Mycobacterium smegmatis*. Journal of ethnopharmacology 2011;138(1):253-5.
- Gambhir SS, Goel RK, Dasgupta G. Anti-inflammatory and antiulcerogenic activity of Amentoflavone. Indian J Med Res 1987; 85: 689- 693.
- Kim HK, Son KH, Chang HW, Kang SS, Kim HP. Amentoflavone, a plant biflavone: a new potential anti-inflammatory agent. Archives of pharmacol research 1998;21(4):406-10.
- Tordera M, Ferrandiz ML, Alcaraz MJ. Influence of anti-inflammatory flavonoids on degranulation and arachidonic acid release in rat neutrophils. Z Naturforsch C 1994; 49(3-4): 235-240.
- Huguet AI, Manez S, Alcaraz MJ. Superoxide scavenging properties of flavonoids in a non-enzymic system. Z Naturforsch C 1990; 45(1-2): 19-24.
- Lee E, Shin S, Lee JY, Lee S, Kim JK, Yoon DY. Cytotoxic activities of Amentoflavone against human breast and cervical cancers are mediated by increasing of PTEN expression levels due to Peroxisome proliferator-activated receptor  $\gamma$  activation. Bull. Korean Chem. Soc 2012;33(7):2219-23.
- Geiger H. The Flavonoids: Advances in Research since 1986. Harborne JB ed. London: Chapman and Hall; 1994.p.95- 115..

16. Mora A, Payá M, Ríos JL, Alcaraz MJ. Structure-activity relationships of polymethoxyflavones and other flavonoids as inhibitors of non-enzymic lipid peroxidation. *Biochemical pharmacology* 1990;40(4):793-7.
17. Cholbi MR, Paya M, Alcaraz MJ. Inhibitory effects of phenolic compounds on CCl<sub>4</sub>-induced microsomal lipid peroxidation. *Experientia* 1991;47(2):195-9.
18. Shi S, Zhou H, Zhang Y, Huang K. Hyphenated HSCC-DPPH for rapid preparative isolation and screening of antioxidants from *Selaginella moellendorffii*. *Chromatographia* 2008; 68: 173-178.
19. Silva GL, Chai H, Gupta MP, Farnsworth NR, Cordell GA, Pezzuto JM, et al. Cytotoxic biflavonoids from *Selaginella willdenowii*. *Phytochemistry* 1995;40(1):129-34.
20. Lee HS, Oh WK, Kim BY, Ahn SC, Kang DO, Shin DI, et al. Inhibition of phospholipase C gamma 1 activity by amentoflavone isolated from *Selaginella tamariscina*. *Planta medica* 1996;62(4):293-6.
21. Lin LC, Kuo YC, Chou CJ. Cytotoxic biflavonoids from *Selaginella delicatula*. *Journal of natural products* 2000;63(5):627-30.
22. Guruvayoorappan C, Kuttan G. Effect of amentoflavone on the inhibition of pulmonary metastasis induced by B16F-10 melanoma cells in C57BL/6 mice. *Integrative cancer therapies* 2007;6(2):185-97.
23. Baureithel KH, Büter KB, Engesser A, Burkard W, Schaffner W. Inhibition of benzodiazepine binding in vitro by amentoflavone, a constituent of various species of *Hypericum*. *Pharmaceutica acta Helvetiae* 1997;72(3):153-7.
24. Gil B, Sanz MJ, Terencio MC, Gunasegaran R, Payá M, Alcaraz MJ. Morelloflavone, a novel biflavonoid inhibitor of human secretory phospholipase A2 with anti-inflammatory activity. *Biochemical pharmacology* 1997;53(5):733-40.
25. Woo ER, Lee JY, Cho IJ, Kim SG, Kang KW. Amentoflavone inhibits the induction of nitric oxide synthase by inhibiting NF-kappaB activation in macrophages. *Pharmacological research : the official journal of the Italian Pharmacological Society* 2005;51(6):539-46.
26. Jung HJ, Park K, Lee I-S, Kim HS, Yeo S-H, Woo E-R, et al. S-phase accumulation of *Candida albicans* by anticandidal effect of amentoflavone isolated from *Selaginella tamariscina*. *Biological & pharmaceutical bulletin* 2007;30(10):1969-71.
27. Lin YM, Flavin MT, Schure R, Zembower DE, Zhao GX. 1998. Biflavonoids and derivatives thereof as antiviral agents. United States Patent 5773462 (June 30, 1998).
28. Lin YM, Flavin MT, Schure R, Chen FC, Sidwell R, Barnard DL, et al. Antiviral activities of biflavonoids. *Planta medica* 1999;65(2):120-5.
29. Lin YM, Flavin MT, Schure R, Zembower DE, Zhao GX. 1999b. Biflavonoids and derivatives thereof as antiviral agents. United States Patent 5948918 (September 7, 1999).
30. Lin YM, Zembower DE, Flavin MT, Schure R, Zhao GX. 2002. Biflavonoids and derivatives thereof as antiviral agents. United States Patent 6339654 B1 (June 4, 2002).
31. Flavin MT, Lin YM, Zembower DE, Zhang H. Robustaflavone, intermediates and analogues and method for preparation thereof and a method for synthesizing the same are provided. United States Patent 6225481 May 2001;1.
32. Flavin MT, Lin YM, Zembower DE, Schure R, Zhao GX. 2002. Biflavonoids and derivatives thereof as antiviral agents. United States Patent 6399654 (June 4, 2002).
33. Ma SC, But PP, Ooi VE, He YH, Lee SH, Lee SF, et al. Antiviral amentoflavone from *Selaginella sinensis*. *Biological & pharmaceutical bulletin* 2001;24(3):311-2.
34. Kang DG, Yin MH, Oh H, Lee DH, Lee HS. Vasorelaxation by amentoflavone isolated from *Selaginella tamariscina*. *Planta medica* 2004;70(8):718-22.
35. Cassels BK, Dajas FJ, Medina JH, Paladini AC, Silveira RH. 1998. Flavonoid and biflavonoid derivatives, their pharmaceutical compositions, their anxiolytic activity. United States Patent 5756538 (May 26, 1998).
36. Cassels BK, Dajas FJ, Medina JH, Paladini AC, Silveira RH. 1999. Flavonoid and biflavonoid derivatives, their pharmaceutical compositions, their anxiolytic activity. United States Patent 6004998 (December 21, 1999)
37. Silva KLDA, dos Santos AR, Mattos PE, Yunes RA, Delle-Monache F, Cechinel-Filho V. Chemical composition and analgesic activity of *Calophyllum brasiliense* leaves. *Therapie* 2001; 56: 431-434.
38. Lee JS, Lee MS, Oh WK, Sul JY. Fatty acid synthase inhibition by amentoflavone induces apoptosis and antiproliferation in human breast cancer cells. *Biological & pharmaceutical bulletin* 2009;32(8):1427-32.
39. Burda S, Oleszek W. Antioxidant and antiradical activities of flavonoids. *Journal of agricultural and food chemistry* 2001;49(6):2774-9.
40. Husain SR, Cillard J, Cillard P. Hydroxyl radical scavenging activity of flavonoids. *Phytochemistry* 1987; 26: 2489-2491.
41. Rozenfel'd LG, Abyzov RA, Bozhko GT et al. The possibilities of protection against local radiation injuries in ORL-oncologic patients. *Vestn Otorinolaringol* 1990; 2:56-58.
42. Quercetin (Monograph). *Altern Med Rev* 1998; 3:140-143.
43. Belinha I, Amorim MA, Rodrigues P, de Freitas V, Moradas-Ferreira P, Mateus N, et al. Quercetin increases oxidative stress resistance and longevity in *Saccharomyces cerevisiae*. *Journal of agricultural and food chemistry* 2007;55(6):2446-51.
44. Yang J-H, Hsia T-C, Kuo H-M, Chao P-DL, Chou C-C, Wei Y-H, et al. Inhibition of lung cancer cell growth by quercetin glucuronides via G2/M arrest and induction of apoptosis. *Drug metabolism and disposition: the biological fate of chemicals* 2006;34(2):296-304.
45. ICH Harmonised Tripartite Guideline, Q2 (R1): Validation of Analytical Procedures: Text and Methodology Current Step 4 version, 2005.