

**PHYTOCONSTITUENTS, PIGMENTS, GAS CHROMATOGRAPHY MASS SPECTROMETRY ANALYSIS, AND ALLELOPATHY EFFECT OF *ALTERNANTHERA FICOIDEA* (L.) P. BEAUV**

RESHMA B PATIL\*, BASAVARAJ A KORE

Department of Botany, Yashwantrao Chavan Institute of Science, Satara - 415 001, Maharashtra, India. Email: reshmagodse09@gmail.com

Received: 04 August 2016, Revised and Accepted: 27 October 2016

**ABSTRACT**

**Objective:** The aim was to investigate phytoconstituents, composition of pigments, gas chromatography mass spectrometry (GCMS) and allelopathy of *Alternanthera ficoidea* (L.) P. Beauv.

**Methods:** Qualitative phytochemical analysis was carried out by the method of Paech and Tracey using five different solvent systems (D.W, ether, chloroform, ethanol, and methanol). Methanol leaf extract of this plant was analyzed using GCMS. Chlorophylls, carotenoid, and polyphenols were estimated by following standard methods. Allelopathy effect was studied using leaf leachets on seeds of jowar and mung.

**Result and Conclusion:** Investigation of above parameters in *A. ficoidea* showed presence of large amount of pharmaceutically important phytochemicals like tannins, saponins, phytols, carotenoids, xanthophylls and polyphenols. It indicates that this weed can be used in herbal medicines and dietary supplements. Since this weed is showing allelopathy effect it should be eradicated from fields before it spreads and occupies the place meant for crop plants.

**Keywords:** Allelopathy, *Alternanthera ficoidea*, Exotic weed, Phytochemicals.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i2.14498>

**INTRODUCTION**

The plants are capable of activating defense mechanisms against other plants by producing hundreds of secondary metabolites. These substances have received special attention because of their agricultural potential as either natural herbicides or templates for new synthetic herbicides [1]. Polyphenols have a major role in defense mechanism and carotenoids protect chlorophyll from cellular damaging. A pigment is a molecule that absorbs and reflects light. The broad array of colors found in plant tissues such as leaves, flowers, and fruits, can be accumulated for by the presence of thousands of different kinds of plant pigments. Chlorophylls and carotenoids have been considered to be responsible for the color of green plants [2-4]. Chlorophylls are of interest to agriculture and ecology where they are indicators of the health status of individual plants and communities and are often used as a quantitative reference in physiological research. They are also permitted as food colors [5]. Chlorophylls have recently attracted interest as phototherapeutic drugs [6,7]. Many lines of research suggest that consuming diet rich in plant pigments may slow the process of cellular aging and reduce the risks of some types of diseases such as cancer, heart diseases, and stroke [8]. Identification of natural plant components could contribute to the discovery of allelopathic agents. Allelopathy is a sophisticated mechanism of plant defense.

Amaranthaceae comprises 65 genera and approximately 1000 described species that originate from the tropical, subtropical and temperate zones of Africa, South America, and Southeast Asia. The genus *Alternanthera* is a prominent member of this family and includes 80 species. *Alternanthera ficoidea* (L.) P.Beauv (synonym *Alternanthera tenella* Colla) [9] is an invasive alien weed found everywhere in India.

**METHODS**

Specimens were collected from Satara and adjoining areas identified with standard literature and expert taxonomist [10].

**Preparation of samples for phytochemical analysis**

Leaf, stem, and roots were shade dried for 3 weeks and then pulverized using an electric blender. The powdered material was stocked in an airtight glass container. A total of 1 g of dried leaf material was extracted in 100 ml each of cold distilled water, ether, chloroform. Ethanol and methanol for 48 hrs at 37°C. This was then filtered with Whatman No. 1 filter paper. The filtrate was concentrated on a steam bath to obtain extracts used for qualitative phytochemical analysis.

**Preliminary screening**

The phytochemical screening of the plant extract was carried out by following the method of Paech and Tracey [11]. MeOH extract was used for gas chromatography mass spectrometry (GCMS) analysis.

**Pigment and polyphenol estimation**

Chlorophylls [12], carotenoid [13], and polyphenols [14] were estimated by following standard methods.

**Preparation of leaf leachete for study of allelopathy**

Leaf leachete was prepared by soaking 100 g of dry leaf powder in 500 ml of distilled water for 24 hrs. Then, filter through Whatman's No. 1 filter paper and allowed to evaporation. The same powder reextracted twice in fresh water and evaporated filtrate is pooled to 100 ml with D.W. and stored in refrigerator in amber colored bottle for further use, leaf leachetes of 1%, 2%, 5%, 10%, 15%, and 25% concentrations were prepared by further dilutions of stock solution.

**Seed germination bioassay**

The healthy seeds of mung bean and jowar were used for seed germination bioassay. Seeds were surface sterilized with 0.02% aqueous mercuric chloride for 2 minutes, thoroughly washed with distilled water and 20 seeds were kept in each sterilized petriplate (9 cm diameter) lined with filter paper. The filter papers in petriplates were moistened with 10 ml of respective concentrations (1%, 2%, 5%, 10%, 15%, and

25%) of leaf leachet. The seeds placed in petri plates moistened with 10 ml distilled water were considered as control. The experiment was arranged in triplicate by using seed germination chamber at 26°C and natural light and dark period. A number of germinated seeds were recorded every day. The germination percentage, root and shoot length, vigor index, fresh, and dry weight of seedlings were recorded on the 7<sup>th</sup> day. Mean time for germination is an index of seed germination speed and velocity, and mean daily germination (MDG) an index of daily germination speed and calculated by germination speed formula [15] was calculated for a week.

## RESULTS AND DISCUSSION

Plants are rich in a wide variety of secondary metabolites tannins, saponins, quinone, anthraquinone, steroids, glycosides, terpenoids, triterpenoids, flavonoids, alkaloids, coumarins, phenolic compounds, and essential oils with antimicrobial properties [16,17]. Plant synthesizes a wide range of chemical compounds which are classified based on their chemical class, biosynthetic origin and functional groups into primary and secondary metabolites.

Secondary metabolites are synthesized during secondary metabolism of plants. Phenolic compounds are the most widely distributed secondary metabolites, ubiquitous in the plant kingdom. The great majority of active phenolic compounds isolated from higher plants are flavonoids and phenolic acids. Studies on phenolic compounds have shown a wide range of biological activities such as anti-inflammatory, hepatoprotective, antioxidant, antitrombotic, vasodilating, and anticarcinogenic [18,19]. All secondary metabolites have a specific function as like saponins have antifungal activity [20].

The previous preliminary phytochemical analysis showed presence of all type of phytochemicals while tannins and saponins were detected in moderate amount [21]. Preliminary phytochemical analysis of *A. ficoidea* shows the presence of alkaloids, flavonoids, cardiac glycosides, terpenoids, tannins, anthraquinones, and saponins (Table 1).

GCMS analysis confirmed the results as shown in Table 2. It showed the presence of five major groups. 3,7,11,15-tetramethyl-2-hexadecan-1-ol6, 9,12,15-docosatetraenoic acid, 9,12,15-octadecatrienoic acid, methyl ester and phytol are the groups which are pharmaceutically important. Phytol is one among the five compounds of the present study. 9,12,15-octadecatrienoic acid, (Z,Z,Z)-linolenic acid (R/T 20.06) possesses anti-inflammatory, insectifuge, hypocholesterolemic, cancer preventive, nematocidal, hepatoprotective, antihistaminic, antieczemic, antiacne, 5-alpha reductase inhibitor, antiandrogenic, antiarthritic and anticoronary properties. n-Hexadecanoic acid - palmitic acid (R/T 17.25) can be an antioxidant, hypocholesterolemic, nematocidal, pesticide, lubricant activities and hemolytic 5-alpha is a reductase inhibitors. Phytol-diterpene (R/T 19.67) is an antimicrobial, anticancer, anti-inflammatory and diuretic agent [22] as in *Cassia italic* [23]. Rani *et al.* [24] observed the presence of phytol in the leaves of *Lantana camara* and Sridharan *et al.* [25] in *Mimosa pudica* leaves. A similar result was also observed in the leaves of *L. camara* [26]. It was observed to have antibacterial activities against *Staphylococcus aureus* by causing damage to cell membranes as a result, there is a leakage of potassium ions from bacterial cells [27] phytol, phenol, 2, 4-bis (1-phenylethyl) - which are all have medicinal properties. Phytol is a key acyclic diterpene alcohol that is a precursor for vitamins E and K1. It is used along with simple sugar or corn syrup as a hardener in candies. Mangunwidjaja *et al.* [28] reported the main components of 9,12 octadecadienoic acid, octadec-9-enoic acid and 9,12-actadecadienoic acid present in *Croton tiglium* seed. These compounds were found to have potential antioxidant and anticancer activities. Polyphenol content is maximum in summer in post-reproductive phase. These plants are also rich in minerals like potassium [29].

Pigments are integrally related to the physiological function of leaves. Chlorophylls absorb light energy and transfer it to the photosynthetic apparatus. Carotenoids (yellow pigments) can also contribute energy

to the photosynthetic system. However, when incident light energy exceeds that needed for photosynthesis, the carotenoids that compose the xanthophyll cycle dissipate excess energy, thus avoiding damage to the photosynthetic system [30]. Because of the importance of pigments for leaf function, variations in pigment content may provide information concerning the physiological state of leaves. Chlorophyll tends to decline more rapidly than carotenoids when plants are under stress or during leaf senescence [31]. When we investigate pigments in pre-reproductive, reproductive and post-reproductive stages at summer and rainy season as shown in Table 3, it shows variability among chlorophyll a, chlorophyll b, and total chlorophyll. Chlorophyll level is more in pre-reproductive stage during summer while carotenoid and xanthophyll content is more in reproductive phase during summer.

The xanthophyll cycle carotenoids seem to be highly flexible, adjusting rapidly to light conditions [32]. In addition to daily changes, a seasonal pattern has been described, with major changes in the V + A + Z pool content and composition occurring between summer and winter in some evergreen coniferous trees (Adams) carotenoids provide colors to flowers, seeds, fruit, and to some fungi, and color has an important role in reproduction: Coloration attracts animals that disperse pollen, seeds, or spores. Main pigments involved in photosynthesis are chlorophylls and carotenoids. Carotenoids have two well-known functions in photosynthesis: Accessory pigments in light harvesting [33], and as photoprotectors against oxidative damages [34]. It has been proposed

Table 1: Qualitative phytochemical analysis of *A. ficoidea*

Phytochemical tests	Source of sample	Solvent used				
		D.W.	Ether	Chloroform	EtOH	MeOH
<b>Alkaloid</b>						
Marqui's test	Leaf	-	+++	+++	++	++
	Stem	-	+++	++	+	++
	Root	-	+++	++	+	-
Hager's test	Leaf	-	+++	++	+	-
	Stem	-	+++	++	+	+
	Root	-	+++	++	+	+
<b>Flavonoid</b>						
Shinoda test	Leaf	++	-	-	-	-
	Stem	+	-	-	-	-
	Root	+	-	-	-	-
Ammonia test	Leaf	++	-	-	-	-
	Stem	+	-	-	-	-
	Root	+	-	-	-	-
<b>Cardiac glycosides</b>						
Baljet test	Leaf	+	+	++	+++	+++
	Stem	-	+	+	++	++
	Root	-	-	+	+	+
<b>Terpenoid</b>						
Salkovashi test	Leaf	++	-	-	++	+++
	Stem	+	-	-	+	++
	Root	+	-	-	+	++
Test for tannin	Leaf	+++	+++	+++	+++	+++
	Stem	+++	+++	+++	+++	+++
	Root	+++	+++	+++	+++	+++
Test for reducing sugar	Leaf	-	-	+++	+	+
	Stem	-	-	+++	+	+
	Root	-	-	+++	+	+
Test for anthraquinones	Leaf	-	+++	-	-	-
	Stem	-	+++	-	-	-
	Root	-	+++	-	-	-
Test for saponins	Leaf	+++	-	+++	+++	+
	Stem	+++	-	+++	+++	+
	Root	+++	-	+++	+++	+

-, Absent, +, Present in small amount, ++, Moderately present, +++, Present in large amount, *A. ficoidea*: *Alternanthera ficoidea*

Table 2: GCMS analysis of *A. ficoidea*

S.No.	Name of the compound	Molecular formula	Molecular weight	Retention time	Area %	Activity
1	3-ethoxy-1,1,1,7,7,7-hexamethyl-3,5,5-tris(trimethylsiloxy) tetrasiloxane	C <sub>17</sub> H <sub>50</sub> O <sub>7</sub> Si <sub>7</sub>	562	18.192	17.98	Unknown
2	3,7,11,15-tetramethyl-2-hexadecen-1-ol	C <sub>20</sub> H <sub>40</sub> O	296	26.333	19.23	Antimicrobial
3	6,9,12,15-docosatetraenoic acid, methyl ester	C <sub>23</sub> H <sub>38</sub> O <sub>2</sub>	346	26.525	5	Unknown
4	9,12,15-octadecatrienoic acid, methyl ester	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	292	26.658	13.43	Antiinflammatory, nematocide, insectifuge, hypocholesterolemic, cancer preventive, hepatoprotective, antihistaminic, antiacne, antiarthritic, antieczemic
5	Phytol	C <sub>20</sub> H <sub>40</sub> O	296	26.825	44.35	Antimicrobial, anticancer, cancer preventive, diuretic antiinflammatory

*A. ficoidea*: *Alternanthera ficoidea*, GCMS: Gas chromatography mass spectrometry

Table 3: Pigment and polyphenol analysis at three different life stages and two seasons

Life stage	Season	Chlorophyll a g/100 g	Chlorophyll b g/100 g	Total chlorophyll g/100 g	Carotenoids g/100 g	C <sub>x+c</sub> g/100 g	Polyphenols g/100 g
Pre reproductive	Rainy	0.140	0.0739	0.214	0.242	0.562	4.5
Reproductive	Summer	0.157	0.112	0.269	0.314	0.952	8.08
	Rainy	0.156	0.0862	0.242	0.260	0.639	5.23
	Summer	0.134	0.121	0.255	0.311	0.709	9.16
Post reproductive	Rainy	0.034	0.0263	0.060	0.070	0.230	6.08
	Summer	0.142	0.113	0.255	0.303	0.750	10.16

that carotenoids as light harvesting compounds evolved from anaerobic organisms, then generalized to all of the aerobic photosynthetic organisms. The physical structure of chloroplasts facilitates the transference of energy absorbed by carotenoids to chlorophyll. When leaves are exposed to high illumination, epoxy xanthophyll groups are removed of violaxanthin to initially form antheraxanthin and then zeaxanthin. This is one of the plant protection mechanisms against light damage. The number of carotenoid molecules is higher in sun-exposed leaves than darkness maintained leaves.

This phenomenon is very important, sun-exposed leaves in a fast-growing stage use not more than 50% of absorbed energy during the stage of maximum radiation (midday), and in some species only 10% is used. Thus, 50-90% of absorbed light is in excess and must be eliminated to avoid cellular damage. Xanthophyll cycle is a process that makes the energy dissipation easy and protects the photosynthetic apparatus observed. Moreover, xanthophyll cycle carotenoids are associated with the energy-harvesting complexes photosystem I (PSI) and PSII [35,36]. It was found that higher carotenoid concentrations decreased the peroxide value of soybean oil by quenching singlet oxygen, and it was shown that longer chromophores favor this reaction. Packer [37] evaluated the antioxidant activity of different carotenoids by *in vitro* assays. It was reported that antioxidant activity depends on the used system. Carotenoids, by their antioxidant effect, can show benefits in such diseases; however, this function is not completely demonstrated *in vivo*. Carotenoids have been considered that provide benefits in age-related diseases, against some forms of cancer (in especial lung cancer), strokes, macular degeneration, and cataracts. However, most studies relate dietary components with sickness incidence or symptoms; thus, these studies cannot establish a direct cause-effect relationship. On the other hand, it is clear that carotenoids in association with other components of fruits and vegetables seem to have a protective effect against some chronic diseases and precancerous conditions. Carotenoids protect lab animals of ultraviolet-induced inflammation and a certain type of cancers.

Historically, carotenoid supplementation has been used in the treatment of diseases produced by light sensitivity, which is usually hereditary: 84% of patients with erythropoietic protoporphyria, consuming diets supplemented with  $\beta$ -carotene, increased by a factor of 3 their ability to

resist sunlight exposition without presenting symptoms. Furthermore, carotenoids have been used in other photosensitivity diseases: Congenital porphyria, sideroblastic anemia, and have shown only limited success in the treatment of polymorphic light eruption, solar urticaria *Hydroa vacciforme*, *Porphyria variegata*, *Porphyria cutanea tarda*, or actinic reticuloid [38,39].

The invasive weeds infesting the crop fields and other ecosystems have become a serious problem today. The agroecosystems show an association of various types of native and invasive weeds in crops and barren lands as well as wastelands. To demonstrate the allelopathic potential of selected native and invasive weeds, seed germination bioassays were conducted using the seeds of test crops such as mung and jowar. The protection and conservation of native plant species can be achieved through studies on ecophysiology, allelopathy, and biological invasion of weeds. Scanty research is done on isolation, identification, and characterization of allelochemicals in different native and invasive weeds. Considering these facts, the present investigation was attempted to focus on allelopathic studies of dominant, invasive weed species *A. ficoidea*. The negative impact of higher concentrations of extracts and leachates of different weeds on various crops had been well documented by several workers such as Mallik and Pellissier [40], Djanaguiraman *et al.* [41], Duary [42], Patil *et al.* [43], Ali *et al.* [44], Punjani [45], Travlos and Paspatis [46], Aziz *et al.* [47], and Tanveer *et al.* [48]. They have reported significant inhibition of root and shoot length in *Picea*, wheat, maize, sorghum, rice, blackgram, greengram, soybean, cowpea, cotton and sunflower due to higher concentration treatments of *Vaccinium*, *Eupatorium*, *Sesamum*, *Prosopis*, *Eucalyptus*, *Acacia*, *Grewia*, *Populus*, *Casuarina*, *Amaranthus*, and *Parthenium*.

*A. ficoidea* shows inhibitory effect on germination of mung and jowar. Stress declined the germination and also delayed the germination process. The negative allelopathic activity increases with increase in leaf leachet concentration. Maximum inhibition was shown by 25% leaf leachete from concentration 1%, 2%, 5%, 10%, 15%, and 25% on germination indices such as germination percentage, plumule length, radical length, plumule fresh weight, plumule dry weight, radicle fresh weight, radicle dry weight, meantime germination, MDG, daily germination speed, seedling length vigour index, and seedling weight vigor index also shows

allelopathic. As reported above, the phenolic compounds present in the invasive weeds might have caused inhibition of the shoot and root growth in mung bean, because higher concentrations of phenols were detected in the leaves and roots of these invasive weeds. The results on dry biomass of treated seedlings in mung bean and jowar indicate decrease due to the application with higher concentration treatments (Tables 4 and 5). However, Mallik and Pellissier, Duary *et al.* and Djanaguiraman *et al.*, Lodha [49] and Ali *et al.* have reported considerable inhibitory effects on dry weight in wheat, rice, black gram, mung bean, cowpea, and *Picea* due to higher concentrations of aqueous leaf extracts and decomposing litter of sunflower, *Eucalyptus*, *Amaranthus*, *Parthenium*, *Sphaeranthus*, sesame, and *Vaccinium*, respectively.

The level of inhibition increased with increase in the concentration of the extracts for all the crops tested. In the present investigation, similar results were obtained regarding the decrease in the dry weight of mung bean and jowar seedlings due to the lower to higher concentration treatment of leaf extracts of *Alternanthera*. This concentration-dependent increase or decrease in dry biomass may be attributed to the explanations given by An *et al.* [50], Orr *et al.* [51] recorded the potential allelopathic influence of two invasive weed species like *Lolium* and *Elaeagnus* on native species of *Acer*, *Populus*, and *Platanus*. They found reduction and delay in seedling emergence, root and leaf biomass in treated plants. They explained that it was due to the action of allelochemicals. Swain *et al.* [52] had also reported that the leachates of root, stem, and leaves of *Physalis* inhibited seed germination of *Parthenium* at higher concentrations.

They have claimed that phenolics were responsible for the reduction in seed germination and seedling growth. According to Todaria *et al.*

(2005) the tannins, phenols, and other secondary metabolites were responsible for inhibition of seed germination in treated plants. Travlos and Paspatis, Aziz *et al.* and Tanveer *et al.* also observed similar inhibition of germination of their respective test crops. Einhellig (1996) [53], Reigosa *et al.* [54] claimed that the allelochemicals present in the extracts, leachates or residues are responsible for the changes in water relations, membrane permeability and enzymatic activities of protein and carbohydrate metabolism during seed germination process, which determine the success of seed germination. It may be the reasons for the stimulation of seed germination in mung bean when, treated with leaf extracts of *Alternanthera*, *Croton* and *Xanthium* [55].

Lodha and Ali *et al.* have reported considerable inhibitory effects on dry weight in wheat, rice, black gram, mung bean, cowpea and *Picea* due to higher concentrations of aqueous leaf extracts and decomposing litter of sunflower, *Eucalyptus*, *Amaranthus*, *Parthenium*, *Sphaeranthus*, sesame, and *Vaccinium*, respectively. The level of inhibition increased with increase in the concentration of the extracts for all the crops tested. In the present investigation, similar results were obtained regarding the increase or decrease in the dry weight of mung bean and jowar seedlings due to the lower and higher concentration treatment of leaf extracts of *A. ficoidea*. The inhibition or stimulation may be due to various types of allelochemicals such as terpenoids, high phenolic compounds present in the extracts of *Alternanthera*.

Allelopathy may provide alternatives to synthetic herbicides for weed control [56,57]. Laboratory bioassay is the first step to investigate the probable involvement of allelopathy [58] aqueous extract bioassays have been widely employed to evaluate allelopathy of a suspected donor

Table 4: Allelopathic effect of *A. ficoidea* on mung seeds

S.No.	Concentrated of leaf leachet/aspects studied (Mung)	Control D.W.	1%	2%	5%	10%	15%	25%
1	Germinated seeds/10 seeds	10	10	9.66	10	10	10	8
2	Germination (%)	100	100	96.6	100	100	100	80
3	Radicle length (cm)	14	3.73	2.76	2.43	2	2	1.5
4	Plumule length (cm)	3.56	1.73	0.96	0.15	0.23	0.1	-
5	Radicle fresh weight (g)	7.62	5.14	4.61	3.7	2.336	0.2	0.05
6	Radicle dry weight (g)	0.885	0.524	0.594	0.0258	0.189	0.005	0.01
7	Plumule fresh weight (g)	20.70	11.76	9.05	9.04	9	6.43	-
8	Plumule dry weight (g)	1.19	1.74	1.70	1.87	1.184	0.73	-
9	MTG/7 days	2	2.5	2.17	5.8	5.8	5.8	0.91
10	MDG/20 seeds	14.28	14.28	13.80	14.28	14.28	14.28	11.42
11	DGS	0.070	0.070	0.072	0.070	0.070	0.070	0.087
12	SLVI	1756	446	372	268	223	200	160
13	SWVI	2832	1690	1320.37	1240	1143	714	408.8

SWVI: Seedling weight vigor index, SLVI: Seedling length vigor index, DGS: Daily germination speed, MDG: Mean daily germination, MTG: Mean time germination, *A. ficoidea*: *Alternanthera ficoidea*

Table 5: Allelopathic effect on Jowar seeds

S.No.	Concentrated of leaf leachet/aspects studied (Jowar)	Control D.W.	1%	2%	5%	10%	15%	25%
1	Germinated seeds/10 seeds	9.66	9.66	8.66±0.72	7.66	7.66	6.66	6.33
2	Germination (%)	96.6	96.6	86.6	76.6	76.6	66.6	63.3
3	Radicle length (cm)	12.83	6.06	3.16	3	2.4	1.96	0.3
4	Plumule length (cm)	6.03	2.2	1.633	1.43	1.26	1.07	0.1
5	Radicle fresh weight (g)	4.53	1.76	1.34	1.14	0.82	0.469	0.09
6	Radicle dry weight (g)	0.568	0.391	0.28	0.18	0.177	0.147	0.05
7	Plumule fresh weight (g)	8.43	2.53	2.06	1.94	2.1	1.44	0.5
8	Plumule dry weight (g)	1.07	0.455	0.431	0.397	0.385	0.26	0.09
9	MTG/7 days	2.06	2.85	2.69	2.56	2.90	2.65	0.6
10	MDG/20 seeds	13.8	13.8	12.38	10.95	10.95	9.52	7.14
11	DGS	0.0724	0.072	0.080	0.091	0.0913	0.105	0.14
12	SLVI	1823.006	798.41	369.338	280.356	280.36	203.13	150
13	SWVI	1296	414.67	294.64	235.928	223.67	127.206	53.77

SWVI: Seedling weight vigor index, SLVI: Seedling length vigor index, DGS: Daily germination speed, MDG: Mean daily germination, MTG: Meantime germination

species. The speed of germination considers the number of germinated seeds between two exposure times, whereas accumulated germination involves the cumulative number of germinated seeds at each exposure time. Results supported the hypothesis that data interpretations depends on the choice of germination index and also that one index might not be suitable for precise description of allelopathic effects on the germination process. Comparison of different indices may provide better justification. This will contribute in making allelopathy a more precise science.

## CONCLUSION

Investigation of *A. ficoidea* showed presence of large amount of pharmaceutically important phytochemicals like tannins, saponins, phytols, carotenoids xanthophylls and polyphenols. It indicates that this weed can be used in herbal medicines and dietary supplements. Since this weed is showing allelopathy effect it should be eradicated at all the stages from fields before it spreads and occupies the place meant for crop plants.

## ACKNOWLEDGMENTS

The authors are thankful to the Principal, Yashvantrao Chavan Institute Of Science, Satara for providing necessary laboratory facilities and encouraging during the current research work.

## REFERENCES

1. Chuah TS, Norhafizah MZ, Ismail S. Phytotoxic effects of the extracts and compounds isolated from Napier grass (*Pennisetum purpureum*) on Chinese sprangletop (*Leptochloa chinensis*) germination and seedling growth in aerobic rice systems. *Weed Sci* 2014;62(3):457-67.
2. Khachick F, Madlagiri B, Beecher GR, Holden J, Lubsy WR, Tenorio MD, et al. Effect of food preparation on qualitative and quantitative distribution of major carotenoid constituents of tomatoes and several green vegetables. *J Agric Food Chem* 1992;40(3):390-8.
3. Valverde J, This H, Vignolle M. Qualitative determination of photosynthetic pigments in green beans using thin-layer chromatography and flatbed scanner as densitometer. *J Chem Educ* 2007;84:1505-7.
4. Wrolstad RE. Handbook of Food Analytical Chemistry. 1<sup>st</sup> ed., Vol. 2. Hoboken, New Jersey: John Wiley & Sons, Inc.; 2005.
5. Guerra RN, Pereira HA, Silveira LM, Olea RS. Immunomodulatory properties of *Alternanthera tenella* Colla aqueous extracts in mice. *Braz J Med Biol Res* 2003;36(9):1215-9.
6. Scheer H. In: Chlorophylls. Boca Raton, FL: CRC Press; 1991.
7. Schwartz SJ, Lorenzo TV. Chlorophylls in foods. *Crit Rev Food Sci Nutr* 1990;29(1):11-7.
8. Clinton SK. Lycopene: Chemistry, biology, and implications for human health and disease. *Nutr Rev* 1998;56:35-51.
9. Anonymous the Plant List, 2014 - Version 1.1. Available from: <http://www.theplantlist.org>. [Last accessed on 2015 Sep 18].
10. Yadav SR, Sardesai MM. Flora of Kolhapur District. Kolhapur: Shivaji University; 2002.
11. Paech D, Tracey MV. Modern Methods of Plant Analysis. 1<sup>st</sup> ed., Vol. 3. Berlin: Springer Verlag; 1955. p. 64-5.
12. Arnon DI. Copper enzyme in isolated chloroplasts: Polyphenol oxidase in *Beta vulgaris*. *Plant Physiol* 1949;24(1):1-15.
13. Lichtenthaler HK, Wellburn AR. Determination of total carotenoids and chlorophylls A and B of leaf in different solvents. *Biol Soc Trans* 1985;47(11):591-2.
14. Folin O, Denis WA. Calorimetric estimation of phenols and phenol derivatives in urine. *J Biol Chem* 1915;22:305-8.
15. Abbasian A, Moemeni J. Effects of salinity stress on seed germination and seedling vigor indices of two halophytic plant species (*Agropyron elongatum* and *A. pectiniforme*). *Int J Agric Crop Sci* 2013;5(22):2669-76.
16. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999;12:564-82.
17. Lewin WH, Elvin-Lewis MP. Medicinal plants as sources of new therapeutics. *Ann Mo Bot Gard* 1995;82(1):16-24.
18. Bruneton J. Pharmacognosy, Phytochemistry. Medicinal Plants. 2<sup>nd</sup> ed. London, New York: Lavoisier; 1999.
19. Wang H, Provan GJ, Helliwell K. Determination of rosmarinic acid and caffeic acid in aromatic herbs by HPLC. *Food Chem* 2004;87(2):307-11.
20. Sodipo OA, Akanji MA, Kolawole FB, Odutuga AA. Saponin is the active antifungal principle in *Garcinia kola*, heckle seed. *Biosci Res Commun* 1991;3:171.
21. Poonguzhali TV, Ramani K. Qualitative phytochemical analysis of *Alternanthera tenella* and *Dipteracanthus prostrates*. *Int J Curr Res* 2012;4(12):210-1.
22. Kumar PP, Kumaravel S, Lalitha C. Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*. *Afr J Biochem Res* 2010;4(7):191-5.
23. Sermakkani M, Thangapandian V. GC-MS analysis of *Cassia italica* leaf methanol extract. *Asian J Pharm Clin Res* 2012;5(2):90-4.
24. Rani PM, Kannan PS, Kumaravel S. GC-MS analysis of *Lantana camara* L. leaves. *JPRD* 2011;2(11):63-6.
25. Sridharan S, Meena V, Kavitha V, Nayagam AA. GC-MS study and phytochemical profiling of *Mimosa pudica* Linn. *J Pharm Res* 2011;4(3):741-2.
26. Kumar MS, Manimegalai S. Evaluation of larvicidal effect of *Lantana camara* Linn against mosquito species *Aedes aegypti* and *Culex quinquefasciatus*. *Adv Biol Res* 2008;2(3-4):39-43.
27. Inoue Y, Hada TA, Shiraiishi K, Hirore H, Kobayashi S. Biphasic effects of geranylgeraniol, terpenone and phytol on the growth of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2005;49(5):1770-4.
28. Mangunwidjaja DS, Kardono SR, Iswantini L B. Gas chromatography and gas chromatography-mass spectrometry analysis of Indonesian *Croton tiglium* seeds. *J Appl Sci* 2006;6(7):1576-80.
29. Patil RB, Kore BA. Availability of mineral elements in an exotic weed *Alternanthera tenella* Colla var. *Tenella* Veldk. *Asian J Pharm Clin Res* 2015;8(1):73-5.
30. Demmig-Adams B, Adams WW 3<sup>rd</sup>. The role of xanthophylls cycle carotenoids in the protection of photosynthesis. *Trends Plant Sci* 1996;1(1):21-7.
31. Gitelson AA, Merzlyak MN. Signature analysis of leaf reflectance spectra: Algorithm development for remote sensing of chlorophyll. *J Plant Physiol* 1996;148(3-4):494-500.
32. Demmig-Adams B, Adams WW 3<sup>rd</sup>, Winter K, Meyer A, Schreiber U, Pereira JS, et al. Photochemical efficiency of photosystem II, photon yield of O<sub>2</sub> evolution, photosynthetic capacity, and carotenoid composition during the midday depression of net CO<sub>2</sub> uptake in *Arbutus unedo* growing in Portugal. *Planta* 1989;177(3):377-87.
33. Adewusi SR, Bradbury JH. Carotenoids in *Cassava*: Comparison of open-column and HPLC methods of analysis. *J Sci Food Agric* 1993;62:375-83.
34. Al-Babili S, von Lintig J, Haubruck H, Beyer P. A novel, soluble form of phytoene desaturase from *Narcissus pseudonarcissus* chromoplasts is Hsp70-complexed and competent for flavinylation, membrane association and enzymatic activation. *Plant J* 1996;9(5):601-12.
35. Armstrong GA, Hearst JE. Carotenoids 2: Genetics and molecular biology of carotenoid pigment biosynthesis. *FASEB J* 1996;10(2):228-37.
36. Olson JA, Krinsky NI. Introduction. The colorful, fascinating world of the carotenoids: Important physiologic modulators. *FASEB J* 1995;9:1547-50.
37. Packer L. Antioxidant action of carotenoids *in vitro* and *in vivo* and protection against oxidation of human low density lipoproteins. *Ann N Y Acad Sci* 1993;691:48-60.
38. Mathews-Roth MM. Carotenoids in erythropoietic protoporphyria and other photosensitivity disease. *Ann N Y Acad Sci* 1993;691:127-38.
39. Ziegler RG. Carotenoids, cancer, and clinical trials. In: Canfield LM, Krinsky NI, Dunastable JA, editors. Carotenoids in Human Health. Vol. 691. New York: Annals of the New York Academy of Sciences; 1993. p. 110-9.
40. Mallik AU, Pellissier F. Effects of *Vaccinium myrtillus* on spruce regeneration: Testing the notion of coevolutionary significance of allelopathy. *J Chem Ecol* 2000;26(9):2197-209.
41. Djanaguiraman M, Ravishankar P, Bangarusamy U. Effect of *Eucalyptus globulus* on greengram, blackgram and cowpea. *Allelopathy J* 2002;10(2):157-62.
42. Duary B. Effect of leaf extract of sesame (*Sesamum indicum* L.) On germination and seedling growth of black gram (*Vigna mungo* L.) And rice (*Oryza sativa* L.). *Allelopathy J* 2002;10(2):153-6.
43. Patil RH, Hunshal CS, Itnal CJ. Effect of *Casuarina* litter leachates on crops. *Allelopathy J* 2002;10(2):141-6.
44. Ali H, Kumar S, Abdulla MK, Sindhu G, Sindhu A. Allelopathic effect of *Amaranthus viridis* and *Parthenium hysterophorus* (L.) On wheat, maize and rice. *Allelopathy J* 2005;16(2):341-6.
45. Punjani BL. Allelopathic effects of *Prosopis chilensis* (Molina) Stuntz. on germination and seedling growth. *Allelopathy J* 2005;16(2):295-300.
46. Travlos IS, Paspatis EA. Allelopathic effects of heliotrope (*Heliotropium*

- europaeum* L.) on *Avena sativa*, *Phaseolus vulgaris*, and *Spirodela polyrhiza*. Allelopathy J 2008;21(2):397-404.
47. Aziz A, Tanveer A, Ali A, Yasin M, Babar BH, Nadeem MA. Allelopathic effect of cleavers (*Galium aparine*) on germination and early growth of wheat (*Triticum aestivum*). Allelopathy J 2008;22(1):25-34.
  48. Tanveer A, Tahir M, Nadeem MA, Younis M, Aziz A, Yaseen M. Allelopathic effects of *Xanthium strumarium* L. on seed germination and seedling growth of crops. Allelopathy J 2008;21(2):317-28.
  49. Lodha V. Germination and seedling vigor of some major crop plants as influenced by allelopathy of *Spheranthus indicus*. Indian J Plant Physiol 2004;9(2):195-8.
  50. An M, Pratley JE, Haig T, Liun DL. Whole-range assessment: A simple method for analysing allelopathic doseresponse data. Nonlinearity Biol Toxicol Med 2005;3(2):245-60.
  51. Orr SP, Rudgers JA, Clay K. Invasive plants can inhibit native tree seedlings: Testing potential allelopathic mechanisms. Plant Ecol 2005;181(2):153-65.
  52. Swain D, Pandey P, Paroha S, Singh M, Yeduraju NT. Effects of *Physalis minima* on *Parthenium hysterophorus*. Allelopathy J 2005;14:275-84.
  53. Einhellig FA. Mechanisms and modes of action of allelochemicals. In: Putnam AR, Tang CS, editors. The Science of Allelopathy. New York: John Wiley & Sons Inc.; 1996. p. 171-88.
  54. Reigosa MJ, Sánchez-Moreiras A, González L. Ecophysiological approach in allelopathy. Crit Rev Plant Sci 1999;18(5):577-608.
  55. Patil HS, Shitole SM, Dhupal KN. Effect of leaf and root extracts of selected weed species on seed germination and seedling growth in mung bean. Int J Curr Res 2013;5(1):94-8.
  56. Wu H, Pratley J, Lemerle D, Haig T. Crop cultivars with allelopathic capability. Weed Res 1999;39(3):171-80.
  57. Vyvyan JR. Allelochemicals as leads for new herbicides and agrochemicals. Tetrahedron 2002;58:1631-46.
  58. Foy CL. How to make bioassays for allelopathy more relevant to field conditions with particular reference to cropland weeds. In: Inderjit KM, Dakshni M, Foy CL, editors. Principles and Practices in Plant Ecology: Allelopathic Interactions. Washington: CRC Press; 1999. p. 25-33.