

HYPOGLYCEMIC ACTIVITY OF 10 MEDICINAL PLANTS EXTRACT IN GLUCOSE INDUCED MICEAHMAD MUHTADI^{1*}, YOLA IRENKA¹, WULAN CHANDRA AYU¹, RINI HENDRIANI¹, ADE ZUHROTUN²¹Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia. ²Department of Biology Pharmacy, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia. Email: a.muhtadi@unpad.ac.id

Received: 04 November 2016, Revised and Accepted: 24 January 2017

ABSTRACT

Objective: To evaluate and compare the hypoglycemic activity of 10 medicinal plants extract that is used empirically to treat hyperglycemia in Indonesia, to determine plant with the most potential as a hypoglycemic agent.

Methods: Activity test was conducted in Swiss Webster glucose-induced mice with glucose tolerance test. The ethanolic extracts of *Mangifera indica* L. leaves, *Persea americana* Mill. leaves, *Acorus calamus* L. rhizome, *Phyllanthus niruri* L. herb, *Syzygium cumini* (L.) skeels bark, *Zingiber officinale* Roscoe rhizome, *Moringa oleifera* Lam. leaves, *Tamarindus indica* L. seed, *Momordica charantia* L. fruits, and *Azadirachta indica* A. juss leaves, which were given orally at the same dose of 300 mg/kg bw. The blood samples were used for determination of glucose level examination using a glucose meter and test strips (Accu-Chek).

Results: The ethanolic extract of *M. indica* L. leaves has the best activity in reduction of blood glucose level in mice, followed by *Z. officinale* Roscoe rhizome, *A. calamus* L. rhizome, *T. indica* L. seed, *M. charantia* L. fruit, *S. cumini* (L.) skeels bark, *P. niruri* L. herb., *P. americana* Mill. leaves, and *A. indica* A. juss leaves.

Conclusion: Results of this study suggest the potential use of *M. indica* L. leaves extract in the therapy of hyperglycemia.

Keywords: Hypoglycemic, Medicinal plants, *Mangifera indica* L. leaves, Glucose tolerance test.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10s2.19473>

INTRODUCTION

Diabetes mellitus (DM) is a metabolic disease characterized by hyperglycemia, caused by lack of insulin production, insulin resistance, or both. It is generally classified into type 1 and 2 diabetes. Type 1 diabetes is insulin dependent for life (5-10%), occurs because of β pancreas cells damage. While type 2 diabetes (90-95%) occurs due to insulin resistance, insufficient production of insulin, or both [1]. This disease can also cause chronic complications include abnormalities in microvascular and macrovascular [2].

In Indonesia, the prevalence of diabetes in adulthood, 20-79 years, was 5.8% of the 9 million total diabetic cases and 4.8 million undiagnosed diabetic cases in 2014 [3]. Oral hypoglycemic drugs, such as biguanide, sulfonylurea, and thiazolidinedionas, are used clinically in the treatment of type 2 diabetes, but it may cause adverse effects and become ineffective in long diabetic treatment. The World Health Organization recommended the use of traditional medicine with basic material of plants for diabetes treatment [4]. Alternative measures using herbal medicine agents may also be considered to achieve lower blood glucose levels, with minimal side effects and low cost [5]. As a rich biodiversity country, Indonesia has 30.000 flowering plants species [6], about 7000 of these species are recognized as medicinal plants [7], 950 species are discovered to have pharmacological properties, 283 species are registered, being cultivated and used by industries [8] and another species directly harvested from forest as raw material [9]. Many plants have potential as an antidiabetic agent that can be proven through ethnobotany information report [10]. The aim of this study is to evaluate and compare the hypoglycemic activity of 10 plants, which are used empirically to treat diabetes in Indonesia. The screened plant parts were mango (*Mangifera indica* L.) leaves [11], avocado (*Persea americana* Mill.) leaves [12], meniran (*Phyllanthus niruri* L.) herb [13], Kelor (*Moringa oleifera* Lam.) leaves [14], ginger (*Zingiber officinale* Roscoe) rhizome [15], jeringau (*Acorus calamus* L.)

rhizome [16], jamblang (*Syzygium cumini* (L.) skeels leaves [17], asam jawa (*Tamarindus indica* L.) seed [18], pare (*Momordica charantia* L.) fruit [17], and Mimba (*Azadirachta indica* A. juss) leaves [19] which were given orally at the same doses (300 mg/kg bw) in mice induced glucose.

METHODS**Plant materials**

The plant materials were collected from several areas, *M. indica* L. leaves, *P. americana* Mill. leaves and *A. calamus* L. rhizome were collected from Lembang, West Java; *P. niruri* L. herb, *S. cumini* (L.) skeels bark, and *Z. officinale* Roscoe rhizome were collected from Subang, West Java; *M. oleifera* Lam. leaves were collected from Karanganyar, Center Java; *T. indica* L. seed were collected from Majalengka, West Java; *M. charantia* L. fruit were collected from Cimahi, West Java; and *A. indica* A. juss leaves were collected from Tawangmangu, Center Java. Those plants were authenticated at Jatinangor Herbarium, Plant Taxonomy Laboratory, Department of Biological Science, UNPAD.

Plant extraction

Plant materials were dried, grounded, extracted (by maceration method in 70% ethanol) and concentrated (using rotary evaporator under reduced pressure [Rotavapor, Buchi]). Extract was dissolved in 2% acacia gum propylene glycol alginate (PGA) to prepare doses and administered by oral gavage.

Animal preparation

The animal house ITB supplied male Swiss Webster mice, weighing 25-35 g. They were housed under standard laboratory condition at a room temperature of 22±3°C and relative humidity of 50-60%, with 12 hrs light and dark cycle. Food and water were provided *ad libitum* according to the laboratory standard. Ethical clearance for experimental animals using was obtained from Ethical Committee of Medical Faculty

UNPAD with Ethical Approval No. 204/UN6.C1.3.2/KEPK/PN/2016 and 206/UN6.C1.3.2/KEPK/PN/2016.

Hypoglycemic activity assay

Oral glucose tolerance test was used to evaluate and compare the hypoglycemic activity of 10 plants ethanolic extract that was tested. Mice were divided into 12 groups (3 mice per group), previously been fasted for 18 hrs. Normal control group was given PGA 2%. Negative control group was given PGA 2% and glucose 1 g/kg bw. Test group was given PGA 2%, glucose 1 g/kg bw and plant extracts tested at an oral dose of 300 mg/kg bw. Blood sampling was performed at 30 minutes after given plant extract (t=0), then at 30 minutes (t=30), 60 minutes (t=60), 90 minutes (t=90), and 120 minutes (t=120) after administration of glucose, determination of glucose level using a glucose meter and strips test (Accu-Chek).

RESULTS AND DISCUSSION

The effect of ethanolic plants extract on blood glucose levels in glucose-induced mice was presented by considering relative blood glucose (%), as shown in Fig. 1 and Table 1. All tested plants extracts might lead the reduction of blood glucose levels at a dose of 300 mg/kg bw, which was related to the duration time of treatment.

As shown in Fig. 1, the negative control group had the highest relative blood glucose levels, indicating that induction method was worked by glucose administration. The relative blood glucose levels decreased in 60 minutes, indicating that the body released insulin hormone to break glucose as a homeostatic process.

Oral administration of 10 medicinal plants within ethanolic extract was performed at a dose of 300 mg/kg, producing various outcomes of blood glucose levels. As shown in Table 1, the highest enhancement of glucose levels occurred at 30 minutes, which was shown in all tested samples *Z. officinale* Roscoe rhizome group was significantly different, compared to negative control, meaning that the extract has hypoglycemic activity better than others at that same period time. While at 60 and 90 minutes, all test groups were significantly different with negative control group.

This showed that the hypoglycemic activity of all tested extracts decreased the blood glucose levels of tested animals. At 120 minutes, all groups had the lowest relative blood glucose levels, compared to all interval times. In particular, *M. indica* L. leaves, *P. americana* Mill. leaves, *P. niruri* L. herb, *Z. officinale* Roscoe rhizome, *A. calamus* L. rhizome, *S. cumini* (L.) skeels bark, *T. indica* L. seed, and *M. charantia* L. fruit group had significantly differences, compared to negative control, indicating

Table 1: Hypoglycemic activity of ethanolic extract of 10 medicinal plants, where the blood sampling was performed at 30 minutes after given plant extract (t=0), at 30 minutes (t=30), 60 minutes (t=60), 90 minutes (t=90), and 120 minutes (t=120) after administration of glucose, determination of glucose level using glucose meter and strips test (Accu-Chek)

Group	Relative blood glucose level (%)				
	Time (minute) mean±SD				
	0	30	60	90	120
Normal control	100	101.11±9.20*	96.99±11.20*	93.87±11.58*	83.18±8.21*
Negative control	100	206.76±7.33	194.79±5.39	177.53±12.76	139.76±40.80
<i>Mangifera indica</i> L. leaves	100	125.53±13.33	104.49±19.85*	89.20±20.39*	83.70±6.50*
<i>Persea americana</i> Mill. leaves	100	164.2±17.81	127.69±9.47*	93.01±3.12*	81.29±7.60*
<i>Phyllanthus niruri</i> L. herb	100	154.28±24.63	109.64±19.25*	94.12±21.60*	74.88±16.16*
<i>Moringa oleifera</i> Lam. leaves	100	184.27±47.70	154.28±31.72*	114.16±34.50*	87.27±28.02
<i>Zingiber officinale</i> Roscoe rhizome	100	120.06±7.88*	109.88±15.83*	93.50±21.16*	83.40±12.69*
<i>Acorus calamus</i> L. rhizome	100	143.96±25.84	99.76±5.30*	88.18±5.30*	80.35±4.23*
<i>Syzygium cumini</i> (L.) skeels bark	100	145.22±41.46	115.83±14.07*	88.13±15.28*	79.41±22.17*
<i>Tamarindus indica</i> L. seed	100	156.40±29.83	103.04±14.44*	95.42±22.39*	65.92±19.29*
<i>Momordica charantia</i> L. fruit	100	144.76±27.02	110.52±30.41*	93.57±16.08*	70.96±13.38*
<i>Azadirachta indica</i> A. juss leaves	100	135.86±12.90	127.47±16.26*	121.75±24.04	89.90±10.08

*Significant compared with negative control (p<0.05)




Fig. 1: Relative blood glucose level (%) of 10 medicinal plants, which are often used as remedies for hyperglycemia in Indonesia. The ethanolic extract was tested for their hypoglycemic activity using glucose tolerance test




Fig. 2: Percentage of decrease blood glucose level (%) of 10 medicinal plants extract tested that induced blood glucose level

that they still had hypoglycemic activity at that time. According to the blood glucose relative level, the percentage of reduction blood glucose level could be calculated with $\frac{Cr(\text{negative control}) - Cr(\text{test group})}{Cr(\text{negative control})} \times 100\%$.

As shown in Table 2 and Fig. 2, all tested extracts decreased blood glucose levels. *Z. officinale* Roscoe rhizome extract had the highest percentage of decrease blood glucose level at 30 minutes. After 60 minutes, the percentage of all tested extracts would increase. In addition, at 90 minutes, all extracts still had hypoglycemic activity, except *M. oleifera* Lam. leaves and *A. indica* A. Juss leaves.

An analysis of variance was conducted to find out differences of hypoglycemic activity each group, followed with Duncan method to test pairwise comparisons between average couples. The result showed that *M. indica* L. has the highest hypoglycemic activity, followed by *Z. officinale* Roscoe, *A. calamus* L., *T. indica* L., *Momordica charantia* L., *S. cumini* (L.) Skeels, *P. niruri* L., *P. americana* Mill., *A. indica* A. Juss, and *M. oleifera* Lam.

M. indica L. leaves have been confirmed to contain mangiferin (1,3,6,7-tetrahydroxyxanthone-C2-beta-D-glucoside, one of xanthone derivatives and C-glucosylxanthones) [20]. Mangiferin was widely distributed especially in the leaves and the bark in the Anacardiaceae and Gentianaceae families [21]. Mangiferin was one of the pharmacologically active compounds, which has been reported to have multiple biological effects and also antidiabetic effects [22-24]. Ethanol extract of the leaves of *M. indica* L. leaves produced a significant hypoglycemic effect at a dose of 250 mg/kg, both in normal and streptozotocin (STZ)-induced diabetic animals. The stimulation of β -cells to release insulin was thought to be part of the mechanism of action [25]. The effect of the aqueous extract of the leaves of *M. indica* L. leaves on blood glucose level in normoglycemic, glucose-induced hyperglycemic, and STZ-induced diabetic rats indicate that the aqueous extract of the leaves of *M. indica* L. possesses hypoglycemic activity. This action may be due to an intestinal reduction of the absorption of glucose [26]. The leaves of *M. indica* L. used for antidiabetic properties using normoglycemic, glucose-induced hyperglycemia, and STZ-induced diabetic mice. The aqueous extract of the leaves of *M. indica* L. possesses hypoglycemic activity [27]. Oral glucose tolerance test in rats mangiferin (10 and 20 mg/kg, i.p.) indicate potent antidiabetic without causing hypoglycemia; mangiferin would then offer a greater therapeutic benefit for the management of DM and diabetic complications associated with abnormalities in lipid profiles [27]. The results from this research suggest that *M. indica* L. leaves can possibly help in the treatment of diabetes.

CONCLUSION

M. indica L. leaves were considered to have the highest hypoglycemic activity in glucose-induced mice, followed by *Z. officinale* Roscoe rhizome, *A. calamus* L. rhizome, *T. indica* L. seed., *M. charantia* L. fruit,

Table 2: Percentage of decreased blood glucose level (%) at 30 minutes (t=30), 60 minutes (t=60), 90 minutes (t=90), and 120 minutes (t=120) after administration of glucose

Group	Time (minute)			
	30	60	90	120
<i>Mangifera indica</i> L. leaves	39.29	46.36	49.75	40.15
<i>Persea Americana</i> Mill. leaves	20.53	34.45	47.61	44.82
<i>Phyllanthus niruri</i> L. herb	25.20	43.72	46.98	46.96
<i>Moringa oleifera</i> Lam. leaves	14.08	20.80	35.70	41.13
<i>Zingiber officinale</i> Roscoe rhizome	41.93	43.59	47.33	38.90
<i>Acorus calamus</i> L. rhizome	30.38	48.78	49.00	42.51
<i>Syzygium cumini</i> (L.) Skeels leaves	29.76	40.54	50.36	43.18
<i>Tamarindus indica</i> L. seed	24.36	47.10	46.25	52.83
<i>Momordica charantia</i> L. fruit	29.99	43.26	47.29	49.23
<i>Azadirachta indica</i> A. Juss leaves	34.29	34.56	31.42	35.68

S. cumini (L.) skeels bark, *P. niruri* L. herb., *P. americana* Mill. leaves, and *A. indica* A. juss leaves.

REFERENCES

- ADA (American Diabetes Association). Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2013;36:S67-74.
- Dipiro JT, Talbert RL, Yee GC, Matzke G, Wells B, Posey LM. *Pharmacotherapy: A Pathophysiologic Approach*. 8th ed. New York: McGraw-Hill; 2011.
- IDF (International Diabetes Federation). Indonesia; 2015. Available from: <https://www.idf.org/membership/wp/indonesia>.
- WHO (World Health Organization). *Prevention of Diabetes Mellitus*. Geneva: WHO; 1994.
- Jouad H, Haloui M, Rhiouani H, El Hilaly J, Eddouks M. Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in the North centre region of Morocco (Fez-Boulemane). *J Ethnopharmacol* 2001;77(2-3):175-82.
- Bappenas. *Strategi Nasional Pengelolaan Keanekaragaman Hayati*. Indonesia: Bappenas; 1993.
- Esai PT. *Indeks Tumbuh-Tumbuhan Obat*. Indonesia: EISAI; 1986.
- Sampoerno H. Pengembangan dan pemanfaatan tumbuhan obat Indonesia. Paper Presented at the National Seminar on Medicinal Plants from Indonesian Tropical Forest. Bogor, Indonesia. 28, April; 1999.
- Zuhud EA, Aziz S, Ghulamahdi M, Andrawulan N, Darusman LK. Dukungan teknologi pengembangan obat asli Indonesia dari segi budaya, pelestarian dan pasca panen. Paper Presented at the Workshop on Agribusiness Development Based on Biopharmaca. Jakarta, Indonesia. Departement Pertanian. 13-15 November; 2001.
- Sinha D, Dwivedi C, Dewangan MK, Yadav R, Rao SP, Chandrakar K, et al. Anti diabetic potential of herbal plants and polyherbal formulation. *Int J Phytother Res* 2014;4 Suppl 3:28-49.
- Petchi RR, Parasuraman S, Vijaya C, Darwhekar G, Devika GS. Antidiabetic effect of kernel seeds extract of *Mangifera indica* (Anacardiaceae). *Int J Pharm Bio Sci* 2011;2 Suppl 1:385-93.

12. Antia BS, Okokon JE, Okon JA. Hypoglycemic activity of aqueous leaf extract of *Persea americana* Mill. Indian J Pharmacol 2005;37 Suppl 5:325-6.
13. Okoli CO, Ibiam AF, Ezike AC, Akah PA, Okoye TC. Evaluation of antidiabetic potentials of *Phyllanthus niruri* in alloxan diabetic rats. Afr J Biotechnol 2009;9(2):248-59.
14. Jaiswal D, Rai PK, Kumar A, Mehta S, Watal G. Effect of *Moringa oleifera* Lam. Leaves aqueous extract therapy. J Ethnopharmacol 2009;123:392-6.
15. Bhandari U, Kanojia R, Pillai KK. Effect of ethanolic extract of *Zingiber officinale* on dyslipidaemia in diabetic rats. J Ethnopharmacol 2005;97:227-30.
16. Prisilla DH, Balamurugan R, Shah R. Antidiabetic activity of methanol extract of *Acorus calamus* in STZ induced diabetic rats. Asian Pac J Trop Biomed 2012;21:941-6.
17. Ajit K, Choudhary BK, Bandyopadhyay NG. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. J Ethnopharmacol 2003;84:105-8.
18. Heidari Z, Sagheb AM, Shahraki M, Moudi B. A stereological study of effects of aqueous extract of *Tamarindus indica* seeds on pancreatic islets in streptozotocin - Induced diabetic rats. Pak J Pharm Sci 2010;23 Suppl 4:427-34.
19. Dholi SK, Raparla R, Mankala SK, Nagappan K. *In vivo* antidiabetic evaluation of neem leaf extract in alloxan induced Rats. J Appl Pharm Sci 2011;1 Suppl 4:100-5.
20. Desai PD, Ganguly AK, Govindachari TR, Joshi BS, Kamat VN, Manmade AH, et al. Chemical investigation of some Indian plants: Part II. Indian J Chem 1996;4:457-549.
21. Yoshimi N, Matsunaga K, Katayama M, Yamada Y, Kuno T, Qiao Z, et al. The inhibitory effects of mangiferin, a naturally occurring glucosylxanthone, in bowel carcinogenesis of male F344 rats. Cancer Lett 2001;163:163-70.
22. Ichiki H, Miura T, Kubo M, Ishihara E, Komatsu Y, Tanigawa K, et al. New antidiabetic compounds, mangiferin and its glucoside. Biol Pharm Bull 1998;21:1389-90.
23. Miura T, Ichiki H, Hashimoto I, Iwamoto N, Kato M, Kubo M, et al. Antidiabetic activity of a xanthone compound, mangiferin. Phytomedicine 2001;8:85-7.
24. Muruganandan S, Srinivasan K, Gupta S, Gupta PK, Lal J. Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. J Ethnopharmacol 2005;97:497-501.
25. Sharma SR, Dwivedi SK, Swarup D. Hypoglycemic potential of *Mangifera indica* leaves in rats. Int J Pharmacol 1997;35:130.
26. Aderibigbe AO, Emudianughe TS, Lawal BA. Antihyperglycaemic effect of *Mangifera indica* in rat. Phytother Res 1999;13:504-7.
27. Aderibigbe AO, Emudianughe TS, Lawal BA. Evaluation of the antidiabetic action of *Mangifera indica* in mice. Phytother Res 2001;15(5):456-8.