

CLINICAL PROFILE AND OUTCOME OF NEONATES ADMITTED TO A SECONDARY-LEVEL NEONATAL INTENSIVE CARE UNIT IN NORTH INDIA

SHAGUFTA YOUSUF¹, SHOWKAT HUSSAIN TALI^{2*}, IFTIKHAR HUSSAIN³

¹Department of Obstetrics and Gynaecology, Adesh Institute of Medical Sciences, Bathinda, Punjab, India. ²Department of Pediatrics, Adesh Institute of Medical Sciences, Bathinda, Punjab, India. ³Department of Social Preventive Medicine, Government Medical College Srinagar, Jammu and Kashmir, India. Email: drshowkatshifa@gmail.com.

Received: 09 May 2017, Revised and Accepted: 06 June 2017

ABSTRACT

Objectives: The objective of this study was to evaluate the performance of a secondary-level neonatal intensive care unit (NICU).

Methods: A total of 336 neonates were enrolled in the study. Their clinical profiles at admission and final outcomes were recorded in a predesigned pro forma.

Results: Hyperbilirubinemia, sepsis, and perinatal asphyxia were the most common reasons for admission. Eighty-five percentage of the neonates could be managed in secondary-level newborn unit with comparable mortality and mortality to a tertiary-level newborn care unit.

Conclusion: Strengthening of secondary newborn care units is a viable option that will help to decrease the burden of tertiary-level NICUs.

Keywords: Neonate, Clinical profile, Outcome, Secondary-level neonatal intensive care unit.

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2017.v10i9.19799>

INTRODUCTION

As of 2015, in India, the neonatal mortality rate is 28/1000 live births [1]. The 2030 Agenda for Sustainable development (WHO) calls for reduction in neonatal mortality to 12/1000 live births by 2030 [2]. Most term neonates can be managed at home under the guidance and supervision of mother or health-care professional. On the other hand, low birth weight and preterm neonates are fragile and vulnerable and they demand high level of skills and technology in a special care nursery or neonatal intensive care unit (NICU) for their intact survival. Apart from high mortality, many avoidable disabilities such as cerebral palsy, mental subnormality, learning disabilities, and recurrent seizures have their origin in perinatal period [3,4]. It is encouraging to note that neonates with a birth weight of 1 kg had a mortality of 95% in 1960 and now have 95% chances of survival. In India where a large number of deliveries are non-institutional, secondary- and community-level hospitals form important pillars of health-care delivery. The data from secondary-level newborn care units are sparse, and most of the data on neonatal mortality and morbidity are from tertiary-level care centers [5-7]. The purpose of this study was to evaluate the performance of a secondary-level hospital and stress the importance of neonatal care at this level.

METHODS

This study was carried out at LD Hospital which is an associated hospital of Government Medical College Srinagar and the main maternity care hospital in the state of Jammu and Kashmir. The hospital has a neonatal care unit attached to it that can be compared to a secondary-level NICU. At the time of study, it was a 25-bedded nursery cum NICU with 15 radiant warmers, 3 phototherapy units, 15 infant trolleys, 4 ventilators, and central oxygen supply. However, these ventilators were less often used due to lack of adequately trained personnel present round the clock. One consultant (pediatrician), one senior resident (MD/DCH Pediatrics), 4 junior residents (MBBS), 4 nurses, and two nursing orderlies would be present during the day duty. One postgraduate student, 2 junior residents, 3 nurses, and 2 nursing orderlies would

be present during the night shifts. A consult would be on call round the clock. There were round-the-clock facilities for emergency X-ray and basic laboratory investigations. The study was commenced after obtaining clearance from hospital's ethical committee and written informed consent from one of the parents or the guardian.

Guidelines followed in the NICU

1. Only inborn neonates were admitted as per hospital policy.
2. Preterm neonates <32 weeks, birth weight <1200 g, preterm with severe respiratory distress syndrome, stage 2 and 3 hypoxic-ischemic encephalopathy, cyanotic congenital heart diseases, and other serious congenital malformations were referred to higher centers whenever possible.
3. Neonates between 1200 g and 1500 g and stable were managed as per standardized protocol. Unstable neonates were referred to higher centers.

Table 1: Demography and other baseline characteristics

Attribute	n (%)
Sex	
Male	200 (59.5)
Female	136 (40.5)
Residence	
Rural	202 (60.1)
Urban	92 (39.9)
Place of delivery	
Home delivery	0
Hospital delivery	336 (100)
Birth weight	
AGA	220 (65.5)
SGA	116 (35.5)
Gestational age (weeks)	
<32	8 (2.4)
32-34	88 (26.2)
>34	240 (71.4)

AGA: Appropriate for gestational age, SGA: Small for gestational age

Table 2: Clinical profile and outcome of neonates (n=336)

Attribute	n (%)		
	Survived	Expired	Transferred to higher center
Preterm (<37 weeks) (n=96)			
Preterm with hyaline membrane disease	2 (0.6)	10 (3)	24 (7.1)
Sepsis	14 (4.1)	10 (3)	
Other morbidities	32 (9.5)	4 (1.2)	
Term (>37 weeks) (n=240)			
Perinatal asphyxia	50 (14.8)	11 (3.3)	24 (7.1)
Sepsis	32 (9.5)	7 (2)	
Pneumonia	12 (3.5)	1 (0.3)	
Hyperbilirubinemia	78 (23.2)	-	
Congenital heart disease	2 (0.6)	1 (0.3)	
Hypoglycemia	6 (2.8)	-	
Other morbidities	16 (4.7)	-	
Total (n=336)	244 (72.6)	44 (13.1)	48 (14.3)
Geography (n=336)			
Urban	77 (22.9)	15 (4.5)	13 (3.9)
Rural	167 (49.7)	29 (8.6)	35 (10.4)

4. Stable neonates more than 1500 g and 34 weeks were generally kept with mother if there was no associated morbidity or high risk for neonatal sepsis. They were started on breast feeds or expressed breast milk by katori/spoon under medical supervision and were monitored for feeding difficulty, respiratory distress, jaundice, and cyanosis for 48-72 hrs. Neonates who would behave abnormally would be shifted to the NICU.

RESULTS

Demographic and other baseline characteristics are shown in Table 1. Clinical profile and outcome of neonates are shown in Table 2.

DISCUSSION

A total of 336 neonates were admitted during the study period. Ninety-six (28.6%) neonates were early preterm (<34 weeks) and 240 (71.4%) were late preterm and term. Out of these, 244 (72.6%) survived, 44 (13.1%) died, and 48 (14.3%) were transferred to higher centers. As many as 35.5% of neonates were small for gestational age. Hyperbilirubinemia (23.2%), perinatal asphyxia (14.8), and sepsis (9.5) were the major reasons for admission. Shah *et al.* [8] reported in their study that the share of preterm in neonatal admission was as high as 24 (Table 1). Modi and Kirubakaran [9] reported suspected sepsis (23.7%), low Apgar score (5.3%), preterm delivery (14.2%), and jaundice (13.4%) as the major reasons for admission.

Our study highlights that as many as 85% of neonates can be managed at secondary level with a survival of 72.6% (Table 2). This is in conformity with the study conducted by Garg *et al.* [10]. In their study, the overall survival rate was 65%. Forty-two (25%) neonates expired and 5.4% were referred to higher centers. Shah *et al.* [8] reported an overall mortality of 20.2% during hospital stay. Bose *et al.* [11] also reported similar findings. Of the 175 neonates, 6 expired, 2 were discharged against medical advice, and 8 were transferred to the nursery in the tertiary hospital.

There was an appreciable difference in urban and rural mortality rates (Table 1). Mortality in rural population was 2 times than that of urban population. This may be due to poor antenatal care of rural women. This is in conformity with the National Family Health Survey 3 [12].

Mortality and morbidity profile was similar to those reported from tertiary care level centers. Modi and Kirubakaran [9] observed that all the neonates admitted to tertiary care level neonatal unit at CMC Vellore, there were 87 early neonatal deaths, 4% (49) of inborn admissions and 18% (38) of out born admissions. Basnet and Shrestha [13] in their study at a tertiary care level hospital also observed neonatal mortality

ranges from 9.46 to 14.88 per 1000 live births per year. Ahmed *et al.* [14] observed a mortality of 17%.

CONCLUSION

Strengthening of secondary level newborn care units is a viable option that will help to decrease the burden of tertiary level NICUs.

REFERENCES

- Available from: <http://www.unicef.org/infobycountry/indiastatistics.html>. [Last accessed on 2017 Apr 04].
- United Nations Transforming our world: The 2030 Agenda for Sustainable Development. New York, NY, USA: United Nations; 2015. Available from: <http://www.tinyurl.com/od9mens>. [Last accessed on 2017 Apr 04].
- Singh M. Care of New the born. 7th ed. New Delhi: CBS Publishers & Distributors; 2010.
- Choure MK, Jadhav RR, Padwal SL. Drug utilization study in neonatal intensive care unit at rural tertiary care hospital. *Asian J Pharm Clin Res* 2017;10(4):102-4.
- National Neonatology Forum. Report of the National-Perinatal Database 2000, New Delhi; 2001.
- Singh M, Deorari AK, Khajuria RC, Paul VK. A four year study on neonatal morbidity in a New Delhi hospital. *Indian J Med Res* 1991;94:186-92.
- Chavan YS, Dattal MS, Khadilkar VV. Causes of early neonatal mortality. *Indian Pediatr* 1992;29:781-3.
- Shah GS, Yadav S, Thapa A, Shah L. Clinical profile and outcome of neonates admitted to neonatal intensive care unit (NICU) at a tertiary care centre in Eastern Nepal. *Paediatr Soc* 2013;33(3). DOI: 10.3126/jnps.v33i3.8447.
- Modi N, Kirubakaran C. Reasons for admission, causes of death and costs of admission to a tertiary referral neonatal unit in India. *J Trop Pediatr* 1995;41(2):99-102.
- Garg P, Krishak R, Shukla DK. NICU in a community level hospital. *Indian J Pediatr* 2005;72(1):27-30.
- Bose A, Sinha S, Choudhary N, Aruldas K, Moses PD, Joseph A. Experiences of neonatal care in a secondary level hospital. *Indian Pediatr* 1999;36(8):802-6.
- National Family Health Survey 3 (NFHS 3) 2005-2006. International Institute for Population Sciences, Mumbai; India, ORC Macro, Maryland. Available from: <http://www.rchiips.org/ARCH-3.htm>. [Last accessed on 2017 May 06].
- Basnet S, Shrestha L. Trends in neonatal mortality at a tertiary level teaching hospital. *J Nepal Pediatr Soc* 2013;33(3). DOI: 10.3126/jnps.v33i3.8957.
- Ahmed A, Lutfi S, Al-Hail M, Al-Saadi M. Antibiotic susceptibility patterns of microbial isolates from blood culture in the neonatal intensive care unit of Hamad medical corporation (HMC), Doha, Qatar. *Asian J Pharm Clin Res* 2013;6(2):191-5.