

EVALUATION OF ANTIDEPRESSANT ACTIVITY OF ETHANOLIC EXTRACT OF *LAGERSTROEMIA SPECIOSA* LEAVES IN ALBINO WISTAR RATS

VANITA KANASE, SUNITA VISHWAKARMA*

Department of Pharmacology, Oriental College of Pharmacy, Navi Mumbai, Maharashtra, India. Email: sunitapharmacy1@gmail.com

Received: 22 September 2019, Revised and Accepted: 12 December 2019

ABSTRACT

Objective: The objective of the study was to evaluate the antidepressant activity of ethanolic extract of dried leaves of *Lagerstroemia speciosa* L. (EELS) on acute restraint stress (ARS)-induced depression-like behavior and biochemical alterations in albino wistar rats.

Methods: Thirty rats were randomly divided into five experimental groups. Group-I (normal control) rats received normal saline (2.0 ml/kg, p.o.) daily for 14 days; Group-II (stress control) rats received normal saline (2.0 ml/kg, p.o.) daily for 14 days and subjected to restraint stress on the 13th day. Group-III (standard drug-treated) rats received imipramine (15 mg/kg, p.o.) daily for 14 days and subjected to restraint stress on the 13th day. Groups-IV and V rats were treated with EELS (100 mg/kg and 300 mg/kg, p.o.) daily for 14 days subjected to ARS on the 13th day. Stress-like behavior was assessed by subjecting the rats to behavioral paradigms such as tail-suspension test (TST) and open field test (OFT), 40 min post-restraint stress procedure. Pretest of 10 min for forced swim test (FST) was also given to each rat simultaneously. Then, 23.5 h later, the relevant samples were administered and the main test performed 30 min later. Oxidative stress parameters such as superoxide dismutase (SOD), catalase (CAT), malondialdehyde (MDA), and extent of lipid peroxidation (LPO) were analyzed in restraint stress-induced animals and control group, following FST on the 15th day.

Statistical Analysis: Expression of data was done as a mean standard error of the mean. The normally distributed data were subjected to one-way analysis of variance followed by Dunnett's test. *p<0.05 was considered statistically significant.

Results: It was observed that *L. speciosa* L. showed a significant dose-dependent decrease in duration of immobility time in TST and FST when compared with the control group in a dose-dependent manner. The results of OFT also showed a dose-dependent increase in locomotor activity. In addition to behavioral tests, EELS also normalized oxidative stress markers such as CAT, SOD, MDA, and LPO in a dose-dependent manner.

Conclusion: The results suggest that the ethanolic extract of *L. speciosa* L. leaves possesses significant antidepressant property, may be recommended as a supplement for the antidepressant activity.

Keywords: *Lagerstroemia speciosa*, Antidepressant, Acute restraint stress, Tail-suspension test, Open field test, Forced swimming test, Imipramine, Biochemical estimation.

© 2020 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2020.v13i2.35823>

INTRODUCTION

Depression is a heterogeneous disorder that affects a person's mood, physical health, and behavior. It is caused not only by changing lifestyle as perceived by the general public but also by some of the allopathic drugs, for example, an anti-hypertensive drug, reserpine that depletes neuronal storage granules of norepinephrine (NE), serotonin, and dopamine (DA), cause clinically significant depression in more than 15% of patients. Patients with major depression have symptoms that reflect changes in brain monoamine neurotransmitters, specifically NE, serotonin, and DA. According to a World Health Report, about 450 million people suffer from a mental or behavioral disorder, yet only a small proportion of them receive even the basic treatment [1]. Depression accounts for about 12% of the global burden of disease which is expected to rise to 15% by 2020. The major problems of existing allopathic antidepressant drugs include delayed clinical benefit, serious side effects, and a response rate of <50%. Commonly used drugs for depression are monoamine oxidase (MAO) inhibitors and tricyclic antidepressants (TCAs). They increase the synaptic concentration of at least two of three neurotransmitters, namely, 5-hydroxytryptamine or serotonin (5-HT), NE, and DA. The combined effect of serotonin selective reuptake inhibitor (SSRI) and serotonin reuptake transporter inhibitor increases the synaptic concentration of 5-HT and its duration of action. Ethanolic extracts of most of the herbal drugs are rich in chemical constituents such as phenolic compounds, flavonoids, triterpenoids, and steroids. Ethanolic extracts have less

harmful effects, availability, and lower cost of medicinal plants versus synthetic substances make them as outstanding and simple selection in the treatment of nervous diseases. Therefore, identification and validation of plant-derived substances for the treatment of various depressive disorders attract the attention of researchers [2].

Lagerstroemia speciosa L. is also known as the crepe myrtle or Pride of India or Banaba belongs to the family (Lythraceae). *Lagerstroemia* is named for the Swedish merchant, Magnus von Lagerstrom, was honored by Linnaeus, *speciosa* means showy, referring to the flowers varying in color from pink to purple. The tree is native to Southeast Asia and is called Banaba in the Philippines [3]. The flowers and leaves are used to make an herbal tea, as they contain corosolic acid, a chemical that has an insulin-like effect of lowering glucose levels in the body. It is also utilized in weight loss management [4].

METHODS**Collection and authentication of plant**

The twigs (leaves, flowers, and fruits) of *L. speciosa* L. were collected from Mahapalika Garden, Bhayandar East, Thane. Sample specimen voucher was submitted to Dr. Rajendra D. Shinde, Director, Blatter Herbarium of St. Xavier's College, Mumbai. The leaves were compared and authenticated by comparison with the Blatter Herbarium Specimen No. JF-1532 of J. Fernandez.

The leaves were washed with tap water and shade dried at normal room temperature with the aid of circulating airflow using a fan, and dried leaves were ground to make a coarse powder.

Preparation of extract

The powdered plant leaves of *L. speciosa* L. were extracted with ethanol in the Soxhlet apparatus. The content of the round bottom flask was emptied in the Petri plate, and the solvent was allowed to evaporate. The extract was evaporated to obtain the dry powder of extract. This crude dry extract was stored in a suitable container and kept in the refrigerator (0–4)°C until use.

Physicochemical analysis of powdered leaves

Physicochemical parameters were carried out on powdered leaves of *L. speciosa* L. for loss on drying, total ash value, acid-insoluble ash value, water-soluble ash value, and determination of extractive values and were investigated as described by the well-established methods [5,6].

Preliminary phytochemical screening of the extract

Preliminary chemical tests were carried out on the ethanolic extract of *L. speciosa* L. for the determination of the presence of phytoconstituents such as alkaloids, flavonoids, cardiac glycosides, saponins, steroids, terpenoids, tannins, and phenolic compounds and were investigated as described by the well-established methods [5,6].

Experimental animals

Female albino wistar rats (200–300 g) were used for the study. The animals were obtained from SA-FORD, Plot No.: V-10, MIDC, Taloja, Navi Mumbai, Dist. Raigad Maharashtra - 410 208. The use of these animals and the study protocols were approved by CPCSEA recognized the Institutional Animal Ethical Committee (IAEC) of Oriental College of Pharmacy, Sanpada, Navi Mumbai - 400 705 under protocol no. OCP/IAEC/2018-19/08. Rats were kept at the animal house in polypropylene cages (three in each cage), at 22±2°C, with 12:12 h dark: light cycle. They were provided with commercial rat feed and water given *ad libitum*. The animals were allowed to acclimatize for 7 days before the study.

Selection of doses

In the literature survey, it was found that the ethanolic extract of *L. speciosa* L. was safe. LD₅₀ of the ethanolic extract is reported to be 3000 mg/kg [7]. The plant is often eaten by animals, which is also an indicator to prove it is less toxic. Thus, for the purpose of the research study, the doses of ethanolic extract of *Lagerstroemia speciosa* (EELS) were finalized 100 mg/kg and 300 mg/kg.

Drugs and chemicals

Imipramine hydrochloride (Abbott, Mumbai, India) was used as reference standards for the antidepressant activity.

Experimental design

Thirty female albino wistar rats were randomly divided into five experimental groups containing six rats each as shown in Table 1. Group-I (normal control) rats received normal saline (2.0 ml/kg, p.o.) daily for 14 days; Group-II (stress control) rats received normal saline (2.0 ml/kg, p.o.) daily for 14 days and subjected to restraint stress on the 13th day. Group-III (standard drug-treated) rats received imipramine

(15 mg/kg, p.o.) daily for 14 days and subjected to restraint stress on the 13th day. Groups-IV and V rats were treated with EELS (100 mg/kg and 300 mg/kg, p.o.) daily for 14 days subjected to acute restraint stress (ARS) on the 13th day.

Depression-like behavior was assessed by subjecting the rats to behavioral paradigms such as tail-suspension test (TST) and open field test (OFT), 40 min post-restraint stress procedure. Pretest of 10 min for forced swim test (FST) was also given to each rat simultaneously. Then, 23.5 h later, the relevant samples were administered and the main test performed 30 min later.

Oxidative stress parameters such as superoxide dismutase (SOD), catalase (CAT), malondialdehyde (MDA), and extent of lipid peroxidation (LPO) were analyzed in restraint stress-induced animals and control group, following FST on the 15th day.

Procedure for ARS

ARS was accomplished by placing rats in an individual plastic rodent restraint device for 12 h. This restrained all physical movements without subjecting the animal to pain. Animals were deprived of food and water during the entire period of exposure to stress. After 12 h, the animals were released from their enclosure and 40 min post-release, the animals were subjected to behavioral tests and then to biochemical estimations. In the normal control group, the rats were kept in the animal cage in the experimental room [8,9].

Behavioral tests

TST

The animals were suspended individually by the end of the tail with micropore adhesive tape (approximately 1 cm) with the head 50 cm from the bottom in a suspension box 40 min post-restraint stress procedure. Rats were suspended for a total of 6 min. During the final 4 min interval of the test, the duration of immobility was recorded. Rats were considered immobile only when they will be hung passively and absolutely motionless. Antidepressant decreases the immobility of the rat in this test [10].

OFT

Open-field apparatus was made as reported, each rat was placed in the center of the open field, and its behavior observed for 5 min. The parameters evaluated were the total number of squares crossed, the number of outer squares (those adjacent to the walls) crossed, and the number of inner squares crossed; the three measures referred to as total (TL), peripheral (PL), and central locomotion, respectively. The numbers of leanings (one or two paws in contact with the wall), rearing (the mouse standing on its two hind paws without touching the walls), grooming (face cleaning, paw licking, fur licking, head scraping, and rubbing), and defecations were also recorded. At the end of each test, the whole area was cleaned with a wet sponge and a dry paper towel [11].

FST

On day 14, all the rats were allowed to swim individually for 10 min for adaptation. Then, 23.5 h later, the relevant samples were administered and the main test performed 30 min later, i.e., on day 15. Rats were

Table 1: Grouping of animals

S. No.	Groups	Test substances	Animal required per group (female albino wistar rats)	Dose	Total number
1.	Normal control	Normal saline	6	2 ml/kg	6
2.	Stress control	Normal saline	6	2 ml/kg	6
3.	Standard control	Imipramine	6	15 mg/kg	6
4.	Test group 1	EELS	6	100 mg/kg	6
5.	Test group 2	EELS	6	300 mg/kg	6
Total animals required					30

EELS: Ethanolic extract of *Lagerstroemia speciosa*

forced to swim in a cylinder (diameter 40 cm, height 60 cm) containing 30 cm of freshwater maintained at 25°C±1°C. Water in the cylinder was changed after each animal to prevent behavioral alteration among animals due to used water. Each animal showed vigorous movement during the initial 2 min period of the test. Duration of immobility will be manually recorded during the next 4 min of a total 6 min testing period by the observer. Rats were considered to be immobile when they floated in an upright position, making only small movements to keep their heads above the water level. Following the swimming session, rats were dried using a cotton towel and returned to home cages after the experiment. A decrease in the duration of immobility is indicative of antidepressant-like effect, whereas an increase of immobility time, when compared with the control group, is associated with depressive-like effects [12,13].

Biochemical estimation

Preparation of brain tissue homogenate

All the animals were sacrificed by euthanasia using the CO₂ chamber as per rodent euthanasia guidelines, after behavioral observations. The brains were quickly removed, washed in ice-cold sterile isotonic saline, and weighed. A 10% (w/v) tissue homogenates were prepared with 0.1 M phosphate buffer (pH 7.4). The supernatant was obtained by centrifugation of the homogenate at 1000 rpm for 20 min at 5°C and used for further biochemical estimation [14].

CAT activity

The supernatant (50 µl) was added to a cuvette containing 2.95 ml of 19 mM/L solution of H₂O₂ prepared in potassium phosphate buffer. The change in absorbance was monitored at 240 nm wavelength at the 1 min interval for 3 min. The presence of CAT decomposes H₂O₂ leading to a decrease in absorbance [15].

Sodium oxide dismutase activity

The SOD activity in the supernatant was measured by the method of Misra and Fridovich. The supernatant (500 µl) was added to 0.800 ml of carbonate buffer (100 mM, pH 10.2) and 100 µl of epinephrine

(3 mM). The change in absorbance of each sample was then recorded at 480 nm in spectrophotometer for 2 min at an interval of 15 s. Parallel blank and standard were run for determination of SOD activity. The reaction mixtures are diluted 1/10 just before taking the readings in a spectrophotometer [16].

Determination of MDA formation

To 1 ml of suspension medium was taken from the 10% tissue homogenate. To 0.5 ml of 30% TCA will be added to it, followed by 0.5 ml of 0.8% thiobarbituric acid (TBA) reagent. The tubes were then be covered with aluminum foil and kept in shaking water bath for 30 min at 80°C. After 30 min, tubes were taken out and kept in ice-cold water for 30 min. These were then be centrifuged at 3000 rpm for 15 min.

The absorbance of the supernatant was read at 540 nm at room temperature against an appropriate blank. Blank consists of 1 ml distilled water, 0.5 ml of 30% TCA and 0.5 ml of 0.8% TBA [17].

Determination of LPO assay

To 0.2 ml of the test sample, 0.2 ml of SDS, 1.5 ml of acetic acid, and 1.5 ml of TBA were added. The mixture was made up to 4 ml with water and then heated in a water bath at 95°C for 60 min. After cooling, 1 ml of water and 5 ml of n-butanol/pyridine mixture were added and agitated smartly. After centrifugation at 4000 rpm for 10 min, the organic layer was taken and its absorbance was scan at 532 nm. The level of lipid peroxides was expressed as nmoles of MDA released/g wet tissue [17].

Statistical analysis

The data obtained from animal experiments were analyzed with InStat Software by GraphPad (version 3.10). It was expressed as mean±standard error of the mean. For statistical analysis, the data were subjected to analysis of variance followed by Dunnett's t-test. Results were considered to be statistically significant at p≤0.05. Significance levels were as follows:

*Indicates p≤0.05 as significant; **indicates p≤0.01 as highly significant; ***indicated p≤0.001 as very significant.

RESULTS

Physicochemical analysis of powdered leaves

The result of Physicochemical analysis of powdered leaves of *Lagerstroemia speciosa* are shown in Table 2.

Qualitative phytochemical screening

The result of quantitative phytochemical analysis of powdered leaves of *Lagerstroemia speciosa* are shown in Table 3.

Antidepressive evaluation

TST

Both the doses of the ethanolic extract of leaves of *L. speciosa* L. showed a dose-dependent decrease in immobility time when compared against

Table 2: Result of physicochemical analysis of powdered leaves of *Lagerstroemia speciosa*

Sr. No.	Test	Result (%)
1.	Loss on drying	60.67
2.	Total ash value	16.39
3.	Acid-insoluble ash value	6.24
4.	Water-soluble ash value	7.58
5.	Petroleum ether soluble extractive value	9.6
6.	Chloroform soluble extractive value	6.4
7.	Ethyl acetate soluble extractive value	2.4
8.	Ethanol soluble extractive value	8.8
9.	Water-soluble extractive value	15.2

Table 3: Result of quantitative phytochemical analysis of powdered leaves of *Lagerstroemia speciosa*

S. No.	Phytoconstituents	Test	Ethanolic extract of the leaves <i>Lagerstroemia speciosa</i> L.
1.	Carbohydrates	Molisch's test	+ve
		Fehling's test	+ve
2.	Proteins and free amino acids	Ninhydrin test	-ve
3.	Alkaloids	Mayer's test	-ve
4.	Cardiac glycosides	Keller-Kiliani test	+ve
5.	Steroids	Liebermann-Burchard test	+ve
6.	Terpenoids	Salkowski test	+ve
7.	Tannins	5% FeCl ₃ solution	+ve
8.	Flavonoids	Shinoda test	+ve
9.	Phenols	Lead acetate test	+ve
10.	Saponins	Foam test	-ve

+ve: Present, -ve: Absent

stress control as well as against imipramine which was used as a standard. The results of TST are shown in Table 4 and Fig. 1.

OFT

Both the doses of the ethanolic extract of leaves of *L. speciosa* L. showed a dose-dependent increase in locomotor activity when compared against stress control as well as against imipramine which was used as a standard. The results of TST are shown in Table 5 and Fig. 2.

FST

Both the doses of the ethanolic extract of leaves of *L. speciosa* L. showed a dose-dependent decrease in immobility time when compared against stress control as well as against imipramine which was used as a standard. The results of TST are shown in Table 6 and Fig. 3.

Biochemical estimations

CAT activity

Evaluation of CAT activity revealed that stressed rats presented a significant decrease in CAT activity, which was significantly prevented by EELS (100 mg/kg and 300 mg/kg) pre-treatment when compared to unstressed group, as shown in Table 7 and Fig. 4.

SOD activity

In the rats pretreated with EELS 100 mg/kg and 300 mg/kg p.o., the level of SOD was significantly increased (p<0.001) as compared to ARS rats. Table 7 and Fig. 5 show significant and dose-dependent recovery on ARS induced reduced level of SOD in the animal due to EELS.

MDA formation

The results depicted in Table 7 and Fig. 6 illustrate that ARS significantly increased MDA level in rat brain as compared to unstressed rats. The results indicated that EELS (100 mg/kg and 300 mg/kg) pre-treatment and imipramine significantly abolished the increase in MDA level caused by ARS.

LPO assay

Quantitative measurement of LPO in the whole brain was assessed based on the amount of MDA formed, the statistical analysis revealed that ARS produced a significant increase in MDA level whereas EELS (100 mg/kg and 300 mg/kg) pre-treatment significantly abolished the increase in MDA level compared to stressed animals. The results are shown in Table 7 and Fig. 7.




Fig. 1: Effect of the ethanolic extract of *Lagerstroemia speciosa* L. leaves on immobility time of tail suspension test in albino wistar rats. a Versus normal control stress group and b versus acute restraint stress group




Fig. 2: Effect of the ethanolic extract of *Lagerstroemia speciosa* L. leaves on locomotor activity of open field test in albino wistar rats. a Versus normal control stress group and b versus acute restraint stress group

Table 4: Effect of ethanolic extract of *Lagerstroemia speciosa* L. leaves on immobility time of tail suspension test in albino wistar rats

S. No.	Treatment	Duration of immobility (s)
1.	Normal control	112.17±12.60
2.	ARS	145.17±12.46
3.	Imipramine (15 mg/kg p.o.)+ARS	79.16±4.67**
4.	EELS (100 mg/kg p.o.)+ARS	82.83±6.90**
5.	EELS (300 mg/kg p.o.)+ARS	70.16±5.18**

Values are the mean±standard error of the mean of n=6 rats/treatment. Significance **p≤0.01. ARS: Acute restraint stress, EELS: Ethanolic extract of *Lagerstroemia speciosa*

Table 5: Effect of the ethanolic extract of *Lagerstroemia speciosa* L. leaves on locomotor activity of open field test in albino wistar rats

S. No.	Treatment	Total	Peripheral	Central	Leaning	Rearing	Grooming	Defecation
1.	Normal control	50.33±3.38	46.00±3.51	4.33±1.49	9.00±2.59	0.83±0.4	3.17±0.60	2.83±1.4
2.	ARS	28.50±2.49**	26.17±2.53	2.17±0.17	11.5±3.4	1.5±1.15	3.00±0.68	0.83±0.65
3.	Imipramine (15 mg/kg)+ARS	49.83±6.09**	43.67±6.70	6.50±2.90	10.17±2.21	3.17±2.21	4.67±1.09	0.67±0.33
4.	EELS (100 mg/kg)+ARS	53.83±4.85*	48.50±5.31	2.33±0.76	8.00±2.48	2.83±0.31	3.50±0.76	3.00±1.03
5.	EELS (300 mg/kg)+ARS	58.33±4.40**	52.50±4.56	3.00±1.03	5.17±1.76	1.17±0.54	1.50±0.50	2.17±0.98

Values are the mean±standard error of the mean of n=6 rats/treatment. Significance **p≤0.01, *p≤0.05. EELS: Ethanolic extract of *Lagerstroemia speciosa*, ARS: Acute restraint stress

DISCUSSION

The incidence of depression in the community is very high and is associated with a lot of diseases. Hence, it is very important to report these problems and find effective remedies. Several drugs are available, all are associated with some limitations and there is a crucial need for alternative medications for these disorders. Medical therapies with herbs may be effective alternatives in the treatment of depression, and the research of their effects has progressed significantly since the past decade [18,19].

Table 6: Effect of the ethanolic extract of *Lagerstroemia speciosa* L. leaves on immobility time of force swim test in albino wistar rats

S. No.	Treatment	Duration of immobility (s)
1.	Normal control	42.50±2.63
2.	ARS	72.83±2.63**
3.	Imipramine (15 mg/kg p.o.)+ARS	49.83±4.38**
4.	EELS (100 mg/kg p.o.)+ARS	49.67±2.70**
5.	EELS (300 mg/kg p.o.)+ARS	30.83±4.03**

Values are the mean±standard error of the mean of n=6 rats/treatment. Significance **p≤0.01. EELS: Ethanolic extract of *Lagerstroemia speciosa*, ARS: Acute restraint stress




Fig. 3: Effect of the ethanolic extract of *Lagerstroemia speciosa* L. leaves on immobility time of force swim test in albino wistar rats. a Versus normal control group and b versus acute restraint stress group




Fig. 4: Effect of EELS pre-treatment on ARS induced changes on catalase activity. NC: Normal control, ARS: Acute restraint stress, EELS: Ethanolic extract of *Lagerstroemia speciosa*. Values are expressed as mean±standard error of the mean (n=6). **p≤0.01 and *p≤0.05, a versus NC group and b versus ARS group

The introduction of drugs such as amitriptyline, fluvoxamine, imipramine, citalopram, venlafaxine, and others has modernized the treatment of depression. The marvellous efficacy of imipramine in these depressive disorders has paved the way for the introduction and use of newer antidepressant agents. However, the safety factor in respect of the imipramine antidepressant drugs has been rather intriguing and hence a definite need is visualized for the introduction of safer antidepressant drugs having no troublesome adverse effects [20].

The ARS was used as a model of depression induced by stress, which has both emotional and physical components in combination to distress the brain's intracellular redox status. The main purpose of using restraint is that it induces an inescapable physical and mental stress which is not accompanied by a conditioned response. It is important to clarify that this study was performed in female rats because women are more susceptible to the development of depressive disorder followed to lifetime stress events than men [21].

The behavior despair test (TST and FST) is quite sensitive and relatively specific to all major classes of antidepressant drugs, including tricyclics, SSRI, MAO inhibitors, and atypicals. Imipramine is a pre-synaptic uptake inhibitor of both nor-adrenaline and serotonin. Since catecholamines and 5-HT have been implicated in the etiology of




Fig. 5: Effect of EELS pre-treatment on ARS induced changes on SOD activity. NC: Normal control, ARS: Acute restraint stress, EELS: Ethanolic extract of *Lagerstroemia speciosa*. Values are expressed as mean±standard error of the mean (n=6). **p≤0.01 and *p≤0.05, a versus NC group and b versus ARS group




Fig. 6: Effect of EELS pre-treatment on ARS induced changes on MDA activity. NC: Normal control, ARS: Acute restraint stress, EELS: Ethanolic extract of *Lagerstroemia speciosa*. Values are expressed as mean±standard error of the mean (n=6). ** p≤0.01, a versus NC group and b versus ARS group

Table 7: Effect of oxidative stress markers in brain

S. No.	Treatment	Catalase	Superoxide dismutase	Malondialdehyde	Lipid peroxidation
1.	Normal control	16.81±1.082	1.12±1.082	0.21±0.030	0.107±0.004
2.	ARS	8.41±1.502	0.57±0.079	0.45±0.059	0.355±0.050
3.	Imipramine (15 mg/kg)+ARS	14.86±1.019**	1.15±0.075**	0.20±0.048**	0.142±0.015**
4.	EELS (100 mg/kg)+ARS	13.4±1.226**	1.23±0.093**	0.18±0.026**	0.133±0.010**
5.	EELS (300 mg/kg)+ARS	15.85±1.485**	1.34±0.095**	0.16±0.025**	0.123±0.012**

Values are the mean±standard error of the mean of n=6 rats/treatment. Significance **p<0.01. a versus NC group and b versus ARS group. EELS: Ethanolic extract of *Lagerstroemia speciosa*, ARS: Acute restraint stress




Fig. 7: Effect of EELS pre-treatment on ARS induced changes on LPO activity. NC: Normal control, ARS: Acute restraint stress, EELS: Ethanolic extract of *Lagerstroemia speciosa*. Values are expressed as mean±standard error of the mean (n=6). ** p<0.01, a versus NC group and b versus ARS group

depression, the positive effect of these drugs in TST and FST seems to be due to increased availability of these neurotransmitters at the post-synaptic receptor sites following their reuptake inhibition. In the current study, ARS significantly increased the immobility time in the FST and TST indicating depressive-like behavior which showed that restraint stress-induced depressive-like behavior as evidenced by increased immobility time in FST and TST. This immobility reflects a state of despair in animals and is claimed to reproduce a condition similar to depression in humans. Animals after antidepressant treatment feel amazing even in desperate situations, and they spend less time with immobility. In the present study, EELS in the dose of 100 and 300 mg/kg showed a significant dose-dependent antidepressant-like effect in behavior despair test (TST and FST), as they reduced the immobility time when compared against stress control as well as against standard imipramine (15 mg/kg) showing that in both the doses, it has significant antidepressant activity.

Similarly, restraint stress caused a significant decrease in central and PL crossings in the OFT, indicating depression-like behavior. The dose of 100 mg/kg and 300 mg/kg of EELS increased the number of central and PL crossings. The observed results showed a significant dose-dependent antidepressant-like effect in OFT, as they increase in locomotor activity when compared against stress control as well as against standard imipramine (15 mg/kg) showing that in both the doses, it has significant antidepressant activity.

The exact mechanism by which extract of EELS leaves exert antidepressant action is not precisely known. The leaves of EELS contain phenolic compounds and many flavonoids such as quercetin, isoquercetin, and kaempferol. Phenolic compounds and flavonoids have been reported to inhibit catechol-O-methyltransferase and MAO enzymes in the brain. Hence, it increases the concentration of catecholamines in synaptosomes and thus attenuates

depression. Oxidative stress is one of the mechanisms involved in the pathogenesis of depression. Many natural products have an antioxidant property which is involved in reversing the deleterious effects of neuronal communication and behavior. CAT and SOD are one of the free radicals suggested, role in stress and depression either by modulating the release of other neurotransmitters, decrease in SOD activity, and decrease in CAT activity induced as a result of ARS, is an index of pro-oxidative conditions. In the present study, 12 h restraint stress-induced significant oxidative damage, as indicated by decreased SOD and decreased CAT, which was significantly normalized by EELS pre-treatment in a dose-dependent manner [22].

MDA and LPO are considered as a critical mechanism in causing cell injury during oxidative stress. Several studies have shown that restraint stress significantly elevated the concentration of MDA levels in the hippocampus of rats. The present study results show a significant concentration of MDA level, evidenced by an increased amount of MDA, which was significantly normalized by EELS pre-treatment in a dose-dependent manner. Thus, the antidepressant-like effect of EELS could be associated with its capacity to prevent the MDA and lipid peroxidative damage, induced by ARS [23].

CONCLUSION

It can be concluded from the study that the ethanolic extract of *L. speciosa* L. leaves possesses significant antidepressant property, which is probably due to flavonoids which play an active role in providing antidepressant-like effect. *L. speciosa* L. plant can be used for the treatment of neurological disorders and may be recommended as a supplement for the antidepressant activity.

ACKNOWLEDGMENT

We are grateful to SA-FORD, Plot No.: V-10, MIDC, Taloja, Navi Mumbai, Dist. Raigad Maharashtra – 410 208 for the donation of albino Wistar rats for the research. We are grateful to our Principal Dr. (Mrs.) Sudha Rathod and Prof. Imtiyaz Ansari for their guidance and support as well as to the Pharmacology Department, Oriental College of Pharmacy, Navi Mumbai.

AUTHORS' FUNDING

The authors received no specific funding for this research project work.

AUTHORS' CONTRIBUTION

We declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be done by the authors. Miss Sunita Vishwakarma performed the research work collected the data and analyzed the data. Dr. (Mrs.) Vanita Kanase proof-read the whole manuscript and suggested the necessary changes and helps in designing the manuscript.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this paper.

REFERENCES

- Santosh P, Venugopl R, Nilakasha S, Kunjibhari S, Mangala L. Antidepressant activity of methanolic extract of *Passiflora foetida* leaves in mice. *Int J Pharm Pharm Sci* 2011;3:6-9.
- Bharathi P, Seshayamma V, Jagannadharao GH, Sivakumar N. Evaluation of antidepressant activity of aqueous extract of *Withania Somnifera* [Aswagandha] roots in albino mice. *IOSR J Pharm Biol Sci* 2015;10:27-9.
- Knox GW, Norcini JG. *Lagerstroemia* cultivars under evaluation at the NFREC-monticello. *Proc Fla State Hort Soc* 1991;104:346-7.
- Turner S. Queen's Crape Myrtle (*Lagerstroemia speciosa*) and Crape Myrtle (*Lagerstroemia indica*). Miami, FL: Richard Lyons Nursery, Inc.; 2016.
- Kokate CK. *Practical Pharmacognosy*. 4th ed. New Delhi, India: Vallabh Prakashan; 2005.
- Khandelwal KR. *Practical Pharmacognosy*. Pune, Maharashtra: Nirali Prakashan; 2000.
- Azad AK, Rahman MK, Sunzida NK. Acute oral toxicity study on Malaysian traditional herb: *Lagerstroemia speciosa* L. (Banaba). *J Pharmacogn Phytochem* 2015;4:228-32.
- Mostofsky DI, Buynitsky T. Restraint stress in biobehavioral research: Recent developments. *Neurosci Biobehav Rev* 2009;33:1089-98.
- Takada T, Yoshinari N, Sugiishi S, Kawase H, Yamane T, Noguchi T. Effect of restraint stress on the progression of experimental periodontitis in rats. *J Periodontol* 2004;75:306-15.
- Lotankar AR, Wankhede S, Sharma JB, Momin AJ. Anti-stress activity of flavonoids rutin and quercetin isolated from the leaves of *Ficus benghalensis*. *Int J Pharm Pharm Res* 2016;5:5-19.
- Seyed AM, Hossein H, Fatemeh G. Evaluation of antidepressant effects of aerial parts of *Echium vulgare* on mice. *Iran J Basic Med Sci* 2007;10:189-96.
- Sulakhiya K, Patel VK, Saxena R, Dashore J, Srivastava AK, Rathore M. Effect of *Beta vulgaris* Linn. Leaves extract on anxiety and depressive-like behavior and oxidative stress in mice after acute restraint stress. *Pharmacognosy Res* 2016;8:1-7.
- Devi T, Das S. Evaluation of the protective effect of ethanolic extract of leaves of *Punica granatum* Linn. On forced swimming induced chronic fatigue syndrome in mice. *Int J Pharm Pharm Sci* 2017;9:207-12.
- Thakara VN, Dhakaneb VD, Patel BM. Potential antidepressant-like activity of silymarin in the acute restraint stress in mice: Modulation of corticosterone and oxidative stress response in cerebral cortex and hippocampus. *Pharmacol Rep* 2016;68:1020-7.
- Aebi H. In: Bergmeyer HU, editor. *Methods of Enzymatic Analysis*. New York: Academic Press; 1974. p. 674.
- Mishra HP, Fridovich I. The oxidation of phenylhydrazine: Superoxide and Mechanism. *Biochemistry* 1976;15:681-3.
- Ohkawa H, Ohisi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351-8.
- Hasrat JA, De Bruyne T, De Backer JP, Vauquelin G, Vlietinck AJ. Isoquinoline derivatives isolated from the fruit of *Annona muricata* as 5-HT_{1A} receptor agonists in rats: Unexploited antidepressive (lead) products. *J Pharm Pharmacol* 1997;49:1145-9.
- Hasrat JA, Pieters L, De Backer JP, Vauquelin G, Vlietinck AJ. Screening of medicinal plants from Suriname for 5-HT(1A) ligands: Bioactive isoquinoline alkaloids from the fruit of *Annona muricata*. *Phytomedicine* 1997;4:133-40.
- Porsolt RD, Anton G, Blavet N, Jalfre M. Behavioural Despair in Rats: A new model sensitive to antidepressant treatments. *Eur J Pharmacol* 1978;47:379-91.
- Freitas AE, Bettio L, Neis VB, Santos DB, Ribeiro CM, Rosa PB, et al. Agmatine abolishes restraint stress-induced depressive-like behavior and hippocampal antioxidant imbalance in mice. *Prog Neuropsychopharmacol Biol Psychiatry* 2014;50:143-50.
- Budni J, Zomkowski AD, Engel D, Santos DB, dos Santos AA, Moretti M, et al. Folic acid prevents depressive-like behavior and hippocampal antioxidant imbalance induced by restraint stress in mice. *Exp Neurol* 2013;240:112-21.
- Kolla N, Wei Z, Richardson JS, Li XM. Amitriptyline and fluoxetine protect PC12 cells from cell death induced by hydrogen peroxide. *J Psychiatry Neurosci* 2005;30:196-201.