

**TAMARINDUS INDICA L. (FABACEAE): EXTENT OF EXPLORED USE IN TRADITIONAL MEDICINE**BALAKRISHNA VUYVALA<sup>1\*</sup>, D SENTHIL KUMAR<sup>2</sup>, THAKKALAPALLY LAKSHMI<sup>1</sup><sup>1</sup>Department of Pharmacology, Guru Nanak Institutions Technical Campus-School of Pharmacy, Hyderabad, Telangana, India. <sup>2</sup>Department of Pharmacology, Annamalai University, Chidambaram, Tamil Nadu, India. Email: balakrishnavuyyala@gmail.com

Received: 21 December 2019, Revised and Accepted: 18 January 2020

**ABSTRACT**

Tamarind (*Tamarindus indica* L.) belongs to the family of Fabaceae (Leguminosae), subfamily *Caesalpinioideae*, is a very important food within the tropics. Medicinal plants are the rear bone of traditional medicine (TM). TM is vital in tropical countries: Contrary to pharmaceuticals, pharmacological, and pharmacotherapy. *T. indica* is employed as TM in India, Africa, Pakistan, Bangladesh, Nigeria, and most of the tropical countries. It is used traditionally in abdominal pain, diarrhea and dysentery, helminths infections, wound healing, malaria and fever, constipation, inflammation, cell cytotoxicity, gonorrhoea, and eye diseases. It is numerous chemical values and is rich in phytochemicals, and hence, the plant is reported to possess antidiabetic activity, antimicrobial activity, antivenom activity, antioxidant activity, antimalarial activity, hepatoprotective activity, antiasthmatic activity, laxative activity, and antihyperlipidemic activity. Thus, the aim of the present review demonstrates the plant contains in leaves, seeds, roots, pulp, fruits, and flowers an excellent sort of bioactive substances that have beneficial effects on human health and therefore the possibility of application in various tropical, pharmaceutical, and industrial sectors.

**Keywords:** Tamarind, Antioxidant, Antidiabetic, Antimicrobial, Anti-inflammatory, hepatoprotective.© 2020 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2020.v13i3.36676>**INTRODUCTION**

*Tamarindus indica* L. of the Fabaceae, subfamily *Caesalpinioideae*, is a crucial food within the tropics. Medicinal plants are the rear bone of traditional medicine (TM). TM is vital in tropical countries: Contrary to pharmaceuticals, pharmacological, and pharmacotherapy [1]. It is often freely and readily available multipurpose tree of which just about every part finds a minimum of some use either medicinal or nutritional. For instance, in Burkina Faso, up to 90% of the population relies to use traditional remedies [2]. Tamarind is indigenous to tropical Africa, but it is been introduced and naturalized worldwide in over 50 countries. Plants are the essential elements of TM and selected as a therapy in greater amounts. *T. indica* is employed as a standard medicine in Asian countries India, Thailand, Pakistan, Bangladesh, Sri Lanka, and most of the tropical countries. In America, Mexico and Costa Rica are the most important producers. Africa on the entire does not produce tamarind on a billboard scale, though it is widely employed by the local people. Minor producing countries in Africa are Senegal, Gambia, Kenya, Tanzania, and Zambia [3].

*T. indica* (Fig. 1) is of moderate to enormous in size, evergreen tree, up to 24 m in stature and 7 m in size that has light yellow and pink flowers [4]. The continued morphologic and subatomic examinations and proceeded with the study will explain the accurate situating of *Tamarindus* in connection to its putatively related genera [5-7]. It is an enormous evergreen tree with an outstandingly wonderful spreading crown and is developed during nearly the whole nation, apart from within the Himalayas and Western dry regions [8].

Practically, all parts of the tree discover some utilization or the opposite in nourishment, concoction, pharmaceutical, and textile enterprises, and as grain, timber, and fuel [9,10]. Commercial ranches are accounted for in Central American nations and in North Brazil [11]. Large fragments of human population and animals within the creating nations experience the ill effects of protein unhealthiness. *T. indica* is wealthy in supplements and assumes a big role in human nourishment, basically within the developing nations [12,13]. *T. indica* contains significant levels of rough protein. *T. indica* likewise contains a big level of protein with numerous fundamental amino acids, which help

to create strong and efficient muscles. *T. indica* is additionally high in starch, which provides vitality, and is rich in minerals, such as calcium, phosphorus, potassium, calcium, and magnesium. *T. indica* can likewise give littler measures of iron and nutrient A. The whole plant of *T. indica* Linn. is employed widely for extensively for contemporary reason; subsequently, it is exceptionally gainful to the human beings [14].

The pulp contains organic acids, such as hydroxy acid, ethanolic acid, malic acid, and carboxylic acid; amino acids; carbohydrate (25–30%) [15]; pectin; protein; fat; some pyrazines (trans-2-hexenal); and a few thiazoles (2-ethylthiazole, 2-methylthiazole) as fragrant [16]; and therefore, the seed polysaccharides are found with a main chain consisting of  $\beta$ -1,4-connected glucose molecules alongside xylose

**Table 1: Amino acid content of *T. indica* fruit pulp and seeds**

S. No.	Amino acid		Fruit pulp mg/g dw	Seeds g/100 g dw
1.	Crude protein (total protein)		116.00	17.30
2.	Aspartic acid	ASP	12.00	1.80
3.	Glutamic acid	GLU	16.70	2.82
4.	Serine	SER	6.88	0.95
5.	Glycine	GLY	5.15	1.67
6.	Histidine	HIS	3.37	0.55
7.	Arginine	ARG	8.74	1.66
8.	Threonine	THR	6.05	0.52
9.	Alanine	ALA	6.20	0.65
10.	Proline	PRO	7.61	0.85
11.	Tyrosine	TYR	4.34	0.95
12.	Valine	VAL	6.97	0.71
13.	Methionine	MET	2.48	0.14
14.	Isoleucine	ILE	5.20	0.67
15.	Leucine	LEU	8.89	1.09
16.	Phenylalanine	PHE	4.78	0.71
17.	Lysine	LYS	8.22	1.05
18.	Cysteine	CYS	1.35	0.35
19.	Tryptophan	TRP	1.04	0.18

*T. indica*: *Tamarindus indica*



Fig. 1: (a) Fruits, (b) leaves, (c) flowers, (d) stem bark of *Tamarindus indica*

Table 2: Mineral content of tamarind fruit pulp

S. No.	Minerals	A	B	C	D	E	F
1.	Aluminum	-	-	-	1.84	-	-
2.	Iron	8.49	6.80	14.00	3.17	2.80	1.69
3.	Sodium	76.66	11.10	-	6.21	28.00	13.95
4.	potassium	62.00	1226.90	-	0.65	628.00	790.11
5.	Magnesium	72.03	128.20	-	0.12	92.00	53.28
6.	Calcium	465.75	17.10	240.00	0.19	74.00	106.88
7.	Cobalt	-	-	-	0.01	-	0.05
8.	Copper	21.83	-	-	0.91	0.09	0.29
9.	Barium	-	-	-	0.20	-	-
10.	Manganese	-	-	-	21.50	-	0.06
11.	Molybdenum	-	-	-	0.01	-	-
12.	Chromium	-	-	-	0.29	-	-
13.	Nickel	0.52	-	-	0.13	-	0.08
14.	Phosphorus	91.00	108.10	-	0.12	113.00	99.49
15.	Zinc	1.06	-	2.30	1.32	0.10	0.09
16.	Lead	-	-	-	0.01	-	-
17.	Titanium	-	-	-	0.02	-	-
18.	Strontium	1.06	-	2.30	1.32	0.10	0.09

Units are mg/100 g dw; -: Not mentioned in the original paper. Source: A: Ishola (1990); B: Saka and Msonthi (1994); C: Nordeide (1996); D: Glew (2005); E: USDA (2007, cited in Almeida, 2009); F: Almeida (2009)

Table 3: Mineral content of tamarind seeds

S. No.	Minerals	A	B	C	D	E	F
1.	Calcium	786.86	172.00	-	185.00	142.00	36.60
2.	Copper	18.97	0.47	0.73	1.16	0.26	2.10
3.	Iron	ND	6.30	13.70	2.67	9.09	45.50
4.	Potassium	610.00	1430.00	-	-	-	1308.00
5.	Magnesium	118.00	214.00	28.20	196.00	201.00	104.00
6.	Manganese	-	0.68	0.70	ND	0.70	12.10
7.	Molybdenum	-	-	-	1.39	-	-
8.	Sodium	19.17	21.30	-	ND	-	8.90
9.	Nickel	ND	-	-	-	-	-
10.	Phosphorous	165.00	312.00	-	228.00	220.00	-
11.	Zinc	3.00	7.100	1.22	2.63	3.12	7.00

Units are mg/100 g dw; ND: not detected; -: Not mentioned in the original paper. Source: A: Ishola (1990); B: Siddhuraju (1995); C: Smith (1996); D: Glew (1997); E: Lockett (2000); F: Ajayi (2006)

(alpha-1,6) and galactose [17,18]; total protein; lipids with fatty oils; and a few keto acids [19]. Within the leaves of the plant, two triterpenes, lupanone and lupeol, were found [20].

The aerial parts of this plant have exhibited the appearance of tartaric acid, ethanolic acid, and succinic acid, gum, pectin, sugar, tannins, alkaloid, flavonoids, sesquiterpenes, and glycosides [21,22]. *T. indica* pericarp and seeds contain phenolic antioxidants to stop cancer. The leaf oil contains 13 components among which limonene and benzyl benzoate were most predominant [23]. Phytochemical screening of

the basis bark of *T. indica* demonstrated the presence of n-hexacosane, eicosanoic acid, octacosanylferulate, 21-oxobehenic acid, and (+)-pinitol. The appearance of the bioactive compound (+)-pinitol during this plant is being accounted for the primary time [24]. The unpredictable constituents of the fruit pulp were furan derivatives (44.4%) and acid (33.3%) of the entire volatiles [25]. The many unsaturated fats of seeds were hexadecanoic acid, monounsaturated fatty acid, linoleic acid, and eicosanoic acid. The unsaponifiable issue from the seed oil of *T. indica* indicated the presence of  $\beta$ -amyrin, compesterol,  $\beta$ -sitosterol, and 7-hydrocarbons [26].





