

## CORONAVIRUS DISEASE 2019 VACCINATION DRIVE: THE PERCEPTIONS AND ACCEPTANCE OF VACCINATION AMONG HEALTH CARE WORKERS

KASI VISWANATH<sup>1\*</sup>, HIREMATH RN<sup>2</sup>, SHARANJIT SINGH BASRA<sup>3</sup>, NIRAJ CHOUREY<sup>4</sup>

<sup>1</sup>Department of ???, ???, Meerut, Uttar Pradesh, India. <sup>2</sup>Department of ???, ???, Bengaluru, Karnataka, India. <sup>3</sup>Department of ???, ???, Amritsar, Punjab, India. <sup>4</sup>Department of ???, ???, Babina, Uttar Pradesh, India. Email: dr.kasiviswanath@gmail.com

Received: 04 March 2021, Revised and Accepted: 20 May 2021

### ABSTRACT

**Objective:** Coronavirus disease 2019 (COVID-19) infection evolved as a worst nightmare across entire humanity especially for health care workers along with all stakeholders of the society ever since the beginning of year 2020. Vaccination campaign is one of the most successful public health interventions and a cornerstone for the prevention of communicable infectious diseases. To have a baseline data and to form the basis for further vaccination drive, we carried out this study with an objective to understand the perception and COVID-19 vaccine hesitancy among health care workers (HCWs).

**Methods:** A correctional study was carried out at one of the multispecialty hospital in North India where COVID-19 vaccination for HCWs was planned and 675 HCWs were taken into the study. Data were collected by means of personal interview using pretested semi-structured questionnaire and Epi Info version 6 was used for analysis.

**Results:** The mean age of the participants was 32.78 years. Majority were male (89.48%), working in government sector (87.7%) and paramedic (88.15%). While 94.67% were accepting vaccine voluntarily while remaining were not confident enough on their own, only 26.96% were ready to take vaccine even on payment. The overall awareness scores of study participants were calculated and found that 79.56% of them had poor awareness level while only 20.44% had satisfactory awareness level. The association of awareness score was statistically significant with job profile (designation), gender, and prior COVID-19 positivity of study participants.

**Conclusion:** The decision of vaccine prioritization to HCWs and frontline workers will boost the morale and spirit of warriors high in future. This study was a preliminary attempt to check any vaccine hesitancy factors among HCWs and act accordingly. Although almost all had volunteered for vaccine, the knowledge about vaccine was poor in majority of HCWs as the time period was recent to vaccine rollout and detailed research findings were not yet out about vaccine. All measures to be taken including intensive information education and communications activities targeting HCWs and general population on regular basis to upgrade the knowledge on vaccine issues. A strong communicating policy in an era of misinformation is highly recommended as social media plays an important role in spreading true as well as false messages jeopardizing the entire gains of successful vaccination and attainment of herd immunity. HCWs to act as role model for general public and patients which would in turn increase the vaccine acceptance among them.

### Keywords:

© 2021 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2021v14i6.41971>. Journal homepage: <https://innovareacademics.in/journals/index.php/ajpcr>

### INTRODUCTION

Coronavirus disease 2019 (COVID-19) is super-infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). COVID-19 infection evolved as a worst nightmare across entire humanity especially for health care workers (HCWs) along with all stakeholders of the society ever since the beginning of year 2020, when the World Health Organization (WHO) declared it as pandemic with rising infection rate and severity across the world. This virus caused respiratory illness ranging from the common cold to severe disease. It caused illness across all age group but severely affected older people and those with underlying medical problems such as cardiovascular disease, diabetes, chronic respiratory disease, and cancer [1]. It was first identified in Wuhan, China, in December 2019 and is an on-going pandemic. In India, the first case reported in Kerala state in January 2020, being novel virus there was no specific treatment and vaccine till end of year 2020.

The control of COVID-19 is a huge challenge due to fear of unknown, India was quick to close its international borders and enforce an immediate lockdown [2] for emergency preparedness and response. As of December 27, 2020, there have been over 79.2 million cases and over 1.7 million deaths reported globally since the start of the pandemic,

whereas in India, alone total cumulative cases were over 10,187,850 and 147,622 deaths reported [3]. There is significant reduction of HCWs capacity through periods of infection by quarantine, isolation guidelines, and round the clock hectic work schedule under stress.

After various public health measures such as nation-wide lockdown, containment zones, isolation centers, quarantine centers to simple social distancing, masking, and hand washing, the world was eagerly waiting for vaccination campaign which is one of the most successful public health interventions and a cornerstone for the prevention of communicable infectious diseases. Notwithstanding vaccine progress, ongoing public acceptance is required to maintain herd immunity, prevent outbreaks of vaccine preventable illnesses, and ensure adoption of novel vaccines [2]. Any new vaccine introduction in short period of span creates lots of doubts and hesitancy among both HCWs and general public as we have seen in H1N1 infection too [4]. Vaccine hesitancy refers to delay in acceptance or refusal of vaccination despite availability of vaccination services. Vaccine hesitancy is complex and context specific, varying across time, place, and vaccines. It is influenced by factors such as complacency, convenience, and confidence [5]. Vaccine hesitancy will create isolation from vaccination for any individual when they triggered by variety of unknown sources of information either through social media or message sharing messengers without health

editorial oversight [6], which is extreme potential threat to public health interventional strategies. The simple forward of message leads full throated castigation which compromises the vaccine development for novel viruses.

At one side, HCWs are strongest influencers in vaccination decisions as common population and patients show greater resistance to accept novel vaccines [7], on the other side, COVID-19 pandemic has reminded all of us of the vital role health workers played to relieve suffering and save lives. Thus, recognizing health worker safety as a priority for patient safety, WHO Member States, and all relevant stakeholders urged to take urgent and sustainable action and in the context of emergency response, giving health workers priority access to newly licensed and available vaccines [8].

The union Health Ministry of India advised all states to commence COVID-19 vaccination drive to HCWs on January 16, 2021, as priority one and approved two vaccines for restricted emergency use in India [9]. The massive Phase 1 vaccination drive carried out with "Covishield manufactures by Serum Institute of India" and indigenously developed "Covaxine" of Bharat Biotech which has been offered two doses for free of cost to all HCWs and frontline workers (FLWs).

In view of above, to have a baseline data and to form the basis for further vaccination drive, we carried out this study with an aim to understand the perception and COVID-19 vaccine hesitancy among HCWs. At our study area, Covishield vaccine was provided by Government of India to all HCWs. COVISHIELD is ChAdOx1-S/nCoV-19 (recombinant) vaccine which again is a recombinant vaccine with replication deficient chimpanzee adenovirus vector encoding the SARS-COV-2 spike glycoprotein. The vaccine is genetically modified in human embryonic kidney 293 cells [10].

## METHODS

A correctional study was carried out at one of the multispecialty hospital in North India where COVID-19 vaccination for HCWs was planned. COVID-19 vaccination was planned for approximately 1010 HCWs with target of 100–150 per day. About 95% of vaccination was achieved in 10 days. On each day before inoculation, pre-tested semi-structured questionnaire was used to gather information about awareness and vaccine acceptance/hesitancy among HCWs and data were collected by means of personal interview.

Expecting the vaccine acceptance and awareness rate of 50%, alpha 5%, and chance error 5%, the sample size calculated was 385. However, since 675 HCWs were willing for the study and gave consent, 675 HCWs were taken into the study. All those who refused for consent and unwilling were excluded from study. Data collected were entered into Excel sheet and Epi Info version 6 was used for analysis. Chi-square was used for proportions comparison at statistical significance at  $p < 0.05$ . To assess collective awareness, set of 12 questions was selected. One mark for given for right answer and zero mark for wrong. Total scores were divided into two categories, poor awareness (<50% marks) and satisfactory awareness (more than 50% marks).

## RESULTS

The mean age of the participants was 32.78 years. Majority were male (89.48%), working in government sector (87.7%) and paramedic (88.15%). About 41 (6.07%) of the participants have suffered with COVID-19 earlier with 2 (0.30%) of them suffered 6 months ago and 39 (5.78%) of them suffered 3 months ago. Thirty-two (4.74%) of them had symptoms related to COVID-19 but not tested while one had their family member with COVID-19-related symptoms earlier. All of them 675 (100%) knew about two variety of vaccines but only 130 (19.26%) knew about the correct type of vaccine available at their center before being administration, implying that irrespective of type of vaccine, all HCWs were ready to take vaccine, which was a good sign of vaccine acceptance.


About 100% knew about two variety of vaccines launched by government and 19.26% knew about the type of vaccines allotted to HCWs at their center. When asked about benefits of taking COVID-19 vaccine, 100% said that it will protect us from COVID-19 infection while 64.59% of them said that it protects family members too. About 74.67% said that both vaccines are equally effective while 19.26% said that Covishield is better than Covaxine. When asked about reasons for thinking the vaccine is best, 27.70% told it's because of lesser side effects, easy availability (4.74%), and good efficacy (9.93%) (Table 2). Majority (62.67%) did not know about when immunity is achieved after taking vaccine while 23.85% told that they will be immune within 1–2 weeks. Surprisingly, 31.26% told that vaccine is effective against mutant strains while 36.30% did not know anything about it. Majority of them rightly knew about side effects which included pain (96.15%), fever (65.48%), allergic reactions (48.04%), and paralysis (8.15%). Even at this point when cases and deaths are declining in this country, 49.19% HCWs agreed that vaccine is required and 26% felt that vaccine is required even for those who contracted COVID-19 disease earlier.

While 94.67% were accepting vaccine voluntarily while remaining were not confident enough on their own, only 26.96% were ready to take vaccine even on payment.

The overall awareness scores of study participants were calculated and found that 79.56% of them had poor awareness level while only 20.44% had satisfactory awareness level (Fig. 1). The association of awareness score was statistically significant with job profile (designation), gender, and prior COVID-19 positivity of study participants (Table 5).

**Table 1: Sociodemographic profile of study participants**

Gender	Frequency	Percent
Female	71	10.52
Male	604	89.48
Age category		
<25 years	148	21.93
25–35 years	309	45.78
35–45 years	151	22.37
Above 45 years	67	9.93
Job category		
Government	592	87.70
Private	83	12.30
Designation		
Medical officer	38	5.63
Nursing staff	42	6.22
Paramedic	595	88.15
Total	675	100.00



**Fig. 1: Awareness level among study participants**

## DISCUSSION

Ever since the COVID-19 pandemic, countries across the world were trying hard for an effective vaccine. Finally, vaccines of Moderna, Pfizer BioNtech, and Oxford Astrazeneca were recently approved [11]. India also approved two vaccines on January 3, 2021, for restricted use under emergency [9] and finally vaccination drive for high-risk individuals was started on January 16, 2021, for HCWs. This study was carried out between January 16, 2021, and January 30, 2021, among HCWs at the COVID-19 vaccination center before getting vaccinated. Our study was

**Table 2: Awareness regarding available COVID-19 vaccine in India**

Variables	Frequency (n=675)	Percent
Best available vaccine in India (vaccine effectiveness)		
Covaxine	41	6.07
Covishield	130	19.26
Both	504	74.67
Reasons for thinking this vaccine is best		
Do not know	389	57.63
Easily available	32	4.74
Good efficacy	67	9.93
Less side effects	187	27.70
After getting vaccine, immunity is achieved		
After 3–5 weeks	46	6.81
After 6–8 weeks	20	2.96
After 8 weeks	25	3.70
Do not know	423	62.67
Within 1–2 weeks	161	23.85
Does vaccine effective against mutant strains		
Do not know	245	36.30
No	219	32.44
Yes	211	31.26
Vaccine side effects		
Pain	649	96.15
Paralysis	55	8.15
Allergic reaction	332	48.04
Fever	442	65.48
Others	–	–

COVID-19: Coronavirus disease 2019

**Table 3: Acceptance of COVID-19 vaccine among study participants**

Variables	Frequency	Percent
Acceptance for vaccine voluntarily		
No	36	5.34
Yes	639	94.67
Acceptance for vaccine voluntarily even if available on payment		
Can't tell, depends on cost	53	7.85
No	440	65.19
Yes	182	26.96

COVID-19: Coronavirus disease 2019

**Table 4: Graded response of study participants on COVID-19 vaccine (On a visual analog scale ranging from 0 [minimum] to 5 [maximum])**

Variables	Grading (%)					
	0	1	2	3	4	5
Vaccine effectiveness	–	–	–	116 (17.19)	208 (30.81)	351 (52.00)
Requirement of vaccine based on current scenario	–	–	–	–	227 (33.63)	448 (66.37)
Vaccine side effects	366 (54.22)	174 (25.78)	135 (20.00)	–	–	–
Requirement of taking vaccine by person who is previously COVID positive	31 (4.59)	176 (26.07)	332 (49.19)	136 (20.15)	–	–

COVID-19: Coronavirus disease 2019

an effort to see the perceptions of HCWs about vaccine acceptance/hesitancy as acceptance by HCWs would influence drastically the vaccine acceptance by general population and would act as a role model to others.

In our study, 100% of participants said that they would continue practices of COVID-19 precautionary measures, even after vaccination also, which was very appreciable. About 80.44% of participants strongly believed that beneficiaries could spread infection and can get infected (73.93%) even after vaccination. When majority believed that they could get infected even after vaccination and can spread infection too, then ideally, they should have understood that quarantine would be essential to some extent even after vaccination but only 43 % of HCWs felt so. This indicates that even HCWs had confused ideas about quarantine basics.

**Vaccine acceptance**

Vaccine rollout in India was done when COVID-19 cases and deaths were on declining trend. However, majority (49.19%) of study participants said that vaccine is required irrespective of present COVID-19 scenario, which was a good sign considering the second wave impacting the other countries during this time. Although only 20% of study participants had satisfactory awareness level, the voluntary acceptance rate for vaccine was very high (94.67%). Such a high acceptance level was very good and would influence general population also. The low awareness level about vaccine is probably because of vaccination rollout for HCWs was quick and in-depth awareness about vaccine details was low initially as detailed research findings were not yet out about vaccine. We could not find any study in this area, which was immediately carried out on vaccine rollout to assess the acceptance in India, as the authors were quick to carry out this study to see any hesitancy among HCWs so that any need to increase actions in terms of information education and communications (IEC) activities be found out at the earliest. Our study emphasizes the need for regular update of knowledge on vaccinations among HCWs, although it is a novel vaccine but true in-depth information about vaccine must be disseminated among HCWs and general populace at the earliest, before any hesitancy is developed among them against vaccine.

Poor awareness score was seen among 83.19% of paramedic and 61.90% of nursing staff then compared to medical officers, warranting specific IEC, and regular update to them about all aspects of COVID-19 as the association between designated HCWs and awareness score was statistically significant. Poor awareness level was also seen among 80% of HCWs who did not have COVID-19 positivity earlier as compared to 68% of those who had prior COVID-19 positivity. The association between prior COVID-19 positivity status and awareness score was statistically significant. This indicates that HCWs who were COVID positive earlier were showing more interest on gathering information about vaccine than compared to COVID-negative HCWs. Thus, these HCWs should be targeted more.

The study carried out by Al-Qerem and Jarab [12] in Middle Eastern population, showed very low intend (36.8%) for vaccine even when



- HJ. Vaccine hesitancy and healthcare providers. *Vaccine* 2016;34:6700-6.
6. Puri N, Coomes EA, Haghbayan H, Gunaratne K. Social media and vaccine hesitancy: New updates for the era of COVID-19 and globalized infectious diseases. *Hum Vaccin Immunother* 2020;16:2586-93.
  7. Wiley KE, Massey PD, Cooper SC, Wood N, Quinn HE, Leask J. Pregnant women's intention to take up a post-partum pertussis vaccine, and their willingness to take up the vaccine while pregnant: A cross sectional survey. *Vaccine* 2013;31:3972-8.
  8. World Health Organization. World Patient Safety Day. Geneva: World Health Organization; 2020. Available from: <https://www.who.int/campaigns/world-patient-safety-day/2020>. [Last accessed on 2021 Feb 20].
  9. Press Statement by the Drugs Controller General of India (DCGI) on Restricted Emergency Approval of COVID-19 Virus Vaccine. PIB GOI; 2021 Available from: <https://www.pib.gov.in/PressReleaseDetail.aspx?PRID=1685761>. [Last accessed on 2021 Feb 20].
  10. World Health Organization. Astra Zeneca ChAdOx1-S/n CoV-19, COVID-19 Vaccine. Geneva: World Health Organization; 2021. Available from: <https://www.who.int/publications/m/item/chadox1-s-recombinant-covid-19-vaccine>. [Last accessed on 2021 Feb 20].
  11. COVID-19 Vaccine. Vaccine Safety, Distribution and FAQs/UC Health. Available from: <https://www.uhealth.org/services/infectious-diseases/coronavirus-covid-19/covid-19-vaccine>. [Last accessed on 2021 Feb 20].
  12. Al-Qerem WA, Jarab AS. COVID-19 vaccination acceptance and its associated factors among a Middle Eastern population. *Front Public Health* 2021;9:632914.
  13. Lazarus JV, Ratzan SC, Palayew A, Gostin LO, Larson HJ, Rabin K, et al. A global survey of potential acceptance of a COVID-19 vaccine. *Nat Med* 2020;27:225-8.
  14. Al-Mohaithef M, Padhi BK. Determinants of COVID-19 vaccine acceptance in Saudi Arabia: A web-based national survey. *J Multidiscip Healthc* 2020;13:1657-63.
  15. Ward JK, Alleaume C, Peretti-Watel P. The French public's attitudes to a future COVID-19 vaccine: The politicization of a public health issue. *Soc Sci Med* 2020;265:113414.
  16. Wang J, Jing R, Lai X, Zhang H, Lyu Y, Knoll MD, et al. Acceptance of COVID-19 vaccination during the COVID-19 pandemic in China. *Vaccines* 2020;8:482.
  17. Sarasty O, Carpio CE, Hudson D, Guerrero-Ochoa PA, Borja I. The demand for a COVID-19 vaccine in Ecuador. *Vaccine* 2020;38:8090-8.
  18. Fisher KA, Bloomstone SJ, Walder J, Crawford S, Fouayzi H, Mazor KM. Attitudes toward a potential SARS-CoV-2 vaccine: A survey of U.S. adults. *Ann Intern Med* 2020;173:964-73.
  19. Neumann-Böhme S, Varghese NE, Sabat I, Barros PP, Brouwer W, van Exel J, et al. Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *Eur J Health Econ* 2020;21:977-82.
  20. Arede M, Bravo-Araya M, Bouchard É, Gill GS, Plajer V, Shehraj A, et al. Combating vaccine hesitancy: Teaching the next generation to navigate through the post truth era. *Front Public Health* 2019;6:381.
  21. Swingle CA. How do we approach anti-vaccination attitudes? *Mo Med* 2018;115:180-1.
  22. Johns Hopkins Bloomberg School of Public Health. COVID-19 Vaccine Acceptance Falling Globally and, in the U.S., Survey Finds; 2021. Available from: <https://www.jhsph.edu/news/news-releases/2021/covid-19-vaccine-acceptance-falling-globally-and-in-the-us-survey-finds.html>. [Last accessed on 2021 Feb 20].

#### Author Queries???

- AQ1: Kindly provide department  
 AQ2: Kindly provide University or Institute name  
 AQ3: Kindly provide department  
 AQ4: Kindly provide history details  
 AQ5: Kindly review the sentence as it seems to be incomplete.  
 AQ6: Kindly provide minimum three keywords  
 AQ7: Kindly cite tables 1, 3 and 4 in the text part  
 AQ8: Kindly provide column head