

CLINICAL SPECTRUM AND OUTCOME OF KEROSENE POISONING IN PEDIATRIC AGE GROUP IN SOUTH KASHMIR: A PROSPECTIVE OBSERVATIONAL STUDY

ANDLEEB MAJEED¹, MOHD ASHRAF BHAT¹, MALIK SHAKIR HUSSAIN¹, SHOWKAT HUSSAIN TALI^{1*}, SHAGUFTAYOUSUF², ZULEIDAIN HASSAN¹, BILAL AHMAD NAJAR¹, SHEIKH MUSHTAQ AHMAD²

¹Department of Pediatrics, Government Medical College, Anantnag, Jammu and Kashmir, India. ²Department of Obstetrics and Gynaecology, Government Medical College, Anantnag, Jammu and Kashmir, India. E-mail: anantnag.drshowkatshifa@gmail.com

Received: 02 April 2017, Revised and Accepted: 05 May 2022

ABSTRACT

Objective: The objective of the study was to study the clinical profile of kerosene poisoning, its complications and outcome in children in South Kashmir, J and K, India.

Methods: This prospective study of children admitted with kerosene poisoning was conducted in a Govt. Medical College Hospital from January 2019 to December 2021. Demographic data, risk factors, clinical features, and the outcome recorded as per predesigned pro forma were analyzed.

Results: A total of 199 children were admitted with acute poisoning and 77 (38.6%) had kerosene poisoning. Majority of the patients was male (77%) in the age group of 1 year–3 years (80.5%) and from an urban background (80.5%). Clinical features mainly included cough (75.3%), vomiting (64.9%), and fever (50.6%). X-ray was showing right lower lobe pneumonia in 23.3% followed by bilateral peri-hilar infiltration (18.1%). The most common risk factors include improper storage and lack of knowledge in family.

Conclusion: Kerosene poisoning is a common acute poisoning in pediatric age group. Male children in the age group of 1–3 years are most commonly affected. Improper storage is the most common reason for the accidental poisoning.

Keywords: Kerosene, Poisoning, Children.

© 2022 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2022v15i6.44822>. Journal homepage: <https://innovareacademics.in/journals/index.php/ajpcr>

INTRODUCTION

Acute poisoning is identified as the fourth leading cause of injury-related mortality in children. Morbidity of poisoning varies across different geographic regions due to variable cultural, social, and economic factors [1]. Kerosene is a hydrocarbon which still remains as a major fuel used for cooking in rural India and its toxicity depends on its constituents and low viscosity. Kerosene poisoning is a preventable cause of morbidity and mortality in the developing countries. Low socioeconomic status, unsafe storage, and large family size are various risk factors reported so far and may change worldwide due to different sociogeographic locations [2].

Ingestion of large quantity of kerosene is rare because of its foul smell and taste. Children can be intoxicated by ingestion, inhalation, or contact of kerosene oil with skin. Primarily, it causes pulmonary complications including chemical pneumonitis, hypoxia, emphysema, and pneumothorax. Signs and symptoms of respiratory system appear within 30 min after aspiration and progress during 48 h and then subside by 1 or 2 weeks. Other manifestations include abdominal pain, gastroenteritis, vomiting, drowsiness, and convulsions [3].

There are many studies on kerosene poisoning worldwide but, to the best of our knowledge, there are no studies on kerosene oil poisoning among children in South Kashmir, J and K, India. This study was performed to determine the prevalence of kerosene poisoning and analysis of clinical profile and its complications in children admitted to a medical college hospital in South Kashmir.

METHODS

This was a prospective observational study conducted between January 2019 and December 2021. All the children admitted with the diagnosis of accidental kerosene ingestion were enrolled for the study.

All the details with special reference to demographic factors, time of ingestion and the probable quantity ingested, clinical manifestations, complications, radiological findings, and transfer rate to a higher center were recorded in a predesigned pro forma and analyzed. Methods to look for a statistical significance were not used as they are not thought to be useful for observation studies [4].

RESULTS

Baseline data and demographic characteristics are depicted in Table 1. Clinical features and complications are depicted in Table 2. Risk factors, recovery, and referral rates are depicted in Table 3.

DISCUSSION

This prospective observational study was conducted at the Department of Pediatrics, Govt. Medical College, Anantnag, J and K, India, over a period of 3 years between January 2019 and December 2021. A total of 77 patients had ingested kerosene. Majority of the patients (80.5%) were in the age group of 1–3 years. The most common clinical features were cough (75.3%) and vomiting (64.9%) followed by fever (50.6%) and dyspnea (45.4%). Only one patient developed convulsions. Half of the patient did not develop any radiological changes. In patients who developed radiological changes, the right lower lobe infiltrates (23.3%) followed by perihilar infiltrates (18.1%) were the most common. Inadequate parental supervision (77.9%) and unsafe storage (75.3%) were the most common risk factors. Almost all (97.4%) recovered after symptomatic treatment and only 2 (2.6%) were shifted to a higher center for further care.

Out of 199 poisoning cases, 77 had ingested kerosene (38.7%). This suggests that the kerosene poisoning is an important cause of acute poisoning in pediatric age group even today. More than half of cases were

Table 1: Baseline data and demographic characteristics

Year/ characteristics	Total poisoning cases	Kerosene poisoning/ cases no. (%age)
2019	73	29 (39.7)
2020	65	25 (38.4)
2021	61	23 (37.7)
Total	199	77 (38.6)
Age	<1 year	0 (0)
	1-3 years	62 (80.5)
	>3 years	15 (19.5)
Gender	Male	55 (72)
	Female	22 (28)
Residence	Urban	62 (80.5)
	Rural	15 (19.5)

Table 2: Clinical and radiological features are depicted

Clinical features	Radiological patterns	No. of children/%age
Fever		39 (50.6)
Vomiting		50 (64.9)
Cough		58 (75.3)
Dyspnea		35 (45.4)
Cyanosis		12 (15.5)
Grunting		10 (12.9)
Drowsiness		5 (6.49)
Convulsion		1 (1.29)
Abdominal pain		15 (19.4)
	Normal	39 (50.6)
	Right lower lobe infiltrate	18 (23.3)
	Bilateral perihilar infiltration	14 (18.1)
	Bilateral lower lobe infiltration	5 (6.49)
	pneumothorax	0 (0)
	Subcutaneous emphysema	1 (1.29)

Table 3: Risk factors, recovery, and referral rates are depicted

Risk factors	Number of patients	Percentage
Unsafe storage	58	75.3
Economic problems in a family	25	32.4
Inadequate supervision of the child	60	77.9
Employed parents	22	28.5
Uneducated parents	15	19.4
Farming parents	5	6.49
Recovered	75	97.4
Transferred to a higher center	2	2.6

in the age group of <3 years. This is the age when kids start exploring the environment and are inquisitive. In addition, small kids are more often brought to medical attention for the development of symptoms at a lower ingested amount. Males outnumbered females (72 vs. 28). Male dominance is observed throughout the world and may be because of differences in socialization. All cases were poisoned accidentally as no patient was with a history of intentional intake of kerosene. Most of poisoned children were from urban areas (80.5%). This may be for the easy access of urban population to our hospital and for the fact that the patients from rural areas are either kept at home for mild or no symptoms, or managed at nearby hospitals or shifted to pediatric hospital at Srinagar (capital city) directly. Reasons for accidentally poisoning were easy availability (and within the reach), improper storage (in cold drink bottles), and less supervision of parents (working parents). Similar findings have been observed by other researchers too [1-6]. The most common reported symptom was cough in 58 children (75%). Majority of the patients (50.6%) had normal X-ray findings. Patients who developed

radiological changes and right lower lobe infiltrates (23.3%) were the most common closely followed by perihilar infiltrates (18.1%). Our findings were consistent with the finding of the other studies [6-9].

CONCLUSION

Kerosene ingestion is an important cause of acute poisoning in children. Male children in the age group of 1-3 years are most commonly affected. Improper storage is the most common risk factor. The most common symptoms are respiratory and gastrointestinal. The right lower lobe infiltrates are the most common radiological finding. Nearly, all the patients recover after supportive care and only a minority need referral to a higher center for critical care.

What is known?

Kerosene poisoning has been important cause of poisoning in children throughout the world.

What our study adds?

1. Kerosene poisoning is an important cause of poisoning childhood poisoning in South Kashmir, India.
2. Kerosene poisoning is an important cause of childhood poisoning even today, especially when the use of kerosene has declined significantly throughout the world.

ACKNOWLEDGMENT

Andleeb Majeed: Involved in conception and designing of the study, drafting of the article, analyzing the data, and final approval. Sheikh Mushtaq Ahmad, Mohd Ashraf Bhat, Malik Shakir Hussain, ZulEidain Hassan, Bilal Ahmad Najar, and Showkat Hussain Tali: All involved in data acquisition, revising the article critically, and final approval. All the authors are accountable for the accuracy of the data.

CONFLICTS OF INTEREST

None.

FUNDING

None.

DISCLAIMER

None.

REFERENCES

1. Dayasiri MB, Jayamanne SF, Jayasinghe CY. Plant poisoning among children in rural Sri Lanka. *Int J Pediatr* 2017;2017:6187487.
2. Kumaravel KS, Rameshbabu B. Kerosene poisoning in childhood: A 3-year retrospective study at a tertiary referral hospital. *Int J Contemp Med Res* 2016;3:1832.
3. Sunilkumar MN, Parvathy VK. An analysis of profile of childhood kerosene poisoning in a tertiary care medical college hospital. *Int J Pediatr Res* 2016;3:214.
4. Brennan P, Croft P. Interpreting the results of observational research: Chance is not such a fine thing. *BMJ* 1994;309:727-30. doi: 10.1136/bmj.309.6956.727, PMID 7950528
5. Lifshitz M, Sofer S, Gorodischer R. Hydrocarbon poisoning in children: A 5-year retrospective study. *Wilderness Environ Med* 2003;14:78-82. doi: 10.1580/1080-6032(2003)014[0078:hpicay]2.0.co;2, PMID 12825880
6. Hamid MH, Butt T, Baloch GR, Maqbool S. Acute poisoning in children. *J Coll Physicians Surg Pak* 2005;15:805-8. doi: 12.2005/JCPSP.805808, PMID 16398977
7. Thalhammer GH, Eber E, Zach MS. Pneumonics and pneumatoceles following accidental hydrocarbon aspiration in children. *Wien Klin Wochenschr* 2005;117:150-3.
8. Annobil SH, Ogunbiyi OA. Pulmonary radiological changes in kerosene poisoning in the Asir region of Saudi Arabia. *Ann Trop Paediatr* 1991;11:391-5. doi: 10.1080/02724936.1991.11747536, PMID 1721801
9. Gupta P, Singh RP, Murali MV, Bhargava SK, Sharma P. Kerosene oil poisoning--a childhood menace. *Indian Pediatr* 1992;29:979-84. PMID 1459719