

TITLE-EVALUATION OF THE EFFICACY OF PRANAYAMA ON THE LUNGS PARAMETER AND BLOOD PRESSURE OF HYPERTENSIVE PATIENTS.

AJIT PAL SINGH CHAUHAN¹, MANOJ KUMAR GUPTA², BABITA SHARMA³, DINESH SINGH GAUR⁴, AJAY KUMAR¹

Reader Department of Sharir Kriya Vijnana, Government Ashtang Ayurvedic College and Hospital, Lokmanya Nagar, Indore, Madhya Pradesh, India. 2-Lecturer Department of Roga and vikriti Vijnana, Government Ashtang Ayurvedic College and Hospital, Lokmanya Nagar, Indore, Madhya Pradesh, India. 3-Lecturer, Department of Sharir Kriya vijnana, Government Ashtang Ayurvedic College and Hospital, Lokmanya Nagar, Indore, adhyapradesh, India. 4-Lecturer, Department of Shalaky, Government Ashtang Ayurvedic College and Hospital, Lokmanya Nagar, Indore, Madhya Pradesh, India. 5- Lecturer, Department of Shalya, Government Ayurvedic College and Hospital, Gurukul Kangni Haridwar Uttarakhand, India. 6-Reader, Department of Kaya Chikitsa, Gaur Brahman Ayurvedic College and Hospital, Rohatak, Haryana, India. Email: drajitschauhan@gmail.com

Received: 7 July 2013, Revised and Accepted: 8 August 2013

ABSTRACT

Yoga is a science which has been practised in India from over thousands of years. Besides the spiritual achievements, the practice of yoga is accompanied by a number of beneficial physiological effects in the body. Yoga and health goes hand in hand. Yoga calms and relaxes the mind and it strengthens and tunes the body and brings them into harmony with each another. Pranayama is an art of controlling the life force of breath [1]. It produces many systemic psycho-physical effects in the body, besides its specific effects on the respiratory functions. This study is designed to study the effects of pranayama (12 weeks) on the pulmonary function parameters. It is one of the best lifestyle modifications which have ever been devised in the history of mankind. There are many classical paths which have been described to reach the ultimate goal of healthy life. It is an ancient yoga technique, a spiritual and physical practice which integrates the mind and body. Pranayama is a type of yogic practice which produces many systemic psycho-physical effects in the body, besides its specific effects on the respiratory functions. So, it has become a standard fare at health clubs and community recreation programmes.

Pulmonary function tests (PFTs) are simple screening procedures which are performed by using a standardized equipment (spirometer) to measure the lung function. This test provides useful information about the minimum levels of the lung function. The breath holding time measures the level of the threshold of the respiratory center to the partial pressure of the carbon dioxide (Pco₂) level.

Pulmonary function tests have been studied in yoga and pranayama practitioners. They have shown that the regular practice of these pranayama techniques have proved to be beneficial for the human body. Pranayama has a favourable influence and it causes a marked improvement in the lung functions. This study is designed to study the effects of pranayama (12 weeks) on the pulmonary function parameters.

In these clinical trial 60 patients of mild to moderate hypertensive patients were randomly selected and divided in to two groups. In each group 30 patients had taken. In Group A Pranayam had been given to the 30 patients of mild to moderate hypertensive patients and In Group B only placebo drug Prana had given to the 30 patients mild to moderate hypertensive patients. In all these two groups, group A was found as the most benefited because it showed significant changes in symptoms and Biochemistry investigations. Whereas in Group B (Placebo group) showed insignificant changes in symptoms as well as in Biochemistry investigations.

Keywords: For Yoga- Pranayama, Yoga, Pulmonary Function Parameters FVC FEV1 FEF (25-75%), and BHT For Hypertension-Vyan vikriti, vyan bala, Raktavritta vata, Raktagata vata, Dhamani Praticaya, Siragata vata, Rasabhara, Dhamani Prapurnata, Vyanavritta vata etc.

INTRODUCTION

Aims and objectives

To evaluate the efficacy of Pranayam on different parameter of respirometer and effect on blood pressure and lungs function.

To overcome the harmful effects of Allopathic drug in those patient which are suffering from Cardio-pulmonary problem like Hypertension, Asthma from long duration.

Materials and methods

The study group consisted of 60 patients of mild to moderate hypertensive patients (32 males and 28 females) were randomly selected and divided in to two groups. who were newly recruited for yoga training at the Lokmanya nagar Center, Indore, Informed consent was taken from all the subjects who volunteered (60 patients of mild to moderate hypertensive patients) for the study. They were motivated to undergo pranava, Nadishuddi and Savitri Pranayama training for 1 hour daily, for 6 days a week. The first phase of the recording of the pulmonary parameters was done at the beginning of their course. The second phase of the recording was done after 12 weeks of the regular pranayama practice. The practice of pranayama was for 1 hour a day in the morning (6.30 am to 7.30 am), for six days per week. All the subjects were compulsorily asked to have a balanced vegetarian diet. They initially performed stretching exercises for 10 mins before starting the pranayama. The

subjects sat in Padmasana. The left arm was held straight and it was placed on the left knee. All the three types of Pranayama i.e. Pranava, Nadi shuddi and Savitri Pranayama were done one after the other. Each one was done for 10 rounds. The study group consisted of 60 young adults (32 males and 28 females) who were newly recruited for yoga training at the lokmanya nagar indore. They were motivated to undergo pranava, Nadishuddi and Savitri Pranayama training for 1 hour daily, for 6 days a week. The first phase of the recording of the pulmonary parameters was done at the beginning of their course. The second phase of the recording was done after 12 weeks of the regular pranayama practice.

INCLUSION CRITERIA

Total 60 mild to moderate hypertensive patients were randomly registered for trial for pranayama, respirometry and Placebo drug. Who were aged between 18-60 years mild to moderate hypertensive patients?

A total 60 mild to moderate hypertensive patients was screened out from local regional area of Lokmanya nagar registered for trial for respirometry and Placebo drug.

Grouping of Patient

Screened patient or the case registered for the study were randomly divided into two groups

Group A: This group of 30 patients had given the trial of pranayam and respirometry.

Group B: This group of 30 patients had given the trial of placebo drug Prana 1gram B.D. per day with Luke warm water.

EXCLUSION CRITERIA

Patients with severe grade of the Hypertension. Moderate Hypertension associated with other diseases like Cardiomyopathy, Cardiac failure, Coronary artery disease, Heart block, Cerebrovascular disease, Encephalopathy, Preclampsia/eclampsia, Renal disease, Diabetes mellitus, Retinopathy. The subjects with a history of Allergic disorders or respiratory disorders. Smoking. Treatment with beta-agonists or the xanthenes group of drugs. Chest deformities like kyphosis and scoliosis.

Systemic diseases and respiratory disorders were ruled out in the selected subjects by taking their detailed history and by their thorough clinical examination.

Measurement of Blood Pressure

To obtain diagnosis: On each occasion at least 2 sets of blood pressure reading, separated by 20-30 minutes intervals was taken.

On the basis of 6th and 7th joint national committee on detection evaluation & treatment of high blood pressure.

JNC 6 Category	SBP / DBP in mm Hg	JNC 7 Category
Optimal	< 120 / 80	→ Normal
Normal	120 – 129 / 80 – 84	→ Pre hypertension
Border line	130 – 139 / 85-89	
Hypertension	≥ 140 / 90	Hypertension
Stage 1 (mild)	140 – 159 / 90 – 99	→ Stage I
Stage 2 (moderate)	160 – 179 / 100 – 109	
Stage 3	≥ 180 / 110	

Subjective (clinical) Parameters

Following symptoms have been screened for diagnosis of Essential Hypertension. Sirahashoola (Headache), Bhrama (Dizziness), Epistaxis and Heaviness of head.

Clinical Assessment

Assessment of Symptoms:

Symptoms of the disease were assessed before and after the treatment on the basis of following criteria.

Not Present/Absence of Symptom	0	0
Very mild	1	25%
Mild	2	50%
Moderate	3	75%
Severe	4	100%

Assessment of Blood Pressure Reduction

The results of the treatment were assessed as striking, wonderful, nice and fair at the end of treatment. The parameters of the assessment were taken as follows: -

Striking: An excellent response to therapy when the fall in D.B.P. was found >20 mm Hg or more, S.B.P. >40 mm Hg.

Wonderful: When the patient was noticed with a good response to therapy when fall in D.B.P. was found 11-20 mm Hg. S.B.P. was 21-40 mm. Hg.

Nice: The response is named Nice when the fall in D.B.P. was 6 -10 m; S.B.P. was 11-20 mm. Hg.

Fair: When the response falls in D.B.P. up to 5 mm Hg, S.B.P. was up to 10 mm Hg.

Anthropometric measurements which were recorded

Age was calculated in years to the nearest birthday.

Height was measured in centimetres while standing. The reading was taken nearest to ½ cm.

Weight was recorded in kilograms; the reading was taken nearest to ½ kg and the weighing machine was appropriately calibrated from time to time.

The pulmonary function tests were determined by using a Medspiror model (computerised spirometry) and the breath holding time was measured by using a stop clock

Pulmonary function tests

The pulmonary functions were tested by using the instrument, 'Medspiror' (a self calibrating computerized spirometer that fulfils the criteria for standardized lung function tests), which was available in the Department of Physiology, SDUMC. The pulmonary functions were tested at the start of the course and after 6 weeks. The parameters which were studied were,

- Forced vital capacity (FVC)
- Forced expiratory volumes (FEV1)
- Peak expiratory flow rate (PEFR)
- Forced expiratory flow 25-75 % (FEF 25-75%)

The procedure

The subjects were familiarized with the set up and detailed instructions and demonstrations were given for our satisfaction. The subjects were made to breathe out forcefully, following deep inspiration, into the mouthpiece which was attached to the pneumatachometer. The expiration was maintained for a minimum period of 3-4 seconds. 3 to 4 trials of maximal inspiratory and expiratory efforts were made and only the highest reading was taken for data processing.

As recommended by Snowbind's workshop, all the readings were taken with the subjects in the standing position [2]. All the tests were carried out at the same time of the day, between 8.30 am to 9.30 am, to avoid the possible variations, because rhythmic changes in the physiological functions were found to be associated with changes in the performance during this period [3].

The tests were done in a quiet room in order to alleviate the emotional and psychological stresses. During the tests, a maximum effort from the subjects was ensured by adequately motivating them to perform at their optimum level. A normal PEFr value of 3-5 L/sec ensured a maximum effort by the subject while performing the test.

BREATH HOLDING TIME RECORDING

Procedure: The subject was asked to hold his/her breath while in the sitting posture. The breath was maintained until the subject could no longer hold the breath voluntarily and the time was noted by using a stop watch. This recorded the BHT.

RESULTS AND ANALYSIS

An evaluation of a non-controlled study with 60 adult subjects was undertaken to study the effect of 12 weeks of pranayama on the lung parameters. The mean age of the study group was 25.76 ± 5.51 years (mean ± SD). A P value of <0.005 was considered significant. The

first phase of the recordings in the study group was taken at the beginning of their course. i.e., before starting the Pranayama practice and the second phase of the recordings was taken after 12 weeks of regular practice of Pranayama. Statistical software: The Statistical software, namely SAS 9.0, SPSS 15.0, Stata 8.0, MedCalc 9.0.1 and Systat 11.0 were used for the analysis of the data and Microsoft word and Excel were used to generate graphs, tables, etc [8, 9, 10, and 11].

Table 1 : Comparison of effect of 6 weeks pranayama on lung parameters.

Variables	Before	After	P value	Effect size
FVC (L/min)	2.66±0.406	3.25±0.48	<0.001**	1.48
FEV1 (L/min)	2.40±0.40	3.00±0.46	<0.001**	1.58
PEFR(L/sec)	6.14±1.08	7.43±1.17	<0.001**	1.24
FEF (25-75%)(L/min)	2.97±0.51	3.78±0.49	<0.001**	1.79
BHT(sec)	38.38±4.38	56.66±9.05	<0.001**	2.79





Fig-1

OBSERVATION AND RESULTS

In these clinical trial 60 patients of hypertension were randomly selected and divided in to two groups. In each group 30 patients had taken. In Group A Pranayam, In Group B placebo drug had given to the patients. Patients with persistent rise of blood pressure with clinical picture of Essential Hypertension have been selected for the research works and result of these 60 patients are shown in the following tables:


parameter	BT	AT	MD	N	MP	SD	SE	t	P	RESULT
Vital capacity	3670.1	3953.4	283.3	60	7.72	425.96	77.76	3.64	P<.001	H.S
Tidal volume	394.00	493.69	99.67	60	25.3	234.7	42.85	2.32	P<.01	S
Expiratory reserve volume	996.00	1106.7	110.00	60	11.11	201.34	36.76	3.01	P<.01	S
Pulse rate	85.33	78.5	6.83	60	8.00	12.81	2.33	2.92	P<.01	S
BP(SBP)	122.73	113.33	9.4	60	7.65	18.16	3.31	2.83	P<.01	S
BP(DBP)	84.33	78.6	5.73	60	6.79	12.30	2.24	2.55	P<.01	S



Total 60 people were randomly registered who the symptoms of Hypertension.

The Effect of Pranayam on 60 hypertensive patients on subjective parameter is given below.(table no-2)

No	parameter	BT	AT	MD	N	MP	SD	SE	t	P	Results
1	Headache	3.25	2.40	.85	60	26.15	1.59	.357	2.37	<0.02	S
2	Dizziness	3.10	2.20	.90	60	29.03	1.65	.369	2.43	<0.02	S
3	Epistaxis	2.59	1.59	1.00	60	38.61	1.46	.378	3.20	<0.01	S
4	Heaviness of Head	2.56	1.20	1.36	60	53.12	1.22	.244	5.57	<0.001	HS



DISCUSSION

Pranayama is a type of yogic breathing exercise. It is a form of physiological stimulation. The regular practice of Pranayama is a form of adaptation to a repeated stimulus. Breathing is the only autonomic function that can be consciously controlled and it is the key in bringing the sympathetic and the parasympathetic nervous system into harmony [12]. Breath is the only function through which we can influence the involuntary nervous system, i.e. we can establish rhythms of breathing with our voluntary nerves and muscles, which will affect the involuntary nervous system [13].

The pulmonary function test (PFTs) is a valuable tool for evaluating the respiratory system. It is a simple screening procedure which can be performed by using standardized equipment to measure the lung function [14].

Pulmonary function tests have been studied in yoga and pranayama practitioners. They have shown that the regular practice of these pranayama techniques have proved to be beneficial for the human body and that it also improved the breath holding time [15,16]. This study is designed to know the effect of pranayama (12 weeks) techniques on the pulmonary function parameters.

In our study, the PFT values were recorded in young healthy subjects before and after 6 weeks of pranayama. The pulmonary function parameters FVC, FEV1, PEFR, FEF25-75% and Breath holding time (BHT) were found to be significantly increased. These results were consistent with those of other studies which were done by Yadav A et al. [17], Upadhyay KD et al. [18] and Chanavirut et al. [19].

All these studies have explained that, during pranayama training, regular inspiration and expiration for prolonged period leads the lungs to inflate and deflate maximally and that it causes strengthening and increased endurance of the respiratory muscles [4,5,6,7]. This maximum inflation and deflation is an important physiological stimulus for the release of surfactants and prostaglandins into the alveolar spaces, which thereby increase the lung compliance [20, 21]. The stretch receptors reflex decrease the tracheobronchial smooth muscle tone activity, which leads to decreased air flow resistance and increased airway caliber, which causes the dynamic parameters of the lung function test to improve.

A study showed that after 4 weeks of the pranayama practice, the FVC, FEV1, FEF 25-75% and the PEFR values had improved in the subjects. In our study, a greater improvement of the pulmonary parameters was observed. This may be because our subjects were young, healthy adults (mean age group -25.76± 5.51 years) who had practised pranayama for 12 weeks [22].

A study by Bhargava MR et al. showed a statistically significant increased breath holding time after the pranayama practice. The same study explained that during pranayama training, regular inspiration and expiration for longer duration would lead to acclimatization of central and peripheral chemoreceptors for both hypercapnoea and hypoxia [23]. Acclimatization of the stretch receptors of the chest, the bronchial walls and the alveoli increase the synchronization between the lung tissue and the cortex. The prolonged inhalation in pranayama leads to an increased breath holding time [24, 25].

A study by Upadhyay A et al. in which the pranayama practice was conducted for duration of 4 weeks, showed increased PEFR, whereas

in our study, with 6 weeks of pranayama practice, the PEFR values showed greater improvement [18]. This showed that as the duration of the pranayama increased, the pulmonary function test parameters also increased proportionately.

It was observed in another study by Bhavani et al. that pranayama produced immediate and significant reduction in the auditory and visual reaction time, indicating an improved sensory motor performance and an enhanced processing ability of the central nervous system [26].

A study by Ravindra et al. on patients with premature ventricular complexes (PVC) and episodes of palpitations, found that pranayama produced an immediate relief in the palpitations and PVC. This improvement could have been because of the reduction 29 of the sympathetic reactivity which was attained by the pranayama training [27].

CONCLUSION

After completion of my study I have found the significant changes in investigation like Systolic BP & Diastolic BP ($p < 0.01$). I have found the significant changes in symptoms like sirah shoola ($p < 0.02$) dizziness ($p < 0.02$) Epistaxis ($p < 0.01$) and Heaviness of head ($p < 0.001$). After completion of my study I have also found the significant changes in investigation like Tidal volume ($p < 0.01$) Expiratory reserve volume ($p < 0.01$) and pulse rate ($p < 0.01$). Pranayama is an ancient yoga technique. The regular practice of Pranayama integrates the mind and the body. It differs from other forms of exercises as it mainly focuses on the sensations in the body. Pranayama thus acts directly on the various functions of the body and affords benefits in a positive way. Our study showed that the pulmonary function test values improved after (12 weeks) pranayama practice. The following may be the reasons for this:

Regular, slow and forceful inspiration and expiration for a longer duration during the pranayama practice, leading to strengthening of the respiratory muscles. Pranayama training causes improvement in the expiratory power and decreases the resistance to the air flow in the lungs.

Pranayama training causes an increase in the voluntary breath holding time. This may be due to acclimatization of the chemoreceptors to hypercapnoea.

Pranayama is a type of yogic breathing exercise. This resultant effect of pranayama can be used as lung strengthening tool to treat many lung diseases like asthma, allergic bronchitis, post pneumonia recoveries, tuberculosis and many occupational diseases. Breathing is an autonomic function that can be consciously controlled and it is the key in bringing the sympathetic and the parasympathetic nervous system into harmony. The multi system benefits of pranayama can be used to treat multisystem disorders like diabetes and hypertension. Can be performed by using standardized equipment to measure the lung function [14].

Pulmonary function tests have been studied in yoga and pranayama practitioners. They have shown that the regular practice of these pranayama techniques have proved to be beneficial for the human body and that it also improved the breath holding time [15,16]. This study is designed to know the effect of pranayama (12 weeks) techniques on the pulmonary function parameters.

In our study, the PFT values were recorded in young healthy subjects before and after 6 weeks of pranayama. The pulmonary function parameters FVC, FEV1, PEFR, FEF25-75% and Breath holding time (BHT) were found to be significantly increased. These results were consistent with those of other studies which were done by Yadav A et al. [17], Upadhyay KD et al. [18] and Chanavirut et al. [19].

All these studies have explained that, during pranayama training, regular inspiration and expiration for prolonged period leads the lungs to inflate and deflate maximally and that it causes strengthening and increased endurance of the respiratory muscles [4, 5, 6, and 7]. This maximum inflation and deflation is an important physiological stimulus for the release of surfactants and prostaglandins into the alveolar spaces, which thereby increase the

lung compliance [20, 21]. The stretch receptors reflexly decrease the tracheobronchial smooth muscle tone activity, which leads to decreased air flow resistance and increased airway caliber, which causes the dynamic parameters of the lung function test to improve.

A study showed that after 4 weeks of the pranayama practice, the FVC, FEV1, FEF 25-75% and the PEFR values had improved in the subjects. In our study, a greater improvement of the pulmonary parameters was observed. This may be because our subjects were young, healthy adults (mean age group -25.76 ± 5.51 years) who had practised pranayama for 12 weeks [22].

A study by Bhargava MR et al. showed a statistically significant increased breath holding time after the pranayama practice. The same study explained that during pranayama training, regular inspiration and expiration for longer duration would lead to acclimatization of central and peripheral chemoreceptors for both hypercapnoea and hypoxia [23]. Acclimatization of the stretch receptors of the chest, the bronchial walls and the alveoli increase the synchronization between the lung tissue and the cortex. The prolonged inhalation in pranayama leads to an increased breath holding time [24, 25].

A study by Upadhyay A et al. in which the pranayama practice was conducted for duration of 4 weeks, showed increased PEFR, whereas in our study, with 12 weeks of pranayama practice, the PEFR values showed greater improvement [18]. This showed that as the duration of the pranayama increased, the pulmonary function test parameters also increased proportionately.

It was observed in another study by Bhavani et al. that pranayama produced immediate and significant reduction in the auditory and visual reaction time, indicating an improved sensory motor performance and an enhanced processing ability of the central nervous system [26].

A study by Ravindra et al. on patients with premature ventricular complexes (PVC) and episodes of palpitations, found that pranayama produced an immediate relief in the palpitations and PVC. This improvement could have been because of the reduction

REFERENCES

- Mishra SP. Yoga and Ayurveda: Their alliedness and scope as positive health sciences. 2nd ed. Varanasi, Chaukhambha Sanskrit Sansthan 1997.
- Mauch AD. Effects of a two week yoga program on the pulmonary function. *BIO* 493.2008; 1-9.
- Madanmohan, Lakshmi J, Udupa K, Bhavanani AB. Effect of yoga training on handgrip, respiratory pressures and pulmonary function. *Indian J Physiol Pharmacol* 2003;47(4):387-92.
- Madanmohan, Udupa K, Bhavanani AB, Shatapathy CC, Sahai A. *Indian J Physiol Pharmacol*. 2004; 48(4): 461-65.
- Madanmohan, Lakshmi J, Udupa K, Bhavanani AB. Effect of yoga training on handgrip, respiratory pressures and pulmonary function. *Indian J Physiol Pharmacol*. 2003; 47(4): 387-92.
- Raghuraj P, Telles S. Muscle power, dexterity skill and visual perception in community home girls who were trained in yoga or sports and in regular school girls. *Indian J Physiol Pharmacol*. 1997; 41: 409-15.
- Dash M, Telles S. Yoga training and motor speed, based on a finger tapping task. *Indian J Physiol Pharmacol*. 1999; 43: 458-62
- Bharagava MG, Gogate, Mascarenhas JF. Autonomic responses to breath holding and its variations following pranayama. *Indian J Physiol Pharmacol*. 1998; 32(4): 257-63.
- Bernard R. *Fundamentals of Biostatistics*, 2000; 5th Edition, Duxbury, 80-240.
- Venkataswamyreddy M. *Statistics for Mental Health Care Research*, NIMHANS publication, India, 108-144.
- Sunderaoo PSS. Richard J. *An Introduction to Biostatistics, A manual for students in health sciences*, New Delhi: Prentice Hall of India. 86-160.
- Grover P, Varma VD, Pershad D, Verma SK. Role of yoga in the treatment of psychoneuron's bull. *PGI*. 1998; 22(2): 68-76.
- Nidhi Jain, Srivastava RD, Singhal A. The effect of the right and left nostril breathing on the cardiorespiratory and the autonomic parameters. *Indian J Physiol Pharmacol*. 2005; 49(4): 469-74.
- Udupa K, Madanmohan, Bhavani AB, Vijayalakshmi P, Krishnamurthy N. Effect of the pranayama training on the cardiac function in normal young volunteers. *Indian J Physiol Pharmacol*. 2003; 47(1): 27-33.
- Joshi LN, Joshi VD and Gokhale LV. Effect of pranayama Shankarappa V. et al., The Effect of Pranayama on the Lung Parameters www.jcdr.net *Journal of Clinical and Diagnostic Research*. 2012 February, Vol-6(1): 27-30 30 practice on the breathing rate and the ventilatory functions of the lung. *Indian J Physiol Pharmacol*. 1992; 36(2): 105-08.
- Makwana K, Khirwadkar N and Gupta H C. Effect of yoga practice on the ventilatory function tests. *Indian J Physiol Pharmacol*. 1988; 32 (3): 203-08.
- Yadav A, Savita S, Singh KP. Role of the pranayama breathing exercises in the rehabilitation of coronary artery disease patients. *Indian J of Traditional Knowledge*. 2009; 3:455-508.
- Upadhyay KD, Malhotra V, Sarkar D, Prajapati R. Effects of alternate nostril breathing exercises on the cardio respiratory functions. *Nepal medical Coll J* 2008; 10(1): 25-27.
- Chanavirut R, Khaidjapho K, Jare P, Pongnaratorn P. Yoga exercise increases chest wall expansion and lung volumes in healthy Thais. *Thai J Physiological sciences*. 2006; 19(1):1-7.
- Hilderbran JN, Georke J, Clements JA. Surfactant release exercised rat lung which is stimulated by air inflation. *J Applied physiol*. 1981; 51: 905-10.
- Smith AP. Prostaglandins and respiratory system prostaglandins; physiological, pharmacological and pathological aspects. Edited by SMM. Karim. 1976; 83-102.
- Yadav A, Savita S, Singh KP. Role of the pranayama breathing exercises in the rehabilitation of coronary artery disease patients. *Indian J of Traditional Knowledge*. 2009; 3:405-08.
- Joshi LN, Joshi VD. Effect of forced breathing on the ventilatory functions of the lung. *J Postgrad Med*. 1998; 44(3); 67-69.
- Bhargava MR, Gogate MG, Mascarenhas. A study of BHT and its variations following Pranayamic exercises. *The Clinician*. 1982; 43-46.
- Jerath R, Edry J, Barnes V, Jerath V. Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that can explain how slow deep breathing shifts the autonomic nervous system. *Medical hypo*. 2008; 67(3):566-571.
- Bhavanani et al. *Ind J Physiol Pharmacol*. 2003; 47:297-300.
- Ravindra et al. *International J Cardiology*. 2006; 108:124-25.