

PHYTOCHEMICAL EVALUATION OF *PUNICA GRANATUM* L. LEAF EXTRACT

P. SREEDEVI¹, K. VIJAYALAKSHMI², R. VENKATESWARI³

^{1,2}Department of Biochemistry, Bharathi Women's College, ³Department of Medical Biochemistry, University of Madras, Taramani Campus
Email: vijji42research@yahoo.co.in

Received: 27 Dec 2016, Revised and Accepted: 27 Mar 2017

ABSTRACT

Objective: This study was conducted to assess the phytochemical constituents in *Punica granatum* L. Leaf extracts (PGL) using standard methods.

Methods: The leaf powder was extracted using solvents namely aqueous, hydroalcohol, ethanol, ethyl acetate and n-hexane. Qualitative and Quantitative phytochemical screenings of PGL were assessed by standard methods.

Results: All the leaf extracts were positive for a wide range of bio-active compounds except n-hexane. The result has showed that the maximum amount of total phenols (394.16 mg/g DW of extract), total tannins (210.5 mg/g DW of extract), flavanoids (147.4 mg/g DW of extract) and total triterpenoids (112 mg/g DW of extract) were noted in ethanolic extract of *P. granatum* leaf (EPGL). The biological assay revealed that relevant amount of carbohydrate, protein, lipid and alkaloid in EPGL.

Conclusion: The findings of this study concluded that the EPGL had potential bioactive substances that may be used as pharmaceutical ingredients for formulation of new or prospective potent drug to cure wide range of metabolic diseases.

Keywords: *Punica granatum*, Phytochemical constituents, Total phenols, Tannins flavonoids and triterpenoids

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)
DOI: <http://dx.doi.org/10.22159/ijcpr.2017v9i4.1159>

INTRODUCTION

Medicinal plants play a major role in meeting the medicinal and health needs of about 70 % of populations in developed and developing countries, which serve as an important resource for the treatment of various maladies and illnesses [1].

Plants synthesize a wide range of chemical compounds which are classified based on their chemical class, biosynthetic origin and functional groups into primary and secondary metabolites. Primary metabolites directly involved in growth and development while secondary metabolites are not involved directly and they have been worked as biocatalysts. Primary metabolites are widely distributed in nature, occurring in one form or another in virtually all organisms. They are chlorophyll, amino acids, nucleotides, carbohydrates and so on, which have a key role in metabolic processes such as photosynthesis, respiration and nutrient assimilation. They are used as industrial raw materials and food additives. Secondary metabolites are synthesized during secondary metabolism of plants. They are the basic source for the establishment of several pharmaceutical industries since they have enormous medicinal properties [2]. The most important secondary metabolites are alkaloids, tannins, flavonoids, phlobatannins, saponins and cardiac glycosides. All secondary metabolites have specific function such as saponins have antifungal activity [3], some alkaloid may be useful against HIV infection [4], flavonoids have strong anticancer activity [5] and tannin have antimicrobial activity.

In the search for phytochemicals that may be of benefit to the pharmaceutical industry, researchers sometimes follow leads provided by local healers in a region [6]. Following such leads, plant parts are usually screened for phytochemicals that may be present. The presence of a phytochemical of interest may lead to its further isolation, purification and characterization. Then it can be used as the basis for a new pharmaceutical product. Successful determination of biologically active compounds from plant material is largely dependent on the type of solvent used in the extraction procedure [7]. This therefore underscores the need to try as much solvents as possible in screening plant parts for phytochemicals.

Punica granatum belongs to the family Punicaceae, commonly known as pomegranate, is a shrub or small tree with several upright, thorny

stems, the leaves are elliptic, roughly 2 inches, the flowers are white or red, double-flowered races, native of Asia and Mediterranean Europe [8]. It is also found in India and more arid regions of Southeast Asia, [9] the East Indies, and tropical Africa. For centuries, the barks, leaves, flowers, fruits and seeds of this plant have been used to ameliorate diseases [10]. The potential therapeutic properties of pomegranate are wide-ranging and include treatment and prevention of cancers, cardiovascular disease, diabetes, dental conditions, erectile dysfunction and prevention from ultra violet (UV) radiation. The pericarp of *P. granatum* is used to treat infections found in human sexual organs as well as mastitis, acne, folliculitis, piles, allergic dermatitis, tympanitis, scalds, diarrhoea and dysentery [11].

Considering all these facts, the present study was designed to investigate the presence of various phytochemicals in the different extracts of *Punica granatum* leaf, a plant which evokes various therapeutic effects.

MATERIALS AND METHODS

Preparation of leaf extracts

Leaves of pomegranate plant (*Punica granatum* L.) were obtained from around the Sathyavedu village, Andhra Pradesh, India. The *Punica granatum* leaf was authenticated by Dr. P. Jayaraman, Director of National Institute of Herbal science, Plant anatomy research centre, Chennai.

The leaves of the plant were carefully removed and thoroughly washed with distilled water to remove dust particles. They were dried in shade and finely powdered using an electric blender. Fifty grams of powdered material was subjected to Soxhlet extraction with 500 ml of n-Hexane, ethyl acetate, ethanol, hydroalcohol and water separately for 8 h. The extracts were evaporated to dryness under controlled temperature (35-40 °C). The extracts were stored in air tight containers under refrigeration. These dried extracts were dissolved in respective solvents and used for further analysis.

Qualitative Phytochemical screening: Phytochemical screening of *Punica granatum* Leaf extracts were assessed by standard methods (12-15).

Test for alkaloids

A fraction of extract was treated with 3-5drops of Wagner's reagent [1.27g of iodine and 2g of potassium iodide in 100 ml of water] and

formation of reddish brown precipitate (or colouration) indicates the presence of alkaloids.

Test for anthraquinone

To 1 ml of plant extract, few drops of 1% HCl were added. Appearance of red colour precipitate indicates the presence of anthraquinone.

Test for carbohydrates

1 ml of plant extract was mixed with alpha naphthol solution and then to the sides of the test tube conc. H₂SO₄ is added. Appearance of violet ring indicates the presence of carbohydrates.

Test for reducing sugar

1 ml of plant extract was mixed with few drops of Benedict's reagent and kept in boiling water bath, observed for the formation of reddish brown precipitate. A positive result shows the presence of reducing sugar.

Test for flavanoids

2 ml of extracts was treated with few drops of 20% sodium hydroxide solution. Formation of intense yellow colour, which becomes colourless on addition of dilute hydrochloric acid, indicates the presence of flavonoids.

Test for phenols

A portion of the extract was treated with aqueous 5% ferric chloride and formation of deep blue or black colour indicates the presence of phenols.

Test for proteins

To the extract, 1 ml of distilled water was added which was then heated with Biuret reagent and observed for the formation of violet/pink colour.

Test for free amino acids

The extract was heated with 0.2 % solution of Ninhydrin which result in the formation of purple colour, suggesting the presence of free amino acid.

Test for coumarins

To 2 ml of the test solution, a few drops of 10% NaOH were added. Appearance of yellow colour indicates the presence of coumarins.

Test for saponins

2 ml of extract was added to 6 ml of distilled water in a test tube. The mixture was shaken vigorously and observed for the formation of persistent foam that confirms the presence of Saponins.

Test for steroids

1 ml of extract was treated with few drops of chloroform, acetic anhydride and conc. H₂SO₄ and formation of dark pink or red colour indicated the presence of steroids.

Test for tannins

2 ml of extract was treated with 10% alcoholic ferric chloride solution and formation of blue or greenish colour solution indicated the presence of tannins.

Test for terpenoids

1 ml of chloroform was added to 2 ml of each extract followed by a few drops of concentrated sulphuric acid. A reddish brown precipitate produced immediately indicated the presence of terpenoids.

Quantitative phytochemical analysis

Estimation of total phenolic content (TP)

The total phenolic compound in the extract was determined by the Folin-Ciocalteu method. An aliquot of sample (0.2 ml) was mixed with 0.5 ml of Folin-Ciocalteu reagent (1: 2 dilution) and 4 ml of sodium carbonate (1M) and allowed to stand for 30 min at room

temperature. The absorbance was measured at 750 nm using a spectrophotometer (Beckman, DU 7400 USA). TP content in the extract was calculated and expressed as mg of Chlorogenic acid equivalent per g of dry weight of extract (mg CGE/g DW) [16].

Estimation of tannins

The tannin present in extracts of *P. granatum* was determined by the method of Peri and Pompei, 1971 [17]. 0.5 ml of extract was made upto 2 ml with distilled water and 2 ml of water serves as blank. To this 0.5 ml of Folin ciocalteu phenol reagent (1:2 dilution) followed by the addition of 5 ml of 35 % sodium carbonate and kept at room temperature for 5 min. Blue colour formed was read at 640 nm. A standard graph (Chlorogenic acid) was plotted, from which the tannin content of the extracts was determined. The total tannin content was expressed in mg CGE/g DW of extract.

Estimation of flavonoids

The TFC (total flavonoid content) of *P. granatum* leaf extract was determined using the aluminium chloride assay by colorimetry. An aliquot (0.5 ml) of extract was taken in different test tubes then 2 ml of distilled water was added followed by the addition of 0.15 ml of sodium nitrite (5% NaNO₂, W/V) and allowed to stand for 6 min, 0.15 ml of aluminium trichloride (10 % AlCl₃) was added and incubated for 6 min, followed by the addition of 2 ml of sodium hydroxide (NaOH, 4% W/V). The solution was well vortexed and absorbance was measured against reagent blank at 510 nm. The total flavonoid content (mg/g) was determined from the calibration curve of quercetin and expressed as mg quercetin equivalents (mg QE/g DW) [18, 19].

Estimation of total triterpenoids

Briefly 200 µl sample solution in a 10 ml volumetric flask was heated to evaporation in a water bath, 1 ml of 5 % (W/V) vanillin-acetic acid solution and 1.8 ml sulphuric acid were added, mixed and incubated at 70 °C for 30 min. Then the solution was cooled and diluted to 10 ml with acetic acid. The absorbance was measured at 573 nm against blank using spectrophotometer. The blank consisted of all reagents and solvents without sample solution. The triterpenoids was determined using the standard ursolic acid and expressed as milligram of ursolic acid equivalent/gram dry weight of extract (mg UE/g DW) [20].

Estimation of carbohydrates

0.2 ml of extract was made up to 1.0 ml with distilled water. 4.0 ml of anthrone reagent (0.2% anthrone in ice cold conc. Sulphuric acid) was added and kept in boiling water bath for 8 min, cooled rapidly and read at 630 nm. The blank consisted of all reagents without sample solution. D-glucose was used as standard. The total sugar content was expressed in terms of percentage of dry weight [21].

Estimation of proteins

0.2 ml of extract was made to 1.0 ml with distilled water. 5.0 ml of alkaline copper reagent was added to all the tubes and allowed to stand for 10 min. Then 0.5 ml of Folin's Ciocalteu reagent was added and incubated in dark for 30 min. The intensity of the colour developed was read at 660 nm. The blank consisted of all reagents without sample solution. The protein content was determined using the standard Bovine serum albumin and expressed in terms of percentage of dry weight [22].

Estimation of total lipids

10 gm. sample was used to extract lipids with 150 ml of petroleum ether for 16 hr., at a solvent condensation rate of 2-3 drops/sec. The obtained extract was concentrated and evaporated at room temperature to dryness. The weight of extract gives the total lipid content which was expressed in terms of percentage of dry weight [23].

Alkaloid determination by harborne (1973) method

5 g of the sample was weighed and added to 250 ml beaker and 200 ml of 10% acetic acid in ethanol was added and covered and allowed to stand for 4 h. This was filtered and the extract was concentrated

on a water bath to one-quarter of the original volume. Concentrated ammonium hydroxide was added drop wise to the extract until the precipitation was complete. The whole solution was allowed to settle and the precipitate was collected and washed with dilute ammonium hydroxide and then filtered. The residue is the alkaloid, which was dried, weighed and expressed in terms of percentage of dry weight [24].

RESULTS AND DISCUSSION

Preliminary phytochemical analysis

The preliminary phytochemical tests are helpful in finding chemical constituents in the plant material that may lead to their quantitative estimation and also in locating the source of pharmacologically active chemical compound.

Table 1: Qualitative phytochemical analysis

S. No.	Phytochemical constituent	Aqueous extract	Hydroalcohol extract	Ethanol extract	Ethyl acetate extract	n-hexane extract
1	Alkaloid	+	+	+	+	-
2	Carbohydrate	+	+	+	+	+
3	Reducing sugar	+	+	+	+	+
4	Flavonoid	+	+	+	+	-
5	Phenol	+	+	+	+	+
6	Protein	+	+	+	+	-
7	Amino acid (free)	-	-	-	-	-
8	Coumarin	+	+	+	+	-
9	Saponin	+	-	-	-	-
10	Tannin	+	+	+	+	+
11	Steroids	-	+	+	+	+
12	Terpenoids	+	+	+	+	-
13	Anthraquinone	-	-	-	-	-

Table 2: Quantification of secondary metabolites of punica granatum leaf extract

S. No	Extract	Total phenols (mg of CGE/g of dry extract)	Tannins (mg of CGE/g of dry extract)	Flavonoids (mg of QE/g of dry extract)	Total triterpenoids (mg of UE/g of dry extract)
1	Water Extract	274.1±0.65	90.9±0.26	77.26±0.8	44.9±1.55
2	Hydro alcohol Extract	117.6±0.76	73.6±0.52	66.3±1.25	71.73±1.67
3	Ethanol Extract	394.16±0.76	210.5±0.5	147.4±1.0	112±2.36
4	Ethyl acetate Extract	240.8±0.28	141.6±0.59	33.3±1.5	37.3±0.92

Values are expressed as mean±SD (n = 4).

Table 3: The alkaloid content and calorific value of ethanolic extract of punica granatum Leaf in percentage

Alkaloid	3.5±1.5
Carbohydrates	19.6±2.3
Proteins	11.86±0.9
Lipids	0.96±1.7

Values are expressed in percentage (n = 4)

The qualitative analysis of bioactive compounds for the five extracts have been analysed in this study and there is wide range of phytochemical compounds present in the five extracts as shown in table 1. The hexane being highly nonpolar in nature was able to extract very less compound characterized like carbohydrates, phenols, steroids and tannins. Ethanolic extract, ethyl acetate and hydro alcoholic extract was found to have a wide range of bioactive compounds like alkaloids, carbohydrates, coumarins, flavonoids, proteins, phenols, reducing sugars, steroids and tannins.

The aqueous extract found to have bioactive constituent said above in addition to saponins except steroids. The presence of bioactive constituents indicates that the *P. granatum* Leaf extract can be used in a multitude of ways for the beneficiary of population.

The results for quantitative analysis of the *P. granatum* leaf extract are illustrated in table 2. Ethanolic extract was found to possess the maximum amount of Phenols (394.16 mg of CGE/g of DW), Tannins (210.5 mg CGE/g of DW), Flavonoids (147.4 mg of QE/g of DW) and total triterpenoids (112 mg of UE/g of DW). Since the yield of bioactive metabolites in a plant extract also varies considerably with the (25, 26) method/solvent of extraction it is plausible that the ethanolic extracts were generally more potent than the aqueous

extracts probably because the active principles in the plant dissolved more readily in and were better extracted by a less polar solvent (ethanol) than water. This is in agreement with many literatures reporting the differences in the activities of extracts obtained from the same morphological part of a plant using different solvents. The methanolic extract of the fruits of *Tetrapleura tetraptera* is more potent than the aqueous [27] extract. Next to ethanolic extract, the water extract have higher content of total phenols (274.1 mg of CGE/g of DW) and flavonoids (77.26 mg of QE/g of DW) except tannins (90.9 mg of CGE/g of DW) and total triterpenoids (44.9 mg of UE/g of DW). The quantity of total phenols (117.6 mg of CGE/g of DW) and tannins (73.6 mg of CGE/g of DW) were found to be lowest in hydroalcohol extract except flavonoids (66.3 mg of QE/g of DW) and total triterpenoids (71.73 mg of UE/g of DW). Considerable amount of total phenols (240.8 mg of CGE/g of DW) and tannins (141.6 mg of CGE/g of DW) existed in ethyl acetate but found to have lowest amount of flavonoids (33.3 mg of QE/g of DW) and total triterpenoids (37.3 mg of UE/g of DW). These results were in agreement with Pande and Akoh [28]. They found that the total polyphenols was 365 mg GAE/g of FW in *P. granatum* leaves. Similar results were given by Hemayet Hossain *et al.*, (29), total phenolic content given as 378.37±0.92 mg of gallic acid/g of dry extract.

The medicinal value of plants lies in some chemical substances that have a definite physiological action on the human body. Different phytochemicals have been found to possess a wide range of activities, which may help in protection against chronic diseases [30].

Flavonoids have been used against the cancer causing tumours and it inhibits the promotion of growth and progression of tumours [31]. Phenols when mixed with the flavonoids compounds in plants are reported to show multiple activities like antioxidant, anti-carcinogenic, anti-inflammatory etc. [32]. Plants with tannins are used for healing of wounds, varicose ulcers, haemorrhoids, frost-bite and burns [33, 34]. Terpenoids are reported to have anti-inflammatory, anti-viral, anti-malarial, inhibition of cholesterol synthesis and anti-bacterial activity [35, 36].

The Alkaloid and Calorific determination of EPGL reveals the biological value of *Punica granatum* Leaves (table 3). Carbohydrate content was found high (19.6 % of dry matter) followed by protein (11.86 % of dry matter) and alkaloid content (3.5% of dry matter) whereas lipid content was found very low i.e., 0.96% of dry matter. The presence of higher protein level in the plant parts shows their possible increased food value or that a protein based bioactive compound could also be isolated in future. *Punica granatum* leaves have a low level of lipid is an indication that it would have little cholesterol. Alkaloids which are one of the largest groups of phytochemicals in plants have amazing effects on humans and this has led to the development of powerful pain killer medications [37]. Also, studies have shown that alkaloid is capable of reducing headaches associated with hypertension [38].

CONCLUSION

Nowadays herbs are extensively used for the research purpose and it possesses more than one chemical entity so it has been widely used for the research investigations. The plant based compounds have the effective dosage response and minimal side effects when compared to the synthetic compounds. Phytochemical screening of *Punica granatum* leaves reveals it as a valuable medicinal plant with numerous medicinal properties. Since the ethanolic extract of *P. granatum* leaves contains more constituents it can be considered beneficial for further investigation. A typical research and developmental work needs to be carried out for its better therapeutic and commercial utilization.

CONFLICT OF INTERESTS

Declare none

REFERENCES

- Ngari EW, Chiuri LW, Kariuki ST, Hockett S. Ethnomedicine of ogiek of river njoro watershed. *Ethnobot Res Appl* 2010;8:135-52.
- Lingarao M, Savithamma N. Phytochemical studies of *svensonia hyderabadensis* (Walp.) Mold-arare medicinal plant: *Der Pharm Lett* 2011;3:51-5.
- Sodipo OA, MA Akanji, FB Kolawole, Odutuga AA. Saponin is the active antifungal principle in *Garcinia kola*, heckle seed. *Biosci Res Commun* 1991;3:171-1.
- Mc Mahon JB, Currens MJ, Gulakowski RJ, Buckheit RWJ, Lackman-Smith C, Hallock YF, *et al.* Michellamine B a novel plant alkaloid, inhibits human immunodeficiency virus-induced all killing by at least two distinct mechanisms. *Antimicrobial Agents Chemother* 1995;39:484-8.
- Noble RI. The discovery of Vinca alkaloids chemotherapeutic agents against cancer: *Biochem. Cell Biol* 1990;68:1544-51.
- Prasanth Tiwari, Bimlesh Kumar, Mandeep Kaur, Gurpreet Kaur, Harleen Kaur. Phytochemical screening and extraction: review. *Int Pharm Sci* 2011;1:98-106.
- Das K, Tiwari RKS, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agent: current methods and future trends. *J Med Plant Res* 2010;4:104-11.
- Egharevba HO, Kunle OF. Preliminary phytochemical and proximate analysis of the leaves of *Piliostigma thionningii* (Schumach.): milne-redhead. *Ethnobot Leaflets* 2010;14:570-7.
- Naqvi SA, Khan MS, Vohora SB. Antibacterial, antifungal, and antihelminthic investigations on Indian medicinal plants: *Fitoterapia* 1991;62:221-8.
- Jayaprakasha GK, Negi PS, Jena BS. Antimicrobial activities of pomegranates, in pomegranates; ancient roots to modern medicines, Eds., CRC Press: Boca Raton, FL, USA; 2006. p. 167-8.
- Singh RP, Chidambara MKN, Jayaprakasha GK. Studies on the antioxidant activity of pomegranate (*Punica granatum*) peel and seed extracts using *in vitro* models: *J Agric Food Chem* 2002;50:81-6.
- Sofowara A. Screening plants for bioactive agents. In: *Medicinal plants and traditional medicinal in Africa*. 2nd edition. Spectrum Book Ltd, Sunshine House, Ibadan Nigeria; 1993. p. 134-56.
- Trease GE, Evans WC. *Pharmacognosy*. 15th ed. Saunders Publishers. London; 2002. p. 42-44, 22-229, 246-249, 304-306, 331-332, 391-393.
- Singleton VL, Rossi JA. Colorimetry of total phenolic with phosphomolybdic acid-phosphotungstic acid reagents. *Am J Enol Viticulture* 1965;16:144-58.
- Vishnoi NR. *Advanced practical chemistry ghaziabad-India*; Yikas Publication House, Pvt Ltd; 1979. p. 447-9.
- El-falleh W, Hannachi H, Tlili N, Yahia Y, Nasri N, Ferchichi A. Total phenolic contents and antioxidant activities of pomegranate peel, seed, leaf and flower. *J Med Plants Res* 2012;6:4724-30.
- Peri C, C Pompei CJ. Estimation of different phenolic groups in vegetable extracts. *Phytochemistry* 1971;19:2187-9.
- Samantha T, Shyamsundarachary R, Srinivas, P, Swamy NS. Quantification of total phenolic and total flavonoid contents in extracts of *oroxylum indicum* L. Kurtz. *Asian J Pharm Clin Res* 2012;5:177-9.
- Patel AN, Bandawane DD, Mhetre NK. Pomegranate (*Punica granatum* Linn.) leaves attenuate disturbed homeostasis and hyperglycemia mediated hyperlipidemia and oxidative stress in streptozotocin induced diabetic rats. *Eur J Integrative Med* 2014;6:307-21.
- Lu ZM, Gong Js, He Z, Xu HY, Dou WF, Shi JS, *et al.* Optimization of total triterpenoids from submergedly cultivated *Androia Camphorata* using response surface methodology. *Nat Prod Res Dev* 2011;23:946-51.
- Hedge JE, Hofreiter BT. In: *Methods in carbohydrates chemistry*. Vol. Eds. Whistler RL, BeMiller JN. Academic Press: New York; 1962. p. 17:420.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with folin phenols reagent. *J Biol Chem* 1957;93:265-75.
- Cheung PCK, Leung AYH, Ang PO. Comparison of supercritical carbon dioxide and Soxhlet extraction of lipids from a brown seaweed, *Sargassum hemiphyllum* (Turn.) C. *Ag J Agric Food Chem* 1998;46:4228-4232.
- Harborne JB. *Phytochemical methods*, London. Chapman and Hall, Ltd; 1973. p. 49-188.
- Clark TE, Appleton CC, Drewes SE. A semi-quantitative approach to the selection of appropriate candidate plant molluscicides—a South African application. *J Ethnopharmacol* 1997;56:1-13.
- Marston A, Maillard M, Hostettmann K. Search for antifungal, molluscicidal and larvicidal compounds from African medicinal plants. *J Ethnopharmacol* 1993;38:215-23.
- Adewunmi CO, Adesina SK, Marquis VO. On the laboratory and field evaluation of the molluscicidal properties of tetrapleura tetraptera. *Bull Anim Hlth Prod Afr* 1982;30:89-94.
- Pande G, Akoh CC. Antioxidant capacity and lipid characterization of six georgia-grown pomegranate cultivators. *J Agric Food chem* 2009;57:9427-36.
- Hemayat Hossain, Tanzir Ahmed, Sariful Islam Howlader, Shubhra kanti Dey, Arpana Hira, Arif ahmed, *et al.* *In vitro* antioxidant potential from the leaves of *punica granatum* Linn. Grown in Bangladesh. *Int J Pharm Phytopharmacol Res* 2012;2:160-6.
- Mir MA, Sawhney SS, Jassal MMS. Qualitative and quantitative analysis of phytochemicals of *taraxacum officinale*. *Wudpecker J Pharm Pharmacol* 2013;2:1-57.
- Stevens JF, Hart HT, Hendriks H, Malingre TM. Alkaloids of some European and macaronesian diode and semepervivodeae (Crassulaceae). *Phytochemistry* 1992;31:3917-24.
- Asha K, Rasika CT, Nirmala RD, Jyoti PS. Antioxidant potential from Stem Bark of *Juglans regia* L. *Ann Biol Res* 2011;2:176-80.

33. Igboko DO. Phytochemical studies on *Garcinia Kola* Heckel. M.Sc. Thesis. University of Nigeria, Nsukka; 1983. p. 202.
34. Maiduyi I. Biochemical and pharmacological studies of active principles of the seeds of *Garcinia kola* Heckel; MSc. Thesis, University of Nigeria, Nsukka; 1983. p. 108.
35. Guangyi Wang, Weiping Tang, Robert R, Bidigare. Terpenoids as therapeutic drugs and pharmaceutical agents: Natural Products; 2005. p. 197-227.
36. McGarvey DJ, Croteau R. Terpenoid metabolism. *Plant Cell* 1995;7:1015-26.
37. Kam PC, Liew A. Traditional Chinese herbal medicine and anaesthesia. *Anaesthesia* 2002;57:1083-9.
38. Ayitey-Smith E, Addae-Mensah I. Phytochemical, nutritional and medical properties of some leafy vegetables consumed by Edo people of Nigeria. *W Afr J Pharmacol Drug Res* 1977;4:7-8.

How to cite this article

- P Sreedevi, K Vijayalakshmi, R Venkateswari. Phytochemical evaluation of *Punica granatum* L. leaf extract. *Int J Curr Pharm Res* 2017;9(4):14-18.