

Original Article

**COMPARATIVE *IN VITRO* ANTICANCER ACTIVITY OF METHANOLIC EXTRACT OF TRADITIONALLY USED MEDICINAL PLANT OF MIZORAM**

TRIDEEP SAIKIA\*, BISWAJIT DAS, PURABI DAS

Girijanada Chowdhury Institute of Pharmaceutical Science, Zemabawk, Aizwal, Mizoram 796017

Email: tsaikia53@gmail.com

Received: 14 Apr 2019, Revised and Accepted: 13 Jun 2019

**ABSTRACT**

**Objective:** The main objective was to find out the *in vitro* comparative anticancer activity of various methanolic extract.

**Methods:** Plants samples such as *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum*, and *Alpinia galanga* were collected, identified and authenticated. By using Trypan blue test Human cervical cancer (Hela) cells were counted. The Human cervical cancer (Hela) cells were treated with methanolic extract of *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum* and *Alpinia galanga*. 20  $\mu$ l of MTT (5 mg/ml) solution was added to cells per well, and the plate was moved to a cell incubator. Measurement was performed using a Spectramax M2 Microplate Reader (Molecular Diagnostic, Inc.) at a wavelength of 570 nm.

**Results:** The Cytotoxicity of studied medicinal plants of Mizoram such as *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum* and *Alpinia galanga* exhibited cytotoxicity in an increased manner with increase concentration against Hela cells. Their IC<sub>50</sub> values were 49.02, 138.5, 199.7 and 209.4  $\mu$ g ml<sup>-1</sup> respectively. The IC<sub>50</sub> of doxorubicin was found to be 3.305  $\mu$ g ml<sup>-1</sup>. The anticancer activity of the leaves related to their content of flavanoids. This study validates the traditional use of plants in the management of cancer.

**Conclusion:** The Cytotoxicity of studied medicinal plants of Mizoram such as *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum* and *Alpinia galanga* exhibited cytotoxicity in an increased manner with increase concentration against HeLa cells.

**Keywords:** Trypan blue test, Cytotoxicity, Human cervical cancer cell (Hela cell), Anti-cancer

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)  
DOI: <http://dx.doi.org/10.22159/ijcpr.2019v11i4.34929>

**INTRODUCTION**

Cancer now a day is very widely spread and prevalent disease. Cancer is the third leading cause of death worldwide, preceded by cardiovascular and infectious diseases. It is becoming very common to every age group. It is very serious as its curability is very less due to unawareness for the symptoms and also the proper medication still not available. It is a major concern in the health care perspective [1]. Cervical cancer is one now becoming common in the northeastern region. These are the main perspective to get through this research work in regards of helping the society. Cancer is one of the most life-threatening diseases in which deregulated proliferation of abnormal cells invades and disrupts surrounding tissues. There has been the success of using clinical therapies such as radiation, chemotherapy, immunomodulation and surgery in the treatment of cancer but these are limited. So, there is a need of alternative strategies in the management of cancer disease. Natural products can play a significant role as secondary metabolites present in them such as terpenoids, phenolic acids, flavonoids, alkaloids, etc. exhibit antioxidant properties which is significant in the role of cancer treatment. Thus, medicinal plants have become a focal point to improve future health needs against cancer with a lesser harmful effect [2]. The methanol extract of the plant like *Mikania micrantha*, *Allium hookeri*, *Alpinia galanga*, *Eryngium foetidum* was also carried out for anticancer activity studies, which includes the study of cytotoxicity by an MTT assay and cell viability determination by Trypan blue exclusion assay. These two studies will give information on how the plant plays a role in fighting cancer cells. The main principle of cytotoxicity of cells by MTT Assay is that the yellow tetrazolium MTT (3-(4, 5-dimethyl thiazolyl-2)-2, 5-diphenyltetrazolium bromide) is reduced by metabolically active cells, in part by the action of dehydrogenase enzymes, to generate reducing equivalents such as NADH and NADPH. The resulting intracellular purple formazan can be solubilized and by spectrophotometric means [3].

**MATERIALS AND METHODS**

**Plant materials**

Plants samples such as *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum*, and *Alpinia galanga* were collected from Mizoram. The plant specimens were authenticated by Dr. A. A. Mao (Scientist-F and H) Botanical Survey of India, Shillong. A voucher specimen of these plants has been deposited in the Department of Pharmacy, with Reference no: BSI/ERC/TECH/2017/589 for future reference. Then samples were subjected for hot methanol extraction. The plant's extract was dried at room temperature until semi-solid or dryness is obtained. The extract was then dissolved in 0.01% methanol in distilled water and filtered through 0.22 $\mu$ . Doxorubicin was used as a positive control with different concentration of 1, 2, 4, 8  $\mu$ g/ml [5].

**Cell lines and culture medium**

A Hela (Human cervical cancer) cell line was procured from National Centre for Cell Sciences (NCCS), Pune, India. Stock cells were cultured in Dulbecco modified eagle media (DMEM) supplemented with 10% inactivated Fetal Bovine Serum (FBS), penicillin (100 IU/ml), streptomycin (100  $\mu$ g/ml) and amphotericin B (5  $\mu$ g/ml) in a humidified atmosphere of 5% CO<sub>2</sub> at 37 °C until confluent. The cells were dissociated with trypsin solution (0.2% trypsin, 0.02% EDTA, 0.05% glucose in PBS). The stock cultures were grown in 25 cm<sup>2</sup> culture flasks, and all experiments were carried out in 96 microtitre plates (Tarsons India Pvt. Ltd., India).

**Sterility test**

It was done to check the extract from the contamination. 35 mm culture disc was plated with Hela in 2 ml of DMEM media and allow the cell to adhere. The crude plant extract was added in the microtitre plate and incubate at CO<sub>2</sub> incubator (5%) for 24 h.

### Cell viability

The cytotoxic effects of the various plants were investigated using the MTT (Sigma, USA) on HeLa cells. The cells were seeded in 96-well plates at a density of  $2 \times 10^4$  cells per well. After incubation for 20–24 h, the cells with 70–80% confluency were treated with the extracts at different concentrations (5, 25, 50, 100 and 200  $\mu\text{g ml}^{-1}$ ) and incubated for 24 h. Then, 20  $\mu\text{l}$  of MTT (5 mg/ml) solution was added to cells per well, and the plate was moved to a cell incubator for another 4 h. The medium was removed, and 150  $\mu\text{l}$  of DMSO was added to the cells. The plate was gently shaken for 15 min to dissolve the formazan crystals generated by proliferating cells, and the measurement was performed using a Spectramax M2 Microplate Reader (Molecular Diagnostic, Inc.) at a wavelength of 570 nm. Relative viability was calculated taking wells with non-treated cells as 100% control. [6]

### Test for cellular proliferation (MTT assay protocol)

Seeded the HeLa at  $3 \times 10^5$  cells/well in a 96 well plate. Cells may be seeded at different densities. At least three wells were left without cells. These wells serve as a control for the minimum absorbance. The plate was incubated overnight at 37 °C in an a humidified incubator, 5% CO<sub>2</sub> for the cancer cells to grow and adhere to the surface. Test compounds were added to the plate. Include replicates for a range of concentrations. Include negative controls (including vehicle control) and a positive control. The final volume will be 100  $\mu\text{l}$  perwell. The plate was incubated for overnight (or for some other appropriate time) at 37 °C in a humidified incubator, 5% CO<sub>2</sub>. MTT reagent (20  $\mu\text{l}$ /100  $\mu\text{l}$  per well of the 96well plate) was added. Incubated at 37 °C for 4 h. 1 volume (100  $\mu\text{l}$ ) of the stop mix solution was added and the plate was rocked at room temperature for a minimum of 1 hour. (Allow time for the formazan precipitate to dissolve). The stop mix solution must be added in a fume hood. A

purple colour should be visible at this stage and should deepen over the 1-hour incubation period. After the 1-hour incubation, ensure the formazan precipitate is dissolved by pipetting each well up and down until no precipitate is visible. Read the plate on a plate reader using wavelength at 572 nm. Tabulate results and calculate the % viability [6, 7]

$$\% \text{ Viability} = \frac{\text{Mean absorbance of sample}}{\text{Mean absorbance of control}} \times 100$$

### Morphological staining

To observe the morphological changes of the cells an inverted phase contrast microscope was used. Cells were inoculated at  $3 \times 10^5$  cell/well in 24 well-micro plates and treated mentioned manner. Other culture wells were treated by H<sub>2</sub>O<sub>2</sub> (100  $\mu\text{M}$ ) as a positive control of apoptosis and necrosis, respectively, as negative control some culture wells were prepared without any treatment. After being cultured for 16h, the culture media removed and cells fixed and stained by the standard hematoxylin-eosin method. The prepared samples were photographed at  $\times 100$ .

### Statistical analysis

One-way analysis of variance (ANOVA), Duncan and Wilcoxon test were used for data analysis. All the results are expressed as the mean  $\pm$  SD, and p-values below 0.05 were considered statistically significant.

## RESULTS AND DISCUSSION

### In vitro anti-cancer activity (MTT assay)

The MTT assay results for 24 h and 48 h incubation of crude drug at the concentrations of 5, 25, 50, 100, and 200  $\mu\text{g/ml}$  showed increased HeLa cell viability as the concentration got diluted.

Table 1: The percentage cell viability of HeLa cell lines against plants extract, the data present as mean  $\pm$  SD, n=3

Conc. ( $\mu\text{g/ml}$ )	Doxorubicin		<i>Mikania micrantha</i>		<i>Allium hookeri</i>		<i>Alpinia galanga</i>		<i>Eryngium foetidum</i>	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
5(0)	0.00	0.00	10.83	1.09	5.08	1.40	6.20	1.07	-8.56	0.31
25(1)	6.07	1.61	25.07	1.17	19.94	2.00	15.19	1.08	2.50	0.52
50(2)	34.11	2.36	49.80	1.80	26.22	0.67	26.98	1.39	12.96	0.22
100(4)	50.00	3.93	78.79	0.25	49.50	1.66	40.01	2.39	37.06	2.74
200(8)	80.72	1.24	81.54	0.72	54.35	1.51	48.22	1.01	44.97	2.14
IC <sub>50</sub>	3.31		49.02		138.50		199.70		209.40	

\*The brackets indicate concentration of doxorubicin.





Fig. 1: The graph represents the percentage of cell cytotoxicity of extracts against, HeLa cells. The brackets indicate the concentration of doxorubicin


### Morphological staining

Morphological study of cell shape changes was performed by direct microscopy, hematoxylin and eosin staining, using an inverted phase-


contrast microscope (400X), it was found that the untreated cells exhibited normal shapes, with clear outline. Although the growth of the Methanolic-extract-treated cells was obviously inhibited. The extract treated cells were round, proliferation was inhibited and slowed.



After 48h of treated cells but before addition of MTT



Control Doxorubicin (4µg/ml) *Mikania Micrantha*




*Alpania galangal*

*Eryngium foetidum*

*Allium hookeri*

After 48h of Treated cells incubated with MTT for 3 h



*Alpania galangal*

*Eryngium foetidum*

Fig. 2: Represents various plants extracts treated HeLa with 100µg/ml. above lane 48hr treated and below represents 48 h treated followed by incubation with MTT. The purple crystal formation indicated viable cells. Lesser purple crystal formed, more cell dead

**CONCLUSION**

The Cytotoxicity of studied medicinal plants of Mizoram such as *Mikania micrantha*, *Allium hookeri*, *Eryngium foetidum* and *Alpinia galanga* exhibited cytotoxicity in an increased manner with increase concentration against HeLa cells. Their IC<sub>50</sub> values were 49.02, 138.5, 199.7 and 209.4 µg ml<sup>-1</sup> respectively. The IC<sub>50</sub> of doxorubicin was 3.305µg ml<sup>-1</sup>.

**ACKNOWLEDGMENT**

The author sincerely thanks Dr. Biswajit Das M. Pharm, Ph. D., Girijananda Chowdhury Institute of Pharmaceutical Science for his constant support and valuable suggestions in completing this manuscript.

**AUTHORS CONTRIBUTIONS**

All the author have contributed equally

**CONFLICT OF INTERESTS**

Declare none

**REFERENCES**

1. Kiranmayi Gali, G Ramakrishnan, R Kothai, B Jaykar. *In vitro* anticancer activity of methanolic extract of leaves of argemone mexicana. Int J PharmTech Res 2011;3:1329-33.
2. Tagne RS, Telefo BP, Nyemb JN, Yemele DM, Njina SN, Goka SMCG, *et al.* Anticancer and antioxidant activities of methanol extracts and fractions of some cameroonian medicinal plants. Asian Pacific J Trop Med 2014;7:442-7.
3. Mosamann T. Rapid colorimetric assay for cellular growth and survival, application to proliferation and cytotoxicity assay. J Immunol Methods 1983;65:55-63.
4. Sakinehsalehi, Saber Z, Seyed MR. Cytotoxic effect of *Prangos Pabularia* extracts on HELA cell line a medicinal plant. Int J Med Res Health Sci 2016;5:547-52.
5. Kostova I. Synthetic and natural coumarins as cytotoxic agents. Curr Med Chem Anti Cancer Agents 2005;5:29-46.
6. Plumb JA. Cell sensitivity assays: the MTT assay. Methods Mol Med 2004;88:165-9.
7. Denecker G, Vercammen D, Declercq W, Vandenabeele P. Apoptotic and necrotic cell death induced by death domain receptors. Cellular Mol Life Sci 2001;58:356-70.