

Original Article

## EVALUATION OF ANTI-PYRETIC AND WOUNDHEALING ACTIVITIES OF EVOLVULUS NUMMULARIUS LEAVES IN ALBINO WISTAR RATS

RAM MOHAN MANDA<sup>1\*</sup>, SRINIVAS REDDY KARKA<sup>2</sup>, GANAPATY SERU<sup>3</sup>

<sup>1</sup>Dept. of Pharmacognosy & Phytochemistry, Talla Padmavathi Pharmacy College, Warangal-506002, Telangana, India, <sup>2</sup>Dept. of Pharmacognosy & Phytochemistry, Vaagdevi Pharmacy College, warangal, Telangana, India, <sup>3</sup>GITAM Institute of Pharmacy GITAM University, Vishakapatnam 530003, Andhrapradesh, India,  
Email: rammohanmanda56@gmail.com

Received: 23 Nov 2014, Revised and Accepted: 24 Dec 2014

### ABSTRACT

**Objective:** The present study was to evaluate the anti-pyretic and wound healing activity of hydro alcoholic extracts of *Evolvulus nummularius* leaves in albino wistar rats.

**Methods:** Anti-pyretic potential was determined by yeast-induced pyrexia and for testing the wound healing property, excision method used in wistar albino rats

**Results:** The present result shows that the hydroalcoholic extract of leaf possesses significant anti-pyretic effect in yeast-provoked elevation of body temperature and The hydro alcoholic extract and leaf juices showed a significant wound healing activity which was well comparable with the standard drug used.

**Conclusions:** The most significant inhibition of wound healing and also the significant reduction in hyperpyrexia in rats when treated with standard drugs as well as different extracts.

**Keywords:** Anti-pyretic, Wound healing activity, *Evolvulus nummularius*, Albino wistar rats.

### INTRODUCTION

*Evolvulus nummularius* Linn. Belongs to the family: Convolvulaceae. It is a small prostrate herb with a small woody branched rootstock; this herb is widely distributed in India, Nepal, Bhutan, Tropical America and Africa. The plant is found as a common weed in lawns, roadsides, grasslands etc. especially in moist places, in India, Madagascan and tropical Africa [1-3]. In Indian traditional folk medicine, the whole plant is used as a medicine for hysteria, to cure burns, cuts, wounds and scorpion stings [4]. In Nepal, the paste of the plant is used to treat scabies [5]. The hot water extract of the dried entire plant is reported to be used in gastric disorder and as sedative [6]. The entire plant is reported to contain evolviside A and Bluteolin and hentriacontane N and triacontane N [7], Sitosterol and its glucoside, stigmasterol, Alpha-mannitol, Ursolic acid and Oleanolic acid [8] and reported to pharmacological activities like anthelmintic [9], anti bacterial and anti oxidant [10] and insecticidal activity [11].

The author has selected plant for to give scientific evidence to the anti pyretic and wound healing based on the folklore claim,

### MATERIALS AND METHODS

#### Plant material

The leaves were collected (3 kg) from the rural belt of Pakala, Warangal district and authenticated by the Botanical Survey of India, Howrah. A voucher specimen (MRM/09/2012) was deposited in the Department of Pharmaceutical Sciences, Andhra University, and Visakhapatnam. The collected plant material was dried under shade and pulverised in a mechanical grinder. The powder was passed through sieve no. 40 and used for further studies.

#### Preparation of extract

The powdered plant material (500 g) was extracted with 2 liters of water-ethanol (1:1) by maceration in closed vessels for 7 days. The solvents were then removed under reduced pressure. The hydro alcoholic extract was brown coloured greasy residue (yield- 5.35% w/w with respect to dry material). The dried extracts (residue) were suspended in 1% gum acacia in normal saline (vehicle) and used for

the Anti-pyretic and wound healing studies. All the experimental procedures were approved by an Institutional animal ethical committee of Talla Padmavathi College of Pharmacy, Warangal, Telangana, India vide approval No. 1505/po/a/11/CPCSEA.

#### Anti-pyretic activity of leaf hydroalcoholic extract of *enummularius*

Hyperthermia was induced in rat sub-cutaneously with 10 ml/kg of 15% aqueous suspension of brewer's yeast, and the rectal temperature was recorded initially and at 18 h. Animals with 1 °C or more elevation in body temperature were used. Hydroalcoholic extract and (100 mg/kg and 200mg/kg, i. p.), paracetamol (100 mg/kg, i. p.) and control vehicle (5 ml/kg, i. p.) were given intraperitoneally after the 18 h yeast injection when the temperature increase was at its peak. The body temperature was measured at 1 h intervals up to 4 h after administration of drug or plant extracts [13].

#### *Evolvulus nummularis* in excised rats

The leaves of *Evolvulus nummularis* are widely used in Indian folk medicine for treating burns. Hence, the author has made an attempt to study the wound healing activity on animal models to support the tribal claim. For testing the wound healing property, a Excision wound model [14] was selected and the procedure was as follows:

The selected Wistar rats were divided into four groups of six in each. The skin hair was removed by using a depilatory cream. Light incisions were made on the cleared surface by cutting the skin of the animals under mild ether anesthesia. The area of the wounds was measured (sq. mm) immediately by placing a transparent polythene graph paper over the wound and then tracing the area of the wound on it. This was taken as the initial wound area reading. All the test samples were applied topically.

**Group-I** served as control. **Group-II** served as reference to which nitrofurazone (0.2 % w/w in simple ointment) was applied topically. **Group-III** animals were treated with the hydro alcoholic extract (10 %w/w in simple ointment I. P. and the Group-IV animals with the juice of the fresh leaves in a similar manner. All the test samples were applied twice daily. The wound area of each animal was measured on 1<sup>st</sup>, 4<sup>th</sup>, 8<sup>th</sup>, 12<sup>th</sup> and 15<sup>th</sup> day. The percentage healing

was calculated from the days of measurements of wound area. The results were presented in table 2.1.2 as Mean  $\pm$  S. E. M. Significance of

difference between control and treated groups was determined using Student's t-test.

**Table 2.1: Anti-pyretic activity of hydro alcoholic extracts of *Evolvulus nummularius***

Groups	Initial temp(c) before yeast	Rectal temp after 18 h of yeast injection				
		0 hr	1 <sup>st</sup> hr	2 <sup>nd</sup> hr	3 <sup>rd</sup> hr	4 <sup>th</sup> hr
Control	37.21 $\pm$ 0.11	38.16 $\pm$ 0.15	38.17 $\pm$ 0.1	38.17 $\pm$ 0.1	38.16 $\pm$ 0.11	38.14 $\pm$ 0.11
Standard (Paracetamol) (100 mg/kg)	36.98 $\pm$ 0.16	38.09 $\pm$ 0.18	37.68 $\pm$ 0.07*	37.32 $\pm$ 0.07**	37.15 $\pm$ 0.03**	36.84 $\pm$ 0.05**
Hydroalcoholic extract(100 mg/. kg)	36.97 $\pm$ 0.1	37.70 $\pm$ 0.05	37.53 $\pm$ 0.06**	37.38 $\pm$ 0.07**	37.18 $\pm$ 0.11	36.94 $\pm$ 0.1**
Hydroalcoholic extract(200 mg/kg)	36.93 $\pm$ 0.10	37.48 $\pm$ 0.04	37.24 $\pm$ 0.14**	37.10 $\pm$ 0.10**	37.0 $\pm$ 0.08**	36.96 $\pm$ 0.12**

Each value represents mean  $\pm$  S. E. M, n = 6. \*P < 0.05, \*\*P < 0.001

**Table 2.1.2: Wound healing activity of the leaf hydroalcoholic extract and fresh leaf juice of *E. nummularius* in excised rats**

Group	Treatment	Percentage inhibition of wound on the day of study				
		1 st	4 th	8 th	12 th	15 th
I	Control	0	19.27 + 3.26	46.87 + 5.87	56.28 + 2.48	62.73 + 1.16
II	Nitrofurazone (0.2% w/w)	0	24.69 + 7.84	57.35 + 7.36	86.24 + 0.82	95.16 + 0.33*
III	Hydroalcoholic extract of <i>E. nummularius</i> (10% w/w)	0	20.6 + 5.68	48.24 + 4.78	71.61 + 2.91	87.12 + 1.28*
IV	Leaf juice of <i>E. nummularis</i>	0	22.46 + 6.75	57.26 + 6.46	83.26 + 1.83	94.24 + 0.48*

Results expressed as Mean + SEM from six observations.\* P < 0.001 on 15<sup>th</sup> day of study.

## RESULTS AND DISCUSSION

The present result shows that the hydroalcoholic extract of leaf possesses significant anti-pyretic effect in yeast-provoked elevation of body temperature in rats, and its effect is comparable to that of paracetamol (standard drug). Furthermore, the hydroalcoholic extract of *E. nummularius* also shows significantly reduced normal body temperature. The studies on wound healing activity revealed that the nitrofurazone treated animals showed 95.16% healing on 15<sup>th</sup> day of study. On the other hand, the extract treated group showed 87.12% healing and the leaf juice treated groups exhibited 94.4% wound healing. Thus the present study justifies its using in the indigenous system of medicine and folklore remedies as anti-pyretic and wound healing activity.

## ACKNOWLEDGMENT

The authors are thankful to the management and faculty of Talla Padmavathi Pharmacy Colleges Warangal for providing necessary facilities to carry out the present research work.

## REFERENCES

- Pullaiah T, Ali Moulali. D, Flora of Andhra Pradesh, volume 2, Scientific Publishers, Jodhpur; 1997. p. 793-4.
- Madhava Chetti K, Sivaji K, Tulasirao K. Flowering plants of chittor district Andhra Pradesh. India, First edition, Tirupati; 2008. p. 16.
- Khare CP. Indian Medicinal Plants, New Delhi, Spinger; 2008. p. 256.
- Jain SK. Dictionary of Indian Folk Medicine and Ethnobotany, New Delhi, Deep. Publication; 1991. p. 92.
- Manandhar NP, Manandhar S. Plants and People of Nepal. Timber Press; 2002. p. 230.
- Chitralkha C, Dey PK, Dey CD. Pharmacological screening of *Valeriana wallichii*, *Lallemantia royleana*, *Breynia rhamnoides* and *Evolvulus nummularius* for sedative and anticonvulsant principle, *Nature Wissenschaften*; 1964. p. 51, 411.
- Biswanath D, Biplab G, Shiho A, Nariko S, Yoshihiro F. Chemical constituents of *Evolvulus nummularius*. *Indian J Chem* 2007;46:492-8.
- Gupta DR, Ahmed B, Dhiman RP, Shoyakugakuzasshi. A Flavone and ts to ne glycosides of *Evolvulus nummularius* 1984;38(4):341-5.
- Dash GK, Bijayini M, Panda A, Patro CP, Ganapaty S. Antihelmintic activity of *Evolvulus nummularius* *Indian. J Nat Prod* 2003;19:24-6.73.
- Pavithra PS, Srevidya N, Verma RS. Antibacterial and antioxidant activity of Methanol extract of *Evolvulus nummularius*. *Indian J Pharmacol* 2009;41:233-336.
- Md Moniruzzaman, Nayeema Parvin, Sabina Sultana, Mohammad Abdullah, Aaur Rahman Khan, Nurul Islam. Evaluation of biological activities of *evolvulus nummularius* through insecticidal, insect repellency and brine shrimp lethality tests. *J Life Earth Sci* 2013;8:101-4.
- Somnath Bhowmik, BK Datta, Sunrit Basu, Sarbadhikary NC. Mandal, Contribution to the less known ethnomedicinally plants used by munda and santal community of india with their ethnomedicinal justification. *World Appl Sci J* 2013;23(10):1408-17.
- JB Perianayagam, SK Sharma, Aney joseph, AJM Chrstina. Evaluation of anti-pyretic and analgesic activity of *Embllica officinalis* Gaertn. *J Ethnopharmacol* 2003;95:83-5.
- Bairy KL, Rao CM. Wound healing profiles of *Ginkgo biloba*. *J Nat Remedies* 2001;1(1):25-7.