

EFFICACY OF PURIFIED PECTINASE OBTAINED FROM PAECILOMYCES VARIOTII IN EXTRACTION AND CLARIFICATION OF JUICE FROM GRAPES AND POMEGRANATE FRUITS

NISHA M. K. *

Department of Botany, Avinashilingam Institute for Home Science and Higher Education for Women (University)
Coimbatore -43, Tamilnadu, India. Email: nishraghunathan@gmail.com

Received: 17 May 2016, Revised and Accepted: 20 July 2016

ABSTRACT

The present investigation was carried out to study the application and the competitiveness of commercial and purified pectinase obtained from *Paecilomyces variotii* in fruit juice (grapes and pomegranate) clarification of different enzyme concentrations like 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, varying incubation time (30, 60, 90, 120 and 160 minutes) at constant temperature of 50°C to optimize the enzymatic treatment for the yield and clarity of the juices. The optimum conditions recommended for enzymatic (crude, purified and commercial pectinase) treatment for clarification and yield of fruit juices were 3.5 mg/20g pulp of enzyme concentration and 180 min incubation time at a constant temperature of 50°C. It was observed that purified pectinase obtained from pectinolytic fungus, *P. variotii* enhanced juice yield and clarity of grape and pomegranate juices and is on par with the commercial pectinase when compared to untreated juices. A maximum yield of 79% and clarity of 19.4 and 19.5% were obtained from grape juice and a significantly high yield of 74% and clarity of 4.9 and 4.8 were achieved from pomegranate juice when compared to the unclarified grape and pomegranate juices (60 and 52% respectively). There was an increase in the yield of 31.6% and 42.3% of the grape and pomegranate juices respectively when treated with purified enzyme than the untreated juices.

Keywords: Pectinase, *Paecilomyces variotii*, Grapes, Pomegranate, Yield, Clarity.

INTRODUCTION

Enzymes are one of the important tools in modern food industry because they simplify many intermediate processes during food processing. Bulk of the industrial enzymes fall into different groups, out of which, the most important group of enzymes is pectinase, used in fruit and vegetable processing industry. Their commercial application was first observed in 1930 for the preparation of wines and fruit juices (Oslen, 2000). Pectinases are one of the important and imminent enzymes of the commercial sector, especially, in the fruit juice industry as a pre-requisite for obtaining well clarified and stable juice with higher yields (Sandriet *al.*, 2011). Pectinases are high molecular weight, negatively charged, acidic glycosidic macromolecules that breakdown complex polysaccharides in plant tissues into simpler molecules with extraordinary specificity, catalytic power and substrate specificity (Approvi and Vuppu, 2012). Pectinases are produced during the natural ripening process of fruits where, it splits polygalacturonic acid into monogalacturonic acid by opening glycosidic linkages. Softening of the cell wall and increase in the yield of juice extract from the fruits takes place during this process. Fungal pectinases are mainly extracellular enzymes, prominent among them being polygalacturonase, which is also most commonly assayed to determine pectinase activity. Pectinase is produced by several fungi including *Aspergillus sp.*, *Botrytis cinerea*, *Fusarium moniliforme*, *Rhizoctonia solani*, *Rhizopus stolonifer*, *Trichoderma sp.*, *Neurospora crassa*, *Penicillium* and *Fusarium* (Joshi *et al.*, 2006). An improved knowledge of the properties of microbial pectinases is important in commercialisation of industrial production and application of these enzymes in various potential fields. Pectinases have attracted attention globally as biological catalysts in numerous industrial processes. These enzymes are used in processing agricultural and agro-industrial waste (Patil and Dayanand, 2006) for the production and clarification of fruit juices to improve the cloud stability of fruit and vegetable juices and nectars, for depectinization in order to produce high density fruit

juice concentrates and for haze removal from wines. As a result, today pectinases are one of the promising enzymes of the commercial sector. Alkaline microbial pectinase reveals a great significance in the current biotechnological arena with wide ranging applications in textile processing, degumming of plant bast fibers, treatment of pectic waste waters, paper making, and coffee and tea fermentations (Pasha *et al.*, 2013). The largest industrial application of pectinases is in fruit juice extraction and clarification. Pectinases contribute to fruit juice viscosity and turbidity. A mixture of pectinases and amylases is used to clarify fruit juices. Treatment of fruit pulps with pectinase also showed an increase in fruit juice volume from banana, grapes and apples (Kaur *et al.*, 2004). With the addition of pectinases, the viscosity of the fruit juice drops, the pressability of the pulp improves, the jelly structure disintegrates and the fruit juice is easily obtained with higher yields. With this background, the present investigation was undertaken to assess the efficacy of the purified pectinase in clarification of fruit juices.

MATERIALS AND METHODS

Collection of fruit samples

Fully ripe fresh grape and pomegranate fruits without any visual blemishes were purchased from local market of Coimbatore, Tamil Nadu. The fruits were washed and rinsed with running water and were ground using a lab mixer for 2-3 min to obtain a homogenous fruit pulp. The grape fruits were extracted from the whole pulp and the pomegranate fruit from the seeds (Figure 1 and 2).

Pre-treatment of extracted fruit pulps

The extracted fruit pulps were pasteurised at 85°C for 3 min to inactivate the natural fruit enzymes and then cooled to 40°C. The fruits are first cut into small pieces and then, pre-treatments like steaming, cooling or heating prior to enzymatic extraction were done to increase juice recovery (Trappey *et al.*, 2008).




Fig. 1:Flow Chart for the extraction of grape juice




Fig. 2:Flow Chart for the extraction of Pomegranate juice

Optimization of enzymatic treatment for the yield and clarity of fruit juice

To optimize the enzymatic treatment, each experiment with 20 g pulp was subjected to the treatment of pectinase obtained from *Paecilomyces variotii* (crude, purified and commercial) of different enzyme concentrations like 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5 mg/20g of pulp, varying incubation time (30, 60, 90, 120 and 160 minutes) at constant temperature of 50°C. At the end of enzymatic treatment, the enzyme in the sample was inactivated by heating the juice at 90°C for 5 min in a water bath.

Evaluation of Juice Yield

The treated juices extracted from the pectinase treated pulp of grapes and pomegranate were centrifuged at 2000 rpm for 10 min using a centrifuge and supernatant was collected and filtered through a muslin cloth spread on a glass funnel and the juice was collected as clear juice. Juice yield was estimated as percentage of juice obtained based on initial pulp. The juice yield was then calculated using the following formula:

Weight of clear juice
 Juice yield % = -----x 100

Weight of sample

Evaluation of juice Clarity

Clarity of the juice was determined by measuring % Transmittance at a wavelength of 660 nm using UV-VIS spectrophotometer according to Tapre and Jain (2014). Distilled water was used as a blank. The percent transmittance was considered as a measure of juice clarity.

Statistical Analysis

Standard errors of means of all the replicates of each variable were computed using Computer Software; Microsoft Excel Data for all experimental data. They were statistically analyzed using 3 way analysis of variance (ANOVA) followed by LSD method to delineate mean differences (Panse and Sukhatme, 1978).

RESULTS AND DISCUSSION

Preliminary experiments were performed to determine the optimum conditions like enzyme concentration and incubation time for maximum yield and clarity of fruit juices. For the optimization of the enzyme treatment, 20 g pulp of grapes and pomegranate were weighed, treated with different concentration and were incubated at a temperature of 40°C for different incubation time.

Optimization of different parameters for the yield and clarity of enzyme treated fruit juices

Effect of enzyme concentration and incubation time on Grape and Pomegranate juice Yield

From Tables 1 and 2, it is clear that with increasing enzyme concentration and incubation time, an increased juice recovery was observed. The yield of grape juice was significantly high with increasing pectinase (crude, purified and commercial) concentrations and incubation time. The results showed significantly high yields of grape juice (69, 79 and 78%) and pomegranate juice (59,74 and 74.5%) using 3.5 mg/ 20g pulp concentration for 180min incubation (crude, purified and commercial enzymes, respectively). Similarly, Thongsombatet *et al.* (2007) obtained a significantly high yield of guava juice using 0.15% pectinase concentration incubated for 2.5 h. Similar result was reported by Ahmed *et al* (2014) who obtained the maximum juice yield at 2.5 hrs. in different concentrations (500, 1000 and 1500 mg.kg⁻¹) as 76, 78 and 80% in guava juice, 76, 78 and 79% in jack fruit juice and 77, 80 and 81% in pine apple juice.

Table - 1 Optimization of enzyme concentration and incubation time on Grape juice Yield (%w/w).

Enzyme concentration mg/ 20g pulp	Crude pectinase					Purified pectinase					Commercial pectinase				
	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min
0.5	64	63.5	64.5	64.5	65.5	71.5	71	71.5	72	73	71	71.5	72.5	74	75
1.0	64	64	64	65.5	65.5	72	72.5	72	73	73	71	73.5	73	74	76
1.5	65	65	65	66.5	66.5	72	73.5	72	72.5	74	71.5	73.5	73.5	74.5	76
2.0	65.5	65.5	66	66.5	67	73	73	73	74	74	71.5	74	74.5	75	77
2.5	65.5	66.5	66.5	67.5	67.5	73.5	73	74	74.5	75	72	74	74.5	75	78
3.0	66	66.5	67.5	67.5	67	74.5	75.5	76	76.5	77.5	72.5	74	76	76.5	78
3.5	66.5	67	67.5	67	69	75.5	77	78	78.5	79	73	75.5	76.5	77	79

Table - 2 Optimization of enzyme concentration and incubation time on Pomegranate juice Yield (%w/w).

Enzyme concentration mg/ 20g pulp	Crude pectinase					Purified pectinase					Commercial pectinase				
	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min
0.5	53	54.5	55.5	56	57	67	68.5	68.5	69	69	67.5	67.5	68	68	69
1.0	53	54	55.5	57	57.5	67	67.5	68	68.5	69	67.5	68.5	68.5	69	69
1.5	53.5	54.5	56	56.5	57	68	68	69	69	70	69	69	69	69	70
2.0	54	55.5	56	57	58	69.5	69	70	70	71	70.5	69.5	70	70.5	71
2.5	54.5	54.5	56	57.5	58.5	69.5	69.5	71	70.5	72	70	71.5	71.5	72	72
3.0	54.5	55	56	57	58.5	71.5	70.5	72	72	73.5	71.5	72	72	73.5	73.5
3.5	55	55.5	56	57.5	59	72	72	73	73	74.5	72	72.5	72.5	74.5	74.5

Evaluation of the enzymes for the yield and clarity of fruit juices (grapes and pomegranate)

Table - 3 Optimization of enzyme concentration and incubation time on Grape juice Clarity (%T)

Enzyme concentration mg/ 20g pulp	Crude pectinase					Purified pectinase					Commercial pectinase				
	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min
0.5	2.1	2.1	2.2	2.3	2.4	16.1	16.1	16.5	16.4	16.5	15.8	16.2	17.0	17.3	17.6
1.0	2.8	2.7	2.7	2.8	2.9	16.3	16.2	16.3	16.4	16.6	16.1	17.1	17.4	17.5	17.6
1.5	3.0	3.2	3.2	3.3	3.2	16.5	16.5	16.7	16.8	16.8	16.4	17.3	17.4	17.4	17.6
2.0	3.7	3.5	3.7	3.8	3.9	16.5	16.7	16.8	16.8	17	16.4	17.3	17.7	17.8	18.2
2.5	4.1	4.2	4.2	4.4	4.3	16.9	16.9	17.2	17.1	17.3	16.4	17.4	17.8	18.1	18.7
3.0	4.3	4.4	4.4	4.7	4.8	17	17.4	17.8	17.8	17.9	17.1	17.3	17.9	18.1	19.2
3.5	4.6	4.6	4.8	4.9	5.2	17.6	17.7	17.8	18.5	19.4	18.2	18.3	18.8	19.4	19.5

Table - 4 Optimization of enzyme concentration and incubation time on Pomegranate juice Clarity (%T).

Enzyme concentration mg/ 20g pulp	Crude pectinase					Purified pectinase					Commercial pectinase				
	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min	30 min	60 min	90 min	120 min	180 min
0.5	0.6	0.6	0.7	0.7	0.7	2.6	2.8	2.8	2.7	2.9	3.1	3.0	3.4	3.4	3.4
1.0	0.8	0.9	1.1	1.1	1.0	2.7	2.7	2.8	2.9	2.9	3.2	3.3	3.4	3.6	3.6
1.5	1.3	1.4	1.4	1.5	1.5	2.7	2.9	2.9	3.1	3.0	3.4	3.5	3.6	3.7	3.8
2.0	1.4	1.5	1.6	1.7	1.7	2.9	3.1	3.1	3.4	3.4	3.3	3.6	3.6	3.8	4.0
2.5	1.7	1.7	1.8	1.9	2.1	3.6	3.6	3.7	3.8	3.8	3.7	3.7	3.8	4.1	4.3
3.0	1.8	1.9	2.0	2.0	2.3	4.1	4.1	4.3	4.4	4.5	3.9	3.9	4.2	4.3	4.3
3.5	2.0	2.1	2.1	2.4	2.7	4.3	4.4	4.5	4.7	4.9	3.9	4.2	4.4	4.7	4.7

Table 5 Yield and Clarity of Grape juice from treated and untreated fruit pulps

Grape juice	Volume of pulp	Volume of juice	Yield (%w/w)	Clarity (%T)
Untreated	20	12.87 ± 1.97	60	0.06
Crude	20	13.07 ± 1.01	69	5.2
Purified	20	15.53 ± 1.12	79	19.4
Commercial	20	15.47 ± 0.76	79	19.5
SEd		1.0601		
CD (p<0.05)		2.4447		

Values are mean ± SD of three samples in each column

Table 6 Yield and Clarity of Pomegranate juice from treated and untreated fruit pulps

Pomegranate juice	Volume of pulp	Volume of juice	Yield %	Clarity%
Untreated	20	10.40 ± 0.80	52	0.08
Crude	20	11.63 ± 0.76	59	2.7
Purified	20	12.63 ± 2.10	74	4.9
Commercial	20	14.77 ± 0.71	74.5	4.7
SEd	-	1.0124	2.3347	-
CD (p<0.05)				

Values are mean ± SD of three samples in each column

From the Tables 5 and 6, it is clear that, purified pectinase obtained from pectinolytic fungus, *P. varioti* enhanced juice yield and clarity of grape and pomegranate juices and is on par with the commercial pectinase when compared to untreated juices. A maximum yield of 79% and clarity of 19.4 and 19.5% were obtained from grape juice and a significantly high yield of 74% and clarity of 4.9 and 4.8 were achieved from pomegranate juice when compared to the unclarified grape and pomegranate juices (60 and 52% respectively). There was an increase in the yield of 31.6% and 42.3% of the grape and pomegranate juices respectively when treated with purified enzyme than the untreated juices. The present result is on par with Singh *et al.* (2012) who observed an increase of 17.5% in bael fruit juice yield from untreated sample at an enzymatic concentration of 20mg/100g pulp, incubation time of 425 min and temperature of 47°C. Similar view was expressed by Srivastava and Tyagi (2013) who reported that the maximum volume of 23.7ml was obtained by pectinase and amylase combination and maximum activity of pectinase enhanced the yield of apple juice upto 34ml/50gm and 25ml/50gm at 5.5 pH and at temperature (45-50°C) respectively. The present findings coincide with the work of Bhardwaj and Garg (2014) who reported that crude pectinase enzyme treatment from *Bacillus* sp. MBRL576 increased the juice volume of 40ml in apple and banana and 50ml in carrot compared to untreated (30, 25 and 40ml) apple, banana and carrot juice respectively and of 25ml in commercial pectinase.

CONCLUSION

Thus it was observed that with an increasing enzyme concentration and incubation time, the yield of the juice increased and also the treated juice became more clear and transparent. The juice yield increased on enzyme treatment as degradation of pectin led to reduction in the water holding capacity of pectin, thus releasing free water into the system and the clarity is due to extended contact between enzyme and substrate. Thus, the present study showed that the usage of purified pectinase obtained from pectinolytic fungus, *P. varioti* enhanced juice yield and clarity when compared to control and also indicated the equal effectiveness and competitiveness of the purified enzyme to that of commercial one.

REFERENCES

- Ahmed B., M.B. Uddin and M. F. Jubayer. 2014. Extraction and standardization of selected fruit juices by enzymatic process. *Peak Journal of Food Science and Technology*, 2(2): 18-27.
- Akesowan and Choonhahirun, 2013. Effect of enzyme treatment on guava juice production using response surface methodology. *Journal of Animal and Plant Sciences*, 23(1): 114-120.
- Apoorvi, C. and S. Vuppu, 2012. Microbially derived pectinases. *Journal of Pharmacy and Biological Science*, 2 (2): 01-05.
- Bhardwaj, V and N. Garg, 2014. Production, purification of pectinase from *Bacillus* sp. MBRL576 isolate and its application in extraction of juice. *International Journal of Science and Research*, (3) 6: 648-652.
- Joshi, V. K., P. Mukesh and N. S. Rana, 2006. Pectin esterase production from apple pomace in solid state and submerged fermentations. *Food Technology and Biotechnology*, 44(2): 253-256.
- Kaur, G., S. Kumar and T. Satyanarayana, 2004. Production, characterization and application of a thermostable polygalacturonase of a thermophilic mould *Sporotrichum Thermophile*. *Bioresource Technology*, 94: 239-243.
- Oslen, H.S., 2000. *Enzymes at work- A concise guide to industrial enzymes and their use*. Novozymes A/S Bagsvaerd, Denmark.
- Panse, V. G and P. V. Sukhatme, 1978. *Statistical methods for agricultural workers*, ICAR, New Delhi, 232-252.
- Pasha, K. M., P. Anuradha and D. Subbarao, 2013. Applications of pectinases in industrial sector. *International Journal of Pure and Applied Sciences and Technology*, 16(1): 89-95.
- Patil, S. R and A. Dayanand, 2006. Optimization of process for the production of fungal pectinases from deseeded sunflower head in submerged and solid state conditions. *Bioresource Technology*, 97(18): 2340-2344.
- Robin, K., K. Sanjay, D. Singh and H. K. Sharma, 2013. Optimization of enzymatic hydrolysis conditions for enhanced juice recovery with optimum quality from alubukhara (*Prunus domestica* L.) fruit. *International Journal of Advanced Research in Engineering and Applied Sciences*, 2(11): 50-67.
- Sandri, I. G., R. C. Fontana, D. M. Barfknecht and M. M. da Silveira, 2011. Clarification of fruit juices by fungal pectinases. *LWT - Food Science and Technology*, 44(10): 2217-2222.
- Sin, H. N., S. Yusof, N. S. A. Hamid and R. A. Rahman, 2006. Optimization of enzymatic clarification of sapodilla juice using response surface methodology. *Journal of Food Engineering*, 73: 313-319.
- Singh, A., S. Kumar and H. K. Sharma, 2012. Effect of enzymatic hydrolysis on the juice yield from bael fruit (*Aegle marmelos* Correa) pulp. *American Journal of Food Technology*, 7(2): 62-72.
- Srivastava, S and K. S. Tyagi, 2013. Effect of enzymatic hydrolysis on the juice yield from apple fruit (*Malus domestica*) pulp. *International Journal of Biotechnology and Bioengineering Research*, 4(4): 299-306.
- Tapre, A. R and R. K. Jain, 2014. Pectinases: Enzymes for fruit processing industry. *International Food Research Journal*, 21(2): 447-453.
- Thongsombat, W., A. Sirichote and S. Chanthachum, 2007. The production of guava juice fortified with dietary fiber. *Songklanakarinn Journal of Science and Technology*, 29 (1): 187-196.
- Trappey, A. F., C. E. Johnson and P. W. Wilson, 2008. Use of a commercial pectolytic enzyme to extract juice from frozen Mayhaw (*Crataegus opaca* Hook.) fruit. *International Journal of Fruit Science*, 7(1): 77-86.

19. Vijayanand, P., S. G. Kulkarni and G. V. Prathibha, 2010. Effect of pectinase treatment and concentration of litchi juice on quality characteristics of litchi juice. *Journal of Food Science and Technology*, 47(2): 235-239.
20. Yannan, S. K., S. Varakumar and O. V. S. Reddy, 2012. Evaluation of antioxidant and sensory properties of mango (*Mangifera indica* L.) wine, *Food Technology and Biotechnology*, 5(3): 359-367