



making near work a favorable and safer routine. Hyperopia has usually been found to be less prevalent than myopia among youths. This could be as a result of the still high accommodative state present in youths which compensate for any latent hyperopia (Megbelayin *et al.*, 2014). Although in young children, hyperopia and astigmatism may be found to be higher than myopia (Jimenez *et al.*, 2012, Krishnamurthy *et al.*, 2014).

**In young children, hyperopia and astigmatism may be found to be higher than myopia (Jimenez *et al.*, 2012, Krishnamurthy *et al.*, 2014).** Hypermetropia is considered to be the most common refractive error in infants. Faderin [10] reported hypermetropia constituting 52.2% of refractive errors in primary schoolchildren in Nigeria while Montes-Mico [11] reported 35.6% prevalence of hypermetropia in Spain. In a survey of the prevalence of refractive errors among children in lower primary school in Kampala district, Kawuma [12] reported astigmatism as the most common single refractive error accounting for 52% of all errors of refraction while Adegbehingbe [9] reported 52.8 in Ile-Ife, Nigeria, Askira [13] reported 22.9% while Presbyopia accounted for 51.3% of patients with refractive errors.

**Table 1: Incidence of refractive errors across the regions of Nigeria**

Region	Total respondents	Number using medicated glasses	Percentage
North-west	31	19	61.29
South-west	97	49	50.51
South-south	295	97	32.88
South-east	300	83	27.67
North-central	122	22	18.03
North-east	19	3	15.79

**Table 2: Distribution of refractive error cases**

S. No.	Refractive error	n	Percentage
1.	Myopia	148	54.21
2.	Hypermetropia	77	28.21
3.	Presbyopia	5	1.83
4.	Astigmatism	43	15.75
	Total	273	100

**Table 3: Distribution of refractive errors across the regions of Nigeria**

	n	%	n	%	n	%	n	%
South-east	47	31.76	21	27.27	1	20.00	11	25.58
South-south	52	35.13	33	42.86	0	0.00	9	20.93
South-west	27	18.24	7	9.09	2	40.00	11	25.58
North-east	3	2.03	0	0.00	0	0.00	0	0.00
North-central	11	7.43	9	11.69	1	20.00	5	11.63
North-west	8	5.41	7	9.09	1	20.00	7	16.28
Total	148	100	77	100	5	100	43	100

**Table 4: Distribution of myopia according to states in Nigeria**

S. No.	State	Number	Percentage	S. No.	State	Number	Percentage
1.	Rivers	27	18.12	11.	Ondo	3	2.01
2.	Lagos	22	14.80	12.	Osun	2	1.34
3.	Delta	21	14.10	13.	Bauchi	2	1.34
4.	Anambra	15	10.10	14.	Kano	2	1.34
5.	Imo	12	8.05	15.	Ebonyi		0.67
6.	Enugu	11	7.38	16.	Oyo		0.67
7.	Abia	9	6.04	17.	Borno		0.67
8.	FCT	8	5.37	18.	Kogi		0.67
9.	Cross River	5	3.36	19.	Sokoto		0.67
10.	Kaduna	4	2.68	20.	Katsina		0.67
					Total	149	100

## METHODS

This study was carried out among the 2<sup>nd</sup> year students of Madonna University Nigeria, Elele campus.

This population was chosen to ensure that whatever visual problems seen were not acquired within the period as a student but was already present before the individual came into the school. This was also ensured further using only respondents who have used medicated glasses for at least 5 years. **Meaning that usage must have commenced before studentship.** One thousand questionnaires were randomly distributed to undergraduate students in various departments in the College of Medicine, including anatomy, physiology, medical laboratory science, optometry, public health, and pharmacy. Furthermore, our distribution of students according to the states of Nigeria is based on residency, not place of origin. As such, this work leans more toward environmental rather than genetic causes. Future analysis will consider the genetic factors. Nine hundred and thirty of these questionnaires were retrieved (93%). Out of that number, 864 (86.4%) met with the conditions stated above and were used for the study. The remaining 66 were discarded. Data were analyzed using simple percentages. Data collected included information on age, sex, state of origin, place of residence with duration, and presence of refractive errors.

## RESULTS

Table 1 shows that the incidence of use of refractive North-west and South-west had the highest incidences of refractive errors with 61.29% and 50.51%, respectively. North-east and North-central had the least incidences with 15.79% and 18.03%, respectively.

From Table 2, myopia had the highest occurrence with 54.21% of all cases, followed by hypermetropia (28.21%), astigmatism (15.75%), and presbyopia (1.83%).

Table 3 shows the distribution of refractive errors across the six geopolitical zones of the country. Myopia was highest in South-south and least in North-east. South-east and South-west had high values. Hypermetropia was also highest in South-south followed by South-east and North-central. It was lowest South-west and North-west.

Table 4 shows the frequency of myopia according to individual states of Nigeria. This frequency, however, only represents a direct numerical counting, comparing a state with the total number of cases. So as expected, states in the south will have more frequency than those in the north as the university is located in Rivers state, South-south part of the country.

Table 5 shows the frequency of hypermetropia by states. As in above, states in the southern part of the country had the highest frequency by number.

Table 6 shows the distribution of presbyopia according to states where cases were recorded in our work.

Table 5: Distribution of hypermetropia according to states in Nigeria

S. No.	State	Number	Percentage	S. No.	State	Number	Percentage
1.	Rivers	19	24.67	10.	Kogi	2	2.60
2.	Delta	8	10.40	11.	Kano	2	2.60
3.	Imo	7	9.09	12.	Cross River		1.30
4.	Anambra	6	7.79	13.	Sokoto		1.30
5.	Edo	6	7.79	14.	Plateau		1.30
6.	Lagos	6	7.79	15.	Kaduna		1.30
7.	FCT	6	7.79	16.	Niger		1.30
8.	Abia	5	6.49	17.	Osun		1.30
9.	Enugu	4	5.19				
					Total	77	100 %

Table 6: Distribution of presbyopia according to states in Nigeria

S. No.	State	Number	Percentage
1.	Lagos	2	50
2.	Anambra	1	25
3.	FCT	1	25
	Total	4	100

Table 7: Distribution of astigmatism according to states in Nigeria

S. No.	State	Number	Percentage	S. No.	State	Number	Percentage
1.	Lagos	11	25.58	8.	Katsina	2	2.33
2.	Imo	7	16.28	9.	Abia		2.33
3.	Delta	6	13.95	10.	Anambra		2.33
4.	FCT	5	11.63	11.	Osun		2.33
5.	Enugu	3	6.98	12.	Plateau		2.33
6.	C. River	2	4.65	13.	Kaduna		2.33
7.	Rivers	2	4.65				
					Total	43	100

Table 7 shows the distribution of astigmatism according to states in Nigeria.

## DISCUSSION

Our results reveal that out of the total of 864 respondents used for this study, 31.60% (273) had a refractive error. It means from our results that the incidence of refractive errors among Nigerian youths in tertiary institution stands at 31.60%. This is comparable to 37.39% reported by Prema, 2011, in a study carried out in India. Our results reveal that North-west region had the highest prevalence of the use of medicated glasses among youths at 61.29%. This is followed by South-west (55.67%), South-south (36.27%), South-east (31.67%), North-central (19.67%), and North-east (15.79%).

The incidence of refractive errors according to the different regions of the country shows that North-west had the highest incidence with 61.29%, South-west 50.51%, South-south 32.88%, South-east 27.67%, North-central 18.03%, and North-east 15.79%. This could be associated with the reported high level of poverty and living conditions in this region of the country. It is in line with the report of Feldkamper and Schaeffel (2003) which shows that environmental factors are associated with the frequency of refractive errors.

The highest occurring refractive error among the six geopolitical regions of Nigeria is myopia (54.21%), followed by hyperopia (28.21%), astigmatism (15.75%), and presbyopia (1.83%). This is similar to the report of Adeoti and Egbewale, 2008, in a study carried out in Ado ekiti where myopia constituted 39.33%, hypermetropia 23.33%, and astigmatism 21.80%. The increasing incidence of refractive errors and consequent use of medicated glasses may be associated with the longer time our youth spend reading books, working on the computer, and watch television according to the report of Prema (2011). This may explain why the incidence was higher in more developed cities

and states of the country than the less developed places. Our study population is undergraduates in one of Nigeria's foremost private university. This population has no difficulty with access to eye clinic or basic medical treatment. South-east and South-south had the highest frequency of myopia. These regions are among the most urbanized in the country. This is in line with the previous works that show a positive correlation between myopia and urbanization/industrialization [10].

## CONCLUSION

Our results suggest that there is a rise in the incidence of refractive errors among our youths, leading to increase in the use of medicated glasses, a phenomenon that was more common among the elderly in our society.

## Recommendation

- We, therefore, recommend that government and its agencies should pay attention to those environmental factors that predispose our youths to visual impairments.
- This study can be extended to specific regions of the country, especially those with observed high incidence from this study.
- Furthermore, attention should be paid to sexual disparity in the manifestation of these refractive errors.

## REFERENCES

1. Adeoti CO, Egbewale BE. Refractive errors in Mercyland specialist hospital, Osogbo, Western Nigeria. Niger Postgrad Med J 2008;15:116-9.
2. Dandona R, Dandona L, Naduvilath TJ, Srinivas M, McCarty CA, Rao GN. Refractive errors in an urban population in Southern India: The Andhra Pradesh eye disease study. Invest Ophthalmol Vis Sci 1999;40:2810-8.
3. Prema N. Causing factors of refractive error in children: Heredity or environment. Indian J Sci Technol 2011;4:???
4. Holden BA, Sulaiman S, Knox K. The challenge of providing spectacles

- 1 in the developing world. *J Community Eye Health* 2000;13:9-10.
- 2 5. National Committee for Prevention of Blindness. National Programme  
3 for Prevention of Blindness: First Five Year Plan. Islamabad, Pakistan:  
4 National Committee for Prevention of Blindness, Ministry of Health,  
5 Special Education and Social Welfare; 1994-98. p. 24.
- 6 6. Feldkammer M, Schaefel F. Interactions of genes and environment in  
7 myopia. *Dev Ophthalmol* 2003;37:34-49.
- 8 7. Zadnik K, Satariano WA, Mutti DO, Cheung YY. The effects of  
9 parental H/o myopia on children eye size. *JAMA* 1994;271:1323-7.
- 10 8. Zadnik K. The Glenn A. Fry Award Lecture (1995). Myopia  
11 development in childhood. *Optom Vis Sci* 1997;74:603-8.
- 12 9. Mutti DO, Mitchell GL, Moeschberger ML, Jones LA, Zadnik K.  
13 Parental myopia, near work, school achievement, and children's  
14 refractive error. *Invest Ophthalmol Vis Sci* 2002;43:3633-40.
- 15 10. Saw SM, Zhang MZ, Hong RZ, Fu ZF, Pang MH, Tan DT. Near-work  
16 activity, night lights, and myopia in the Singapore-China study. *Arch  
17 Ophthalmol* 2002;120:620-7.
- 18 11. Dunaway D, Berger I. Worldwide Distribution of Visual Refractive  
19 Errors and what to Expect at a Particular Location. ????. Presentation  
20 to the International Society for Geographic and Epidemiologic  
21 Ophthalmology; 2006. p. 1-9.
- 22 12. Tebebah T. Pattern of eye diseases in Port Harcourt and an oil producing  
23 rural community. *Niger J Ophthalmol* 1995;3:6-8.
- 24 13. Ayed T, Sokkah M, Charfi O, El Matri L. Epidemiologic study of  
25 refractive errors in schoolchildren in socioeconomically deprived  
26 regions in Tunisia. *J Fr Ophthalmol* 2002;25:712-17.
- 27 14. Chuka-Okosa CM. Refractive Errors among students of a postprimary  
28 institution in a rural community in South-Eastern Nigeria. *West Afr J  
29 Med* 2005;24:62-5.
- 30 15. Curtin BJ. *The Myopias: Basic Science and Clinical Management*.  
31 Philadelphia, PA: Harper and Row; 1985. p. 39-59.
- 32 16. Mutti DO, Zadnik K. Age-related decreases in the prevalence of  
33 myopia: Longitudinal change or cohort effect? *Invest Ophthalmol Vis  
34 Sci* 2000;41:2103-7.
- 35 17. Saw SM, Gazzard G, Koh D, Farook M, Widjaja D, Lee J, *et al*.  
36 Prevalence rates of refractive errors in Sumatra, Indonesia. *Invest  
37 Ophthalmol Vis Sci* 2002;43:3174-80.
- 38 18. Raju P, Ramesh SV, Arvind H, George R, Baskaran M, Paul PG, *et al*.  
39 Prevalence of refractive errors in a rural South Indian population. *Invest  
40 Ophthalmol Vis Sci* 2004;45:4268-72.
- 41 19. Adegbehingbe BO, Majekodunmi AA, Akinsola FB, Soetan EO.  
42 Pattern of refractive errors at Obafemi Awolowo University Teaching  
43 Hospital, Ile-Ife, Nigeria. *Niger J Ophthalmol* 2003;11:76-9.
- 44 20. Faderin MA, Ajaiyeoba AI. Refractive errors in primary school children  
45 in Nigeria. *Niger J Ophthalmol* 2001;9:10-4.
- 46 21. Montes-Mico R, Ferrer-Blasco T. Distribution of refractive errors in  
47 Spain. *Doc Ophthalmol* 2000;101:25-33.
- 48 22. Kawuma M, Mayeku R. A survey of the prevalence of refractive errors  
49 among children in lower primary schools in Kampala District. *Afr  
50 Health Sci* 2002;2:69-72.
- 51 23. Askira BH, Akobundu MN. Refractive errors in Maiduguri. *BOMJ*  
52 2006;3:7-9.

## APPENDIX

## Questionnaire

This research work is been carried out in Madonna University, Elele campus on FREQUENCY OF REFRACTIVE ERRORS AND USE OF MEDICATED GLASSES AMONG NIGERIA YOUTHS. Please kindly respond to the questions as genuinely as possible. Be assured that your confidentiality shall be respected. As the information given cannot be traced to you in any way. All information are strictly for research purposes. Thank you for your cooperation.

**Section A: Personal information (please tick the appropriate box).**

1. Sex: M  F
2. Age: 15-20  21-25  26-30  31-35
3. State of origin:
4. Where have you resided for the most part of your life?
- State  City/Town

5. For how long have you or did you stay there?

**Section B:**

1. Do you use glasses? Yes  No
2. Are your glasses medicated? Yes  No
3. Please specify the type.
- A. Myopia (short-sightedness)
- B. Hypermetropia (long-sightedness)
- C. Presbyopia
- D. Astigmatism
- E. Others (please specify)
4. How long have you used it?  Years
5. How many siblings do you have?
6. How many of them use glasses?

Author Queries???

AQ1: Kindly confirm article type.

AQ2: Kindly check and confirm the author name and its affiliations.

AQ3: Kindly provide history details.

AQ4: Kindly provide structured abstract.

AQ5: Kindly cite author in the reference list.

AQ6: Kindly the author and year doesn't match with reference list.

AQ7: Kindly check the text provided here which is exactly the same as provided under the same heading.

AQ8: Kindly review the sentence as it seems to be incomplete.

AQ9: Kindly provide page number

AQ10: Kindly cite references 1 and 14-23 in text part and also cite in chronological order.

AQ11: Kindly provide location