

Original Article

EFFECT OF THE JOINT SUPPLEMENTATION OF VITAMIN C AND VITAMIN E ON NICKEL HEAMATOTOXICITY AND NEPHROTOXICITY IN MALE SWISS ALBINO MICE

FAOUZI DAHDOUH¹, SALAH ATTALAH², MOHAMED REDA DJABAR¹, ZINE KECHRID³

¹Laboratory of Toxicology, Department of Biology, Faculty of Sciences, Badji Mokhtar University, Annaba, Algeria, ²Department of Animal Biology, Ethnobotany Palynology and Ethnopharmacology- Toxicology Laboratory; Faculty of Natural Science and Life, Montouri University Constantine, Algeria, ³Laboratory of Biochemistry and Applied Microbiology, Department of Biochemistry, Badji Mokhtar University, Annaba, Algeria.
Email: kechridzine@yahoo.fr

Received: 28 Feb 2016 Revised and Accepted: 20 Apr 2016

ABSTRACT

Objective: The aim of this study was to investigate the effect of vitamins C and E separately and in combination against nickel-induced alterations in haematological indices and kidney dysfunction.

Methods: Male Swiss albino mice were divided into eight equal groups: Control, vitamin C (Vit C), vitamin E (Vit E), vitamin C and vitamin E (Vit C+Vit E), nickel (Ni), nickel and vitamin C (Ni+Vit C), nickel and vitamin E (Ni+Vit E), and nickel plus vitamins C and E (Ni+Vit C+Vit E). Vitamin C (1g/l) was given to mice through their drinking water. Vitamin E (1g/kg) and nickel as nickel sulfate (2.7 mg/kg) were supplemented in diet for four weeks.

Results: Nickel caused a significant decrease in body weight, food and water consumption along with significant increase in the absolute and relative kidney weights. Haemoglobin, red blood cells count (RBC), hemoglobin (Hb) concentration, platelet counts (Plt) and packed cell volume (PCV) were significantly diminished, while white blood cells count (WBC) increased in nickel exposed mice. The renal damage induced by nickel was evidenced by a significant increase in the levels of serum urea, creatinine and uric acid. However, vitamins C and E in combination more significantly ameliorated the altered histopathological and biochemical changes in the kidney as well as hematological parameters of Ni intoxicated mice than either vitamin C or E.

Conclusion: The study showed that vitamin C and E combination effectively attenuated Ni-induced heamatotoxicity and nephrotoxicity in mice.

Keywords: Nickel, Vitamins C, Vitamin E, Heamatotoxicity, Kidney injury, Histopathology

© 2016 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

INTRODUCTION

Nickel is a toxic metal that released into the environment by industrial activities, such as the production of batteries and paints, and also a by-product of metal alloys and medical implants. Thus nickel is an important material in industries, and hence, public health environmental problems become inevitable following exposure to nickel compounds via contaminated water and foodstuffs [1, 2]. Nickel cannot be metabolized and, therefore, accumulates in the body, where it can accumulate to high levels in certain organs, in particular, the liver and the kidney, leading to serious health effects [3]. The kidney is particularly predisposed to nickel-induced toxicity and carcinogenicity due to its involvement in nickel toxicokinetics and excretion through urine, as well as it serves as a major organ of Ni accumulation [4]. Several reports are available showing that nickel-induced nephrotoxicity is strongly related with histopathological and serum biochemistry alterations [5, 6].

Moreover, nickel is a known cytotoxic metal that induces kidney cell damages [7, 8]. Although the toxic mechanisms of nickel toxicity and carcinogenicity are still poorly understood. On the other hand, severe adverse heamtotoxic effects including disorders of the bone marrow, haematopoietic systems and the onset of anemia have been noticed in Ni exposed animals [9, 10]. The hematotoxic effect of nickel is basically due to over stimulation of metallothionein and subsequent inducing production of reactive oxygen species (ROS) that leads to oxidative damage in erythrocytes [11, 12]. Recently, the exogenous supplementation of antioxidant molecules such as vitamins C and E have received a wide attention due to their ability to quench reactive oxygen species (ROS) and thereby protecting cellular damage [13]. Vitamin E (dl- α -tocopherol) known as a lipid soluble antioxidant, improves antioxidant defense, prevents lipid peroxidation chain reactions and it can scavenge molecular oxygen, peroxide and hydroxyl radicals and atomic oxygen radicals [14], while vitamin C (ascorbic acid) is known as a potent antioxidant, belongs to the water soluble class of vitamins. Vitamin C

has the ability to scavenge molecular oxygen and hydroxyl radicals and atomic oxygen radicals [15]. Numerous research workers have studied the beneficial effect of vitamin C or vitamin E against nickel toxicity following *in vivo* animal studies [16, 17] and *in vitro* studies [18] over the last few years. To our knowledge, this was the first study on the protective effect of vitamins C and E combination against nickel-induced nephrotoxicity and heamatotoxicity in mice.

MATERIALS AND METHODS

Animals

Sixty-four male Swiss albino mice weighing 29–33 g (Animal Unit of "Pasteur" Algiers Institute, Algeria) were used in this study. Animals were housed in plastic cages under a light-dark cycle 12/12 hour, a temperature 22 ± 2 °C and humidity 40% with *ad-libitum* access to food and water. All experiments were carried out in accordance with ethical approval (AFRO. 123, 2009). Unless otherwise noted, all chemicals were obtained from Sigma Chemical Company (St Louis, France).

Experimental design

Mice were randomly divided into eight groups of eight mice each and subjected to various daily treatment regimes:

Group I: (Control mice) animals fed a standard diet and given bidistilled drinking water.

Group II (Vit C): Mice received a standard diet and vitamin C in bidistilled drinking water (1g/l).

Group III (Vit E): Animals received standard diet enriched with vitamin E (1g/kg of diet).

Group IV (Vit C+Vit E): Mice received vitamin E (1g/kg of diet) and vitamin C (1g/l of bidistilled drinking water).

Group V (Ni): Mice received standard diet supplemented with nickel sulfate (2,7g NiSO₄/kg diet).

Group VI (Ni+Vit C): Mice received standard diet supplemented with nickel sulfate (2,7g NiSO₄/kg) and vitamin C in their drinking water (1g/l).

Group VII (Ni+Vit E): Animals received standard diet supplemented with nickel sulfate and vitamin E (1g Vit E+2,7g NiSO₄/kg diet).

Group VIII (Ni+Vit C+Vit E): Mice received regimen the same as Group VII (1g Vit E+2,7g NiSO₄/kg diet), except that their drinking water contained vitamin C (1g/l).

Vitamin C, vitamin E and nickel sulfate doses were chosen depending on previously established protocols [19], [20] and [21] respectively. Throughout the study, body weight, food intake and water consumption were monitored every three days. After thirty days, animals were killed for blood collection and kidney was removed immediately, fixed in bouin's solution and used for histological examination.

Biochemical analysis

Serum biochemical markers: uric acid, creatinine and urea measured as a functional marker for nephrotoxicity. They were assessed using commercial diagnostic kits (Spinreact, Spain, ref: Creatinine-1001111, urea-1001329 and uric acid-1001011). Hematological parameters: red blood cell (RBC) count, white blood cell (WBC) count, hemoglobin (Hb) concentration, packed cell volume (PCV), and platelet count (Plt) were estimated utilizing fully automated hematological cell counter (Coulter Counter Lympho, serial « T » 540).

Histological evaluation

Slices of the kidney from each experimental group were dissected and immediately fixed in neutral buffered formalin for 24 h, processed by using a graded ethanol series, and then embedded in paraffin. The paraffin sections were cut into 5 µm thick slices and stained with haematoxylin and eosin for light microscopic examination.

Statistical analysis

Results were displayed as mean±standard error of the mean (S. E. M), and were subjected to statistical significance evaluation. Comparisons of multiple groups of each parameter separately were tested by one-way analysis of variance with Turkey post hoc test. The level of significance was set at p<0.05.

RESULTS

Effect of treatment on body weight, absolute and relative kidney weights

In table 1, the body weight of animals exposed to nickel was significantly decreased (p<0.01) as compared to control mice. Treatment with vitamin C and/or E along with nickel showed an improvement in body weight gain (p<0.01, p<0.001) compared to nickel group.

The absolute and relative kidney weights were significantly increased (p<0.01, p<0.001) in nickel treated animals and conversely a significant decrease was noticed (p<0.05, p<0.01) in the combined treatments, with more efficacy in vitamins C and E than either vitamin C or E along with nickel in comparison with nickel exposed mice.

Effect of treatments on Food and Water Intake

As seen in table 1, there was a significant decrease in water (p<0.001) and food intake (p<0.01) in nickel treated mice as compared to control group. Groups: Ni+Vit C, Ni+Vit E and Ni+Vit C+Vit E also showed a significant decrease (p<0.05) of these parameters when compared with control group, but when compared with nickel group, a significant increase of food and water consumption was observed; Ni+Vit C, Ni+Vit E (p<0.05) and Ni+Vit C+Vit E (p<0.01). Hence, the studied parameters were more improved in Ni+Vit C+Vit E treated the group as compared with either of Ni+Vit C or Ni+Vit E treatments.

Table 1: Effect of nickel and its combination with vitamin C and/or vitamin E on body weight, kidney weights (absolute and relative), food and water intake


Groups	Body weight (g)		Kidney weights		Food intake g/100g bwt	Water intake (ml/day)
	Initial	Final	Absolute (g)	Relative g/100g bwt		
Control	30±0.6	41±1.11	1.2±0.05	0.33±0.03	6.62±0.34	7.81±0.37
Vit C	31±0.71	38±0.22	1.19±0.02	0.34±0.01	6.09±0.26	7.06±0.24
Vit E	33±1.1	40±0.17	1.2±0.03	0.32±0.02	5.86±0.21	7.18±0.11
Vit C+Vit E	29±1.03	40±0.23	1.18±0.02	0.33±0.02	6.12±0.30	7.43±0.08
Ni	29±1.05	18±0.13**	1.85±0.07**	1.17±0.02***	3.52±0.15**	4.46±0.17***
Ni+Vit C	32±0.9	26±1.21*##	1.4±0.02**#	0.7±0.04*#	4.7±1.2*##	6.75±0.5*#
Ni+Vit E	33±1.07	28±0.73*##	1.38±0.03**#	0.68±0.06*#	4.9±1.08*##	6.83±0.3*#
Ni+Vit C+Vit E	31±0.66	30±0.45*###	1.3±0.02*##	0.56±0.04*##	5.15±0.6*##	6.92±0.38*##

Values are given as mean±SEM of eight mice each group. Statistically Significant differences from control: *p<0.05, **p<0.01, ***p<0.001; from Ni: #p<0.05, ##p<0.01, ###p<0.001; from Ni+Vit C+Vit E: \$p<0.05, \$\$p<0.01.

Effect of treatment on renal function

Kidney dysfunction was noted by a significant increase in serum urea (p<0.001), creatinine (p<0.001) and uric acid (p<0.01) levels of nickel exposed mice as compared to the control mice. Similarly, these functional kidney markers were significantly increased (p<0.05) in Ni+Vit C, Ni+Vit E and Ni+Vit C+Vit E when compared with the control group.

But when compared with nickel group, a significant decrease of the above mentioned parameters was noticed [serum urea: Ni+Vit C, Ni+Vit C+Vit E (p<0.01) and Ni+Vit E (p<0.05); serum uric acid: Ni+Vit C (p<0.05), Ni+Vit E and Ni+Vit C+Vit E (p<0.01); serum creatinine: Ni+Vit C, Ni+Vit C+Vit E (p<0.01) and Ni+Vit E (p<0.05)]. In addition, vitamin C and vitamin E concomitant treatment with nickel effectively reduced more the above biochemical parameters when compared to nickel along with either vitamin C or vitamin E (fig. 1).






Fig. 1: Serum kidney biochemical markers; creatinine, urea and uric acid in control and experimental animals

Values are given as mean±SEM of eight mice each group. Significantly differences from control: *p<0.05, **p<0.01, ***p<0.001; from Ni: #p<0.05, ##p<0.01, ###p<0.001; from Ni+Vit C+Vit E: §p<0.05, §§p<0.01.

Table 2: Effect of nickel and its combination with vitamin C and/or vitamin E on Red blood cells (RBC), white blood corpuscle (WBC) and blood platelet (Plt) counts, hemoglobin concentration (Hb) and packed cell volume (PCV)

Groups	Red blood cells (10 ⁶ cell/μl)	Packed cell volume (%)	Heamoglobin (mg/dl)	White blood cells (10 ³ cell/μl)	Platelet counts (10 ³ cell/μl)
Control	7.28±0.4	44.12±0.9	13.37±0.6	6.06±0.22	721.6±42.7
Vit C	6.97±0.36	43.27±1.63	12.84±0.5	6.11±0.17	709±31.7
Vit E	7.03±0.43	42.61±1.7	12.79±0.44	6.04±0.1	713±31.4
Vit C+Vit E	7.06±0.12	43.23±2.01	12.9±0.26	6.01±0.08	711±32.3
Ni	5.46±0.65***	32.7±0.13***	10.1±0.08**	8.05±0.40***	426±44.6***
Ni+Vit C	6.71±0.48*§§	37.02±4.5*§	11.3±0.57*§§	6.98±0.28*#	676.08±33*##§§
Ni+Vit E	6.63±0.5*##§§	36.7±3.44*#§	11.1±0.37*#§	6.84±0.34*##§§	662±36.2*##§§
Ni+Vit C+Vit E	6.79±0.43*##	38.9±3.25*##	11.85±0.59*##	6.39±0.22*##	697±35.2*##

Values are given as mean±SEM of eight mice each group. Statistically Significant differences from control: *p<0.05, **p<0.01, ***p<0.001; from Ni: #p<0.05, ##p<0.01, ###p<0.001; from Ni+Vit C+Vit E: §p<0.05, §§p<0.01.

A kidney from nickel treatment (E) showed severe proximal tubule degeneration and tubular necrosis (up than 75%) (double arrows), multiple foci of hemorrhage and inflammation with the presence of lymphocytes in interstitial tissue (single arrow).

In the combined treatments (F, H and G sections), slight lesion of the tubules with only mild tubular necrosis and interstitial inflammation were observed, when compared with untreated mice and those receiving only Ni. The severity of tissue alterations ranged from 60%, 40% and 30% in Ni+Vit C (F), Ni+Vit E (G) and Ni+Vit C+Vit E (H) respectively.

DISCUSSION

The uncontrolled release of hazardous substances, including heavy metals, may have serious effects on kidney function [22] and

Effect of treatments on hematological parameters

As showed in table 2, nickel treatment resulted significant decrease of red blood cells (RBCs), platelet counts, packed cell volume (PCV %) (p<0.001) and hemoglobin (Hb) concentration (p<0.01) concomitant with significant increase of white blood cell (WBC) count (p<0.01) in nickel group when compared to control group. Also, Ni+Vit C, Ni+Vit E and Ni+Vit C+Vit E treatments showed a significant increase (p<0.05) of those parameters in comparison with control group, but when compared with nickel group, a significant increase of red blood cells: Ni+Vit C, Ni+Vit E (p<0.05) and Ni+Vit C+Vit E (p<0.01); packed cell volume: Ni+Vit E (p<0.05) and Ni+Vit C+Vit E (p<0.01); heamoglobin: Ni+Vit C and Ni+Vit E (p<0.05) and Ni+Vit C+Vit E (p<0.01); blood platelet: Ni+Vit C, Ni+Vit E and Ni+Vit C+Vit E (p<0.05) and a decrease of white blood cells: Ni+Vit C, Ni+Vit C+Vit E (p<0.05) and Ni+Vit E (p<0.01).

Whereas, treatments with both vitamins C and E exhibited a good protection than either vitamin C or E separately against nickel haematotoxicity. No significant changes were noticed in the studied parameters in either vitamin E or vitamin C treated animals when compared with untreated animals.

Histological results

The average score obtained for each animal and then for each group were considered as final damage tissue score (fig. 2). Normal histology of the glomeruli and tubules in cortex and medulla was observed in the kidney of control mice (fig. 2A). The histoarchitectural patterns of kidney were almost normal in mice treated with Vit C and Vit E alone or in combination (fig. 2B-2D).

Histopathological studies showed that nickel-induced proximal tubule degeneration and tubular necrosis (double arrows), multiple foci of hemorrhage and inflammation with the presence of lymphocytes in interstitial tissue (single arrow) (fig. 3E). Indeed, pathological lesions induced by Ni were remarkably reduced by the administration of vitamin C or vitamin E, but both vitamins along with Ni proved to be more efficient in reducing nickel-induced severe tissue alterations (fig. 3F-3H).

hematopoietic system [23]. Early reports suggest vitamins C and E as potent antioxidants, that are able to reduce oxidant threat mediated renal damage induced by heavy metals, such as cadmium [24] and mercury [25]. Therefore, this study was devoted to determining the beneficial effects of both vitamins on nickel-induced kidney injury and hematotoxicity in mice. The decrease in final body weight, food and water intake along with significant increase in kidney weight was seen in nickel-exposed group.

These findings are concomitant with previously reports [26]. The decreased body weight is possibly due to the overall increased degeneration of lipids and proteins and the significant increase in both absolute and relative kidney weights may explain the selective accumulation of nickel in the kidney [27]. All these physiological changes observed in nickel treated mice were attenuated by

treatment with vitamin C and E separately or in combination. In this context, the present study confirmed that both vitamins have ameliorative effect against nickel-alteration in body weight of mice. Chatterjee *et al.* [28] suggested that enhancement of body weight

would result in a concomitant increase in the daily food consumption and promotion of protein synthesis. Similar data were also reported in vitamin E co-administration with silver [29] and nickel nanoparticles [30].




Fig. 2: Light photomicrograph of kidney sectional histology from control mice and vitamin C and/or E treated mice (H&E. X 40): Control group (A), Vitamin C (C), Vitamin E (D) and Vitamins C and E (E) treated groups, showing no histological changes were observed in the kidney of vitamin C and/or vitamin E treated animals (B-D) when compared with control kidney (A)




Fig. 3: Light photomicrograph of sectional kidney histology from mice treated with Ni only and those treated with vitamin C/E in combination with Ni (H&E. X 40)

In this study, we noticed renal impairment in association with an increase in serum urea, uric acid and creatinine, as well as renal histopathological changes in Ni-treated mice. The increased level or

uric acid could be linked to increased degradation of purines and pyrimidines, over production or inability of its excretion [31]. The supplementation of vitamin E and/or vitamin C along with nickel

improved the altered biochemical markers and kidney histology induced by nickel, which somehow may result in the antiradical/antioxidant activity. Accordingly, these findings are in line with those obtained by Appenroth *et al.* [32], who found that vitamin E and C combination attenuated the cisplatin-induced nephrotoxicity in rats. The protective effects of these antioxidants are owed to their ability to reduce oxidative stress in the kidney the following exposure to several metals [33]. The hematological indices also showed that treatment with nickel-induced significant decrease in red blood cells (RBC), hemoglobin (Hb) concentration, percentage packed cell volume percentage (PCV%) and platelets count.

Thus, the results indicated that exposure to nickel could lead to anemia. In addition, it was suggested that nickel may adversely affect the hematopoietic process and bone marrow activity resulting in a reduction of RBC and Hb, which is likely due to iron deficiency or chemically induced anemia [34]. Nickel is also found to induce oxidative injury in erythrocytes following generation of reactive oxygen species (ROS)[35]. Also, the low hemoglobin concentration in this study may be as a result of a decrease in the succinyl and glycine pools, as well as the key enzymes such as, ALAD (Aminolevulinic Acid Dehydratase) that are required in the hem biosynthesis [36]. In contrast, white blood cell (BWC) count appeared a significant increase in nickel-exposed animals. This result is in line with earlier reports on rats with potential exposure to arsenic [37] and to cadmium [38], where the plausible explanation of the elevation in white blood cells is due to their contribution to the inflammatory response to toxic metals. Vitamin C and E alone or in combination improved the studied hematological parameters approximately to their normal values. These vitamins are likely involved in activating blood cell productions and hemoglobin synthesis by stimulating bone marrow, as well as they, act as antioxidants to reduce oxidative injury in erythrocytes.

CONCLUSION

The present study indicated that vitamins C and vitamin E combination proved to be more efficiently than either vitamin C or vitamin E to alleviate nickel-induced nephrotoxicity and hematotoxicity. Furthermore, it provides biological evidence supporting the beneficial interaction (synergistic) between the antioxidant vitamins in protecting nickel toxicity.

ACKNOWLEDGEMENTS

The authors greatly knowledge the Ministry of Higher Education and Scientific Research for supporting this research (project reference number: N ° FOI 120070054).

CONFLICT OF INTERESTS

Declared none

REFERENCES

- De Brouwere K, Buekers J, Cornelis C, Schlekot CE, Oller AR. Assessment of indirect human exposure to environmental sources of nickel: oral exposure and risk characterization for systemic effects. *Sci Total Environ* 2011;419:25-36.
- Leblanc JC, Guérin T, Noël L, Calamassi-Tran G, Volatier JL, Verger P. Dietary exposure estimates of 18 elements from the 1st french total diet study. *Food Addit Contam* 2005;22:624-41.
- Cempel M, Janicka K. Distribution of nickel, zinc, and copper in rat organs after oral administration of nickel (II) chloride. *Biol Trace Elem Res* 2002;90:215-26.
- Bardack S, Dalgard CL, Kalinich JF, Kasper CE. Genotoxic changes to rodent cells exposed *in vitro* to tungsten, nickel, cobalt and iron. *Int J Environ Res Public Health* 2014;11:2922-40
- Boulila S, Elfeki A, Oudadesse H, Elfeki H. Substitution effects of a carbonated hydroxyapatite biomaterial against intoxication chloride nickel-exposed rats. *Toxicol Mech Methods* 2015;25:155-65.
- Jadhav SH, Sarkar SN, Patil RD, Tripathi HC. Effects of subchronic exposure via drinking water to a mixture of eight water-contaminating metals: a biochemical and histopathological study in male rats. *Arch Environ Contam Toxicol* 2007;53:667-77.
- Permenter MG, Lewis JA, Jackson DA. Exposure to nickel, chromium, or cadmium causes distinct changes in the gene expression patterns of a rat liver derived cell line. *PLoS One* 2011;6:e27730. Doi:10.1371/journal.pone.0027730. [Article in Press]
- Huang X, Zhuang Z, Frenkel K, Klein CB, Costa M. The role of nickel and nickel-mediated reactive oxygen species in the mechanism of nickel carcinogenesis. *Environ Health Perspect* 1994;102 Suppl 3:281-4.
- Tikare SN, Yendigeri S, Gupta AD, Dhundasi SA, Das KK. Effect of garlic (*Allium sativum*) on hematology and erythrocyte antioxidant defense system of albino rats exposed to heavy metals (nickel II & chromium VI). *Indian J Physiol Pharmacol* 2012;56:137-46.
- Adjroud O. The toxic effects of nickel chloride on liver, erythropoiesis, and development in Wistar albino preimplanted rats can be reversed with selenium pretreatment. *Environ Toxicol* 2013;28:290-8.
- Tkshelashvili LK, Tsakadze KJ, Khulusauri OV. Effect of some nickel compounds on red blood cell characteristics. *Biol Trace Elem Res* 1989;21:337-42.
- Das KK, Saha S. L-ascorbic acid and α tocopherol supplementation and antioxidant status in nickel or lead-exposed rat brain tissue. *J Basic Clin Physiol Pharmacol* 2010;21:325-46.
- Emilie P, Maurice H, Gilles C, Joel P, Jacques M, Dalila C. Oxidative stress and dystrophy facioscapulohumeral: effects of vitamin C, vitamin E, zinc gluconate and selenomethionine supplementation. *Free Radical Biol Med* 2014;75 Suppl 1:S14.
- Pascoe GA, Fariss MW, Olafsdottir K, Reed DJ. A role of vitamin E in protection against cell injury. Maintenance of intracellular glutathione precursors and biosynthesis. *Eur J Biochem* 1987;166:241-7.
- Cathcart RF. 3rd. Vitamin C: the nontoxic, non-rate-limited, antioxidant free radical scavenger. *Med Hypotheses* 1985;18:61-77.
- Das KK, Buchner V. Effect of nickel exposure on peripheral tissues: role of oxidative stress in toxicity and possible protection by ascorbic acid. *Rev Environ Health* 2007;22:157-73.
- Hattiwale SH, Saha S, Yendigeri SM, Jargar JG, Dhundasi SA, Das KK. Protective effect of L-ascorbic acid on nickel-induced pulmonary nitrosative stress in male albino rats. *Biometals* 2013;26:329-36.
- Kasprzak KS, Diwan BA, Kaczmarek MZ, Logsdon DL, Fivash MJ, Salnikow K. Effects of ascorbic acid on carcinogenicity and acute toxicity of nickel subsulfide, and on tumor transplants growth in gluconolactone oxidase knock-out mice and wild-type C57BL mice. *Toxicol Appl Pharmacol* 2011;257:32-7.
- Yildirim O, Büyükbıngöl Z. *In vivo* effect of vitamin C with cobalt on oxidative stress in experimental diabetic rat kidney. *Diabetes Nutr Metab* 2003;16:208-13.
- Alberts DS, Peng YM, Moon TE. Alpha-Tocopherol pretreatment increases adriamycin bone marrow toxicity. *Biomedicine* 1978;29:189-91.
- Schiffer RB, Sunderman FW Jr, Baggs RB, Moynihan JA. The effects of exposure to dietary nickel and zinc upon humoral and cellular immunity in SJL mice. *J Neuroimmunol* 1991;34:229-39.
- Barbier O, Jacquillet G, Tauc M, Cougnon M, Poujeol P. Effect of heavy metals on, and handling by, the kidney. *Nephron Physiol* 2005;99:105-10.
- Brucka-Jastrzębska E, Protasowicki M. Effects of cadmium and nickel exposure on haematological parameters of common carp, *Cyprinus carpio* L. *Acta Ichthyol Piscatoria* 2005;35:29-38.
- Karabulut-Bulan O, Bolkent S, Yanardag R, Bilgin-Sokmen B. The role of vitamin C, vitamin E, and selenium on the cadmium-induced renal toxicity of rats. *Drug Chem Toxicol* 2008;31:413-26.
- Muthu K, Krishnamoorthy P. Effect of Vitamin C and Vitamin E on mercuric chloride-induced reproductive toxicity in male rats. *Biochem Pharmacol* 2012;1:102.
- Milton Prabu S, Shagirtha K, Renugadevi J. Quercetin in combination with vitamins (C and E) improves oxidative stress and renal injury in intoxicated cadmium rats. *Eur Rev Med Pharmacol Sci* 2010;14:903-14.

27. Obone E, Chakrabarti SK, Bai C, Malick MA, Lamontagne L, Subramanian KS. Toxicity and bioaccumulation of nickel sulfate in Sprague-Dawley rats following 13 w of subchronic exposure. *J Toxicol Environ Health A* 1999;57:379-401.
28. Chatterjee K, Chakarborty D, Majumdar K, Bhattacharyya A, Chatterjee GC. Biochemical studies on nickel toxicity in weanling rats--influence of vitamin C supplementation. *Int J Vitam Nutr Res* 1979;49:264-75.
29. Yin N, Yao X, Zhou Q, Faiola F, Jiang G. Vitamin E attenuates silver nanoparticle-induced effects on body weight and neurotoxicity in rats. *Biochem Biophys Res Commun* 2015;458:405-10.
30. Kong L, Tang M, Zhang T, Wang D, Hu K, Lu W, *et al.* Nickel nanoparticles exposure and reproductive toxicity in healthy adult rats. *Int J Mol Sci* 2014;15:21253-69.
31. Amudha K, Pari L. Beneficial role of naringin, a flavanoid on nickel-induced nephrotoxicity in rats. *Chem Biol Interact* 2011;193:57-64.
32. Appenroth D, Fröb S, Kersten L, Splinter FK, Winnefeld K. Protective effects of vitamin E and C on cisplatin nephrotoxicity in developing rats. *Arch Toxicol* 1997;71:677-83.
33. Atef M, Al-Attar. Antioxidant effect of vitamin E treatment on some heavy metals-induced renal and testicular injuries in male mice. *Saudi J Biol Sci* 2011;18:63-72.
34. Hosokawa S1, Nishitani H, Umemura K, Tomoyoshi T, Sawanishi K, Yoshida O. Relationship between serum nickel concentrations and anaemia in chronic haemodialysis patients. *Int Urol Nephrol* 1987;19:447-51.
35. Novelli EL, Rodrigues NL, Nagahashi AM, Sforcin JM, Ribas BO. Nickel chloride effects on erythrocyte generation of superoxide radical. *Braz J Med Biol Res* 1990;23:643-5.
36. Das Gupta A, Das SN, Dhundasi SA, Das KK. Effect of garlic (*Allium sativum*) on heavy metal (nickel II and chromium VI) induced alteration of serum lipid profile in male albino rats. *Int J Environ Res Public Health* 2008;5:147-51.
37. Klibet F, Boumendjel A, Khiari M, El Feki A, Abdennour C, Messarah M. Oxidative stress-related liver dysfunction by sodium arsenite: alleviation by pistacia lentiscus oil. *Pharm Biol* 2016;54:354-63.
38. Karmakar R, Bhattacharya R, Chatterjee M. Biochemical, a haematological and histopathological study in relation to time-related cadmium-induced hepatotoxicity in mice. *Biometals* 2000;13:231-9.