

Original Article

DETERMINATION OF BIOACTIVE CONSTITUENTS OF LEAVES OF *CORCHORUS AESTUANS* (L.) BY GC - MS ANALYSIS

DHANALAKSHMI R\*, MANAVALAN R

Department of Pharmacy, FEAT, Annamalai University, Annamalai Nagar 608002, Chidambaram, Tamilnadu, India.  
Email: dhanama@gmail.Com

Received: 26 Jun 2014 Revised and Accepted: 18 Aug 2014

ABSTRACT

**Objective:** To investigate the bioactive constituents of ethanol extract of *Corchorus aestuans* (L.) using GC-MS technique.

**Methods:** The ethanol extract of leaves were subjected to GC- MS analysis using GC Clarus 500 (Perkin Elmer) with Elite- 5MS (5%Diphenyl / 95% Dimethyl poly siloxane, 30 mm x 0.25 mm x0.25µm df). The 2 µl extract sample injected and the components were separated using Helium (1 mL/min) as the carrier gas. Bioactive constituents were detected by the Turbo mass gold detector (Perkin Elmer) with the aid of the Turbomass 5.2 software.

**Results:** The analysis revealed the presence of fourteen different bioactive constituents namely 3, 7,11,15-tetramethyl-2-hexadecen-1-ol (5.6%), Trans-2-undecen-1-ol (1.26%), E-7-Tetradecenol (1.97%), n-Hexadecanoic acid (25.82%), Phytol (22.34%), 9,12,15-octadecatrienoic acid, methyl ester, (Z,Z,Z)- (20.23%), Docosanoic acid, ethyl ester (1.99%), 1-Eicosanol (2.11 %), 9,9-dimethoxybicyclo[3.3.1] nona-2,4-dione (0.60%), Heptadecanoic acid, heptadecyl ester (0.95%), Pentadecanoic acid, 2,6,10,14-tetramethyl-methylester(0.91%), 3-Hexadecyloxy carbonyl-5-(2-hydroxyethyl)-4-methylimidazolium ion (0.90%), Squalene (8.03%), Vitamin E (7.24%).

**Conclusion:** The presence of various bioactive compounds confirms the application of *Corchorus aestuans* in various disorders. However, isolation of individual phytochemical constituents may proceed to find a lead for herbal therapy.

**Keywords:** *Corchorus aestuans*, Ethanol extract, GC-MS analysis, Bioactive constituents.

INTRODUCTION

Nature has been a source of medicinal plants for thousands of years. Medicinal plants are the chemical gold mines existing in the ecosystems and play a dominant role in the development of novel drugs for the prevention and treatment of disease in men and animals [1]. World Health Organization (WHO) supports the use of medicinal plants, provided it is proven to be efficacious, safe, less toxic, available and reliable natural resource [2]. Phytochemical constituents are responsible for medicinal activity of plants species [3]. The chemical features of these constituents differ considerably among different species. A knowledge of the chemical constituents of plants is desirable not only for the discovery of therapeutic agents, but also because such information may be of great value in disclosing new sources of economic phyto compounds for the synthesis of complex chemical substances and for discovering the actual significance of folkloric remedies [4]. Hence, for the discovery of lead compounds for use as therapeutic drugs, the active principals in medicinal plants needs to be identified [5]. Extraction is the main step for the recovery and isolation of bioactive phytochemical from plant materials, before component analysis [6]. *Corchorus aestuans* (L.) (Syn. *Corchorus acutangulus* Lam), family Tiliaceae is an annual herb occurring throughout the warmer parts of India as weed in the microclimatic areas of river banks where sandy soil is available and also in shade conditions undisturbed areas [7]. They are reported to exhibit various pharmacological activities such as stomachic, anti-inflammatory and pneumonia [8]. But there have been no reports on the bioactive constituents from ethanol extract of leaves of *Corchorus aestuans* Linn. Hence, the present study is to determine the bioactive constituents present in the Ethanol extract of *Corchorus aestuans* leaves by Gas chromatography and Mass spectroscopy (GC-MS) technique.

MATERIALS AND METHODS

Collection and identification of plant materials

Fresh leaves of the selected plant *Corchorus aestuans* were collected from Thirunelveli district, Tamil Nadu, India. The plant materials were taxonomically identified and authenticated by Dr. V. Chelladurai,

Research officer - Botany (scientist C), Central council for research in Ayurveda and Siddha, Govt. of India; Thirunelveli. The plant was thoroughly washed in running tap water to remove soil particles and adhered debris and finally washed with sterile distilled water. The leaves of the plant alone were segregated and dried under shade, pulverized by a mechanical grinder into fine powder. The powdered materials were stored in air tight polythene bags till use.

Preparation of extracts

The powdered plant materials were extracted with ethanol (99.9%). The extraction was done by hot continuous percolation method in Soxhlet apparatus for 24 hrs [9]. The extract was concentrated by using a rotary evaporator till dry powder was obtained. The final residue thus obtained was then subjected to GC-MS analysis [10].

Gas Chromatography-Mass spectrometry (GC-MS) analysis

The ethanol extract of *Corchorus aestuans* leaves was analyzed through GC-MS for the identification of different compounds. The GC-MS analysis was carried out by using Clarus 500 (Perkin - Elmer) Gas chromatograph equipped and coupled to a mass detector Turbo mass gold (Perkin - Elmer) spectrometer with an Elite - 5MS (5% Diphenyl / 95% Dimethyl poly siloxane, 30m x 0.25 mm x 0.25 µm df) of capillary column. The oven was set to an initial temperature 110°C for 2 min, further increased up to 200 °C at the rate of 10 °C/min. Finally temperature was raised up to 280°C, at the rate of 5°C /min for 9 min. Helium gas (99.999%) was used as the carrier gas at constant flow rate of 1 ml/min. An aliquot of 2µl of sample was injected into the column with the injector temperature at 250°C (Split ratio of 10:1). The electron ionization system with ionizing energy of 70 eV was used. Mass spectral scan range was set at 45-450 (m/z). Total GC running time was 36 minutes.

Identification of compounds

Interpretation of mass spectrum of GC-MS was conducted using the database of National Institute Standard and Technique (NIST Version-Year 2005) having more patterns. The relative percentage amount of each component was calculated by comparing its average

peak area to the total areas. The spectrum of the unknown component was compared with the spectrum of the component stored in the NIST data library (version 2005). The name, molecular weight, molecular formula and structure of the components of the test material were determined.

## RESULTS AND DISCUSSION

GC-MS chromatogram of the ethanol extract of leaves of *Corchorus aestuans* (Fig. 1) clearly showed fourteen peaks indicating the presence of fourteen phytochemical compounds.




Fig. 1: The GC - MS Chromatogram of ethanol extracts of leaves of *Corchorus aestuans*

The identification of the phytochemical compounds was confirmed based on the peak area, retention time and molecular formula. The active principles with their retention time (RT), molecular formula (MF), molecular weight (MW) and peak area in percentage (%) were tabulated in Table 1. The components corresponding to the peaks were determined as follows 3,7,11,15-tetramethyl-2-hexadecen-1-ol (5.67%), Trans-2-undecen-1-ol (1.26%), E-7-Tetradecenol (1.97 %), n-hexadecanoic acid (25.82 %), Phytol (22.34 %), 9,12,15-octadecatrienoic acid, methyl ester, (Z,Z,Z)- (20.23 %), Docosanoic acid, ethyl ester (1.99 %), 1-Eicosanol (2.11 %), 9,9-dimethoxybicyclo[3.3.1]nona-2,4-dione (0.60 %), Heptadecanoic acid, heptadecyl ester (0.95 %), Pentadecanoic acid, 2,6,10,14-tetramethyl-, methyl ester (0.91 %), 3-hexadecyloxy carbonyl-5-(2-hydroxyethyl)-4- methylimidazolium ion (0.90 %), Squalene (8.03 %), Vitamin E (7.24%).

The spectrum sketch out of GC-MS confirmed the presence of fourteen components with the retention time 10.79, 11.04, 11.22, 12.47, 13.91, 14.58, 14.89, 16.70, 17.55, 18.55, 20.29, 21.65, 23.18, 27.39 min respectively which is shown in Figure 1. The phytochemical compounds recognized through GC-MS analysis showed many biological activities are listed in Table 2. The biological activities listed are based on Dr. Duke's Phytochemical and Ethnobotanical Databases created by Dr. Jim Duke of the Agricultural Research Service/USDA.

Table 1: Phytochemical compounds identified in ethanol extract of leaves of *Corchorus aestuans*

S. No.	RT	Name of the compound	Molecular Formula	Molecular Weight	Peak Area %
1.	10.79	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C <sub>20</sub> H <sub>40</sub> O	296	5.67
2.	11.04	trans-2-Undecen-1-ol	C <sub>11</sub> H <sub>22</sub> O	170	1.26
3.	11.22	E-7-Tetradecenol	C <sub>14</sub> H <sub>28</sub> O	212	1.97
4.	12.47	n-Hexadecanoic acid	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256	25.82
5.	13.91	Phytol	C <sub>20</sub> H <sub>40</sub> O	296	22.34
6.	14.58	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	292	20.23
7.	14.89	Docosanoic acid, ethyl ester	C <sub>24</sub> H <sub>48</sub> O <sub>2</sub>	368	1.99
8.	16.70	1-Eicosanol	C <sub>20</sub> H <sub>42</sub> O	298	2.11
9.	17.55	9,9-Dimethoxybicyclo[3.3.1]nona-2,4-dione	C <sub>11</sub> H <sub>16</sub> O <sub>4</sub>	212	0.60
10.	18.55	Heptadecanoic acid, heptadecyl ester	C <sub>34</sub> H <sub>68</sub> O <sub>2</sub>	508	0.95
11.	20.29	Pentadecanoic acid, 2,6,10,14-tetramethyl-, methyl ester	C <sub>20</sub> H <sub>40</sub> O <sub>2</sub>	312	0.91
12.	21.65	3-Hexadecyloxy carbonyl-5-(2-hydroxyethyl)-4- methylimidazolium ion	C <sub>24</sub> H <sub>45</sub> N <sub>2</sub> O <sub>3</sub>	409	0.90
13.	23.18	Squalene	C <sub>30</sub> H <sub>50</sub>	410	8.03
14.	27.39	Vitamin E	C <sub>29</sub> H <sub>50</sub> O <sub>2</sub>	430	7.24

Table 2: Biological activities of phytochemical compounds identified in ethanol extract of leaves of *Corchorus aestuans*

S. No.	Name of the compound	Compound Nature	**Biological Activity
1.	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	Terpene Alcohol	Anti microbial, Anti-inflammatory activity
2.	trans-2-Undecen-1-ol	Alcohol	No activity reported
3.	E-7-Tetradecenol	Alcohol	Anti microbial activity
4.	n-Hexadecanoic acid	Palmitic acid	Antioxidant, Hypocholesterolemic
5.	Phytol	Diterpene	Nematicide, Pesticide, Lubricant, Antiandrogenic Hypocholesterolemic, Antimicrobial, Anticancer, Cancer preventive, Diuretic Anti inflammatory
6.	9,12,15-Octadecatrienoic acid, methyl ester, (Z,Z,Z)-	Fatty acid ester compound	Anti inflammatory, Hypocholesterolemic, Cancer preventive, Hepatoprotective, Nematicide, Insectifuge
7.	Docosanoic acid, ethyl ester	Behenic acid ethyl ester	No activity reported
8.	1-Eicosanol	Alcohol	Anti microbial activity
9.	9,9-Dimethoxybicyclo[3.3.1]nona-2,4-dione	Ketone	No activity reported
10.	Heptadecanoic acid, heptadecyl ester	Fatty acid	No activity reported
11.	Pentadecanoic acid, 2,6,10,14-tetramethyl-, methyl ester	Ester compound	No activity reported
12.	3-Hexadecyloxy carbonyl-5-(2-hydroxyethyl)-4- methylimidazolium ion	Carbonyl	No activity reported
13.	Squalene	Triterpene	Antibacterial, Antioxidant, Pesticide, Antitumor, Cancer preventive, Immunostimulant, Chemo preventive, Lipoxygenase-inhibitor
14.	Vitamin E	Vitamin	Antiageing, Analgesic, Antidiabetic, Anti inflammatory, Antioxidant, Antileukemic,

\*\*Source: Dr. Duke's Phytochemical and Ethnobotanical Databases

The individual fragmentation patterns of necessary components were illustrated in Figures A-J.




Fig. A-J): The individual fragmentation pattern of the important compounds

## CONCLUSION

In the present study, fourteen phytochemical constituents have been identified from the ethanol extract of leaves of *Corchorus aestuans* by Gas Chromatogram - Mass Spectrometry (GC - MS) analysis. The presence of these phytochemical constituents justifies the use of this plant for various ailments by traditional practitioners. Isolation of individual photochemical constituents and subjecting it to biological activities are being undertaken.

## CONFLICT OF INTERESTS

Declared None

## ACKNOWLEDGEMENT

The authors express their deep sense of gratitude to the University Grants Commission, New Delhi, for financial assistance and Dr. S. Kumaravel, Senior Scientist, Indian Institute of Crop Processing Technology, Thanjavur, India for providing facilities for GC-MS studies

## REFERENCES

1. Jain SK. Medicinal plants. National Book Trust; India: 1979. p. 1.
2. Lalitha Eswari M, Vijaya Bharathi R, Jayshree N. Preliminary Phytochemical Screening and Heavy Metal Analysis of Leaf

- Extracts of *Ziziphus oenopia* (L) Mill. Gard. Int J of Pharm Sci and Drug Res 2013;5(1):38-40.
3. Savithamma N, Linga Rao M, Beenaprabha. Phytochemical studies of *Dysophylla myosuroides* (Roth.) Benth. In. wall and *Talinum cuneifolium* (Vahl.) Willd. Res J Phyto (2011);5(3):163-9.
  4. Milne A. Inhalational and local anesthetics reduce tactile and thermal responses in *Mimosa pudica* linn. Masui 1993;1190-3.
  5. Vuorela P, Leinonen M, Saikku P, Tammela P, Rouha JP, Wennberg T, et al. Curr Med Chem 2004;11:1375- 89.
  6. Karimi E, Jaafar HZE. HPLC and GC-MS determination of bioactive compounds in microwave obtained extracts of three varieties of *Labisia pumila* benth. Molecules 2011;16:6791- 805.
  7. Gamble JS. Flora of the presidency of Madras, Vol. 1, Bishen Singh Mahendra Pal Singh 23 - A, New Connaught Place, Dehradun (India). Published under the Secretary of State for India in Council 4. s 1997.
  8. CP Khare (Ed. ). Indian Medicinal Plants An Illustrated Dictionary
  9. Harborne JB. Phytochemical methods. In Chapman &, Hall. New York 1984;11:4-5.
  10. Kalimuthu K. Prabakaran R. Preliminary phytochemical screening and GC-MS analysis of Methanol extract of *ceropegia pusilla*. Int J of Res in Applied Natural and Social Sci 2013;1 (3):49-58.