

Short Communication

ANTIMICROBIAL ACTIVITY SCREENING OF *M. MINUTA* EXTRACTS

BINDU A. R¹, ROSEMARY J¹, AKHILA. S*²

¹Department of pharmaceutical Sciences, Mahatma Gandhi university, Cheruvandoor campus, ²Department of pharmaceutical Sciences, RIMSR, rubber board P. O, Mahatma Gandhi university, Kottayam.
Email: akhilamadathil@yahoo.com

Received: 14 Aug 2014 Revised and Accepted: 15 Sep 2014

ABSTRACT

Objective: Infectious diseases are a leading cause of global disease burden with high morbidity and mortality especially in the developing world. Furthermore, there have been threats of new diseases during the past three decades due to the evolution and adaptation of microbes and the re-emergence of old diseases due to the development of antimicrobial resistance and the capacity to spread to new geographic areas. Since antiquity, humans have used plants to treat common infectious diseases and some of these traditional medicines are still included as part of the habitual treatment of various maladies. Objective of the study was to evaluate, one of the traditionally used drugs, *Marselia minuta* for its antimicrobial activity.

Methods: Extracts of *M. minuta* was prepared using successive solvent extraction method and the antibacterial activity of thus prepared petroleum ether, chloroform, ethyl acetate, alcohol and aqueous extracts of *M. minuta* was performed using agar well diffusion method. Four bacterial strains were used to assess the antimicrobial properties of the plant extracts viz *Escherichia coli*, *Streptococcus pneumonia*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

Results: All the extracts showed activity towards the tested microorganisms except petroleum ether extract. Ethyl acetate extract was more active towards *S. aureus* and *P. aeruginosa*. Alcohol and chloroform extract was active toward *E. coli* and *P. aeruginosa*. Aqueous extract was active against *E. coli* and *S. aureus*. minimum inhibitory concentration were calculated for each extract and the Significant activity was observed for ethyl acetate extract (*E. coli*-11.33 mm, *S. aeruginosa*-16.0 mm) and alcoholic fractions (*E. coli*-15.33 mm and *P. aeruginosa*-10.66 mm) for concentration 500 mcg/ml.

Conclusion: Thus, the result in the present experiment ensures the traditional use of *M. minuta* as antibacterial agent. Further, the most active extracts can be subjected for isolation of the therapeutic antimicrobials and its formulation which will be relatively safer than synthetic medicines and offering profound therapeutic benefits at the more affordable price.

Keywords: *M. minuta*, Antimicrobial activity, Infectious disease, Traditional drugs.

Infectious diseases, also known as transmissible diseases or communicable diseases comprise clinically evident illness, resulting from the infection, presence and growth of pathogenic biological agents in an individual host organism. Infectious pathogens include some viruses, bacteria, fungi, protozoa, multi cellular parasites, and aberrant proteins known as prions. Worldwide, infectious disease is the number one cause of death accounting for approximately one-half of all deaths in tropical countries [1]. Death from infectious disease, ranked 5th in 1981, had become the 3rd leading cause of death in 1992, an increase of 58%.

There has been remarkable progress in the prevention, control and even eradication of infectious diseases with improved hygiene and development of antimicrobials and vaccines. However, infectious diseases still remain a leading cause of global disease burden with high morbidity and mortality especially in the developing world. Furthermore, there have been threats of new diseases during the past three decades due to the evolution and adaptation of microbes and the re-emergence of old diseases due to the development of antimicrobial resistance and the capacity to spread to new geographic areas [2].

Since antiquity, humans have used plants to treat common infectious diseases and some of these traditional medicines are still included as part of the habitual treatment of various maladies. The isoquinoline alkaloid emetine obtained from the underground part of *Cephaelis ipecacuanha*, and related species, has been used for many years as an amoebicidal drug as well as for the treatment of abscesses due to the spread of *Escherichia histolytica* infections. Another important drug of plant origin with a long history of use is quinine. This alkaloid occurs naturally in the bark of *Cinchona* tree is used as antimalarial drug. It is estimated that today, plant materials are present in, or have provided the models for 50% Western drug. The

increasing failure of chemotherapeutics and antibiotic resistance exhibited by pathogenic microbial infectious agents has led to the screening of several traditionally used medicinal plants for their potential antimicrobial activity [3]. *Marsilea* is a genus of approximately 65 species of aquatic ferns of family marsileaceae and in ayurveda, some of these species are recommended for treatment of psychopathy, diarrhea, cough, bronchitis and skin infections [4]. Hence, in the present study one of the traditionally used, yet not scientifically investigated, species, *Marsilea minuta* was screened for its antimicrobial activity.

Extraction of whole plant of *Marsilea minuta* was carried out using solvents of increasing polarity by hot continuous extraction method using soxhlet apparatus. 200 g of shade dried plant were taken and size reduced, extracted with 1L of petroleum ether in the round bottom flask and extraction was continued for 10 hrs. The extract obtained was collected and concentrated. The concentrated extract was then weighed and stored. Then the residue obtained after extraction was packed into the body of the soxhlet and then extracted with chloroform, ethyl acetate and ethanol in the same manner. Each time before extracting with the next solvent, the drug material was dried at room temperature. Each extract was concentrated by distilling off the solvent and then evaporating to dryness.

Finally the marc obtained was packed into a round bottom flask and refluxed with water. The extract obtained was concentrated in vacuum drier to obtain aqueous extract. All the extracts were weighed and stored for further studies [5]. All the extracts were subjected to antibacterial study using well diffusion method. Mueller Hinton Agar Medium (MHA) was used for the study [6,7]. Dehydrated medium supplied by HI media laboratories Pvt. Ltd., Mumbai were used for well diffusion studies for testing antibacterial activity of the plant extracts. The reconstituted medium has the following composition.

Beef infusion -300 ml, Casein hydrolysate -17.4 gm, starch -10 gm, Agar -10 gm and Distilled water -1000 ml pH -7.4

Four bacterial strains were used to assess the antimicrobial properties of the plant extracts. The isolates were procured from local clinical laboratories and was identified according to guidelines in the Bergey's Manual of Systematic Bacteriology and by studying their culture, morphological and biochemical properties. Bacterial strains tested were confirmed as *Escherichia coli*, *Streptococcus pneumonia*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. The wells were made using a 5 mm cork borer sterilized using flame and alcohol. The bacterial isolates were first grown in a nutrient broth for 18 hrs before use. A sterile swab was dipped into nutrient broth culture and was rotated firmly against upper inside wall of the tube to expel the excess fluid. The entire surface of Mueller Hinton agar medium was lawned with the swab soaked with inoculums. As soon as the MHA was partly solidified, the plates were inverted and left for 2 hrs. When cooled, wells were made on the solidified medium in the plate. 100 µl of the varying concentrations (100, 200, 300, 400 and 500 µg/ml) of the

extracts were introduced into the wells with the help of micropipettes.

The plates were incubated over night at 37°C. Microbial growth was determined by measuring the diameter of zone of inhibition. For each bacterial strain, controls maintained where DMSO was used instead of the extract. The experiment was done in triplicate and the results obtained by measuring the zone diameter.

The minimum inhibitory concentration of all extracts was found out by graphical method [7]. By successive solvent extraction method, petroleum ether extract (PEE), chloroform extract (CHE), ethyl acetate extract (EAE), alcoholic extract (ALE), and aqueous extract (AQE) of *M. minuta* were prepared. Antibacterial activity of petroleum ether, chloroform, ethyl acetate, alcohol and aqueous extracts of *M. minuta* were determined and experiments were done in triplicate and the results were expressed as mean standard deviation. The results obtained were tabulated in the table 1 and the minimum inhibitory concentration obtained for each extract was tabulated in table 2. The photographs of the plates were depicted in the figures 1 and 2.

Table 1: Zone of inhibition obtained for different extracts of *M. minuta* against four microorganisms

Extracts	Microorganism	Zone of inhibition of various extracts (mm) at different concentration				
		100mcg/ml	200mcg/ml	300mcg/ml	400mcg/ml	500mcg/ml
PEE	<i>E. coli</i>	-	-	-	-	-
	<i>S. aureus</i>	-	-	-	-	-
	<i>P. aeruginosa</i>	-	-	-	-	-
	<i>S. pneumonia</i>	-	-	-	-	-
CHE	<i>E. coli</i>	-	-	6.33±0.58	8.0± 1.0	8.66±0.57
	<i>S. aureus</i>	-	-	-	7.66±0.577	8.0± 1.0
	<i>P. aeruginosa</i>	-	-	6.33±0.58	8.66±5.8	10.33±0.58
	<i>S. pneumonia</i>	-	-	-	-	5.33±0.58
EAE	<i>E. coli</i>	-	7.33±0.58	8.33±0.58	10.0±1.0	11.33±0.58
	<i>S. aureus</i>	8.66±0.5	9.66±0.58	10.0±1.0	13.33±0.58	14.33±0.58
	<i>P. aeruginosa</i>	8±1.0	8.66±0.58	11.0±1.0	13.33±0.58	16.0±1.0
	<i>S. pneumonia</i>	7.33±0.5	8±1.0	10.66±0.5	13.66±0.5	17.66±0.58
ALC	<i>E. coli</i>	-	6.33±0.58	9.0±1.0	12.33±0.58	15.33±0.57
	<i>S. aureus</i>	-	6.33±0.57	7.66±0.58	9.66±0.57	12.33±0.58
	<i>P. aeruginosa</i>	-	5.66±0.58	8±1.0	9.33±0.58	10.66±0.58
	<i>S. pneumonia</i>	-	-	6.66±0.58	8.33±0.58	10±1.00
AQE	<i>E. coli</i>	6.66±0.5	9±1.0	9.66±0.58	11.66±0.58	12±1.00
	<i>S. aureus</i>	8±1.0	-	8±1.00	9.33±0.58	10.00±0.58
	<i>P. aeruginosa</i>	-	-	-	6.66±0.58	8.66±0.50
	<i>S. pneumonia</i>	-	-	-	-	6.0± 1.0
STD	Zone of	inhibition of standard (mm) at different concentration				
	Microorganism	5mcg/ml	10 mcg/ml	20 mcg/ml	40 mcg/ml	80 mcg/ml
Strepto-	<i>E. coli</i>	-	-	7.00±1.00	7.66±0.58	9.00±0.57
	<i>S. aureus</i>	6.33±0.5	7±1.00	7.66±0.57	8.66±0.57	9.66±0.58
myc-in	<i>P. aeruginosa</i>	7	6.66±0.58	7.00±1.00	8.33±0.58	9.33±0.58
	<i>S. pneumonia</i>	-	-	7.33±0.58	8.66±0.57	9.00±1.00
<i>Values are expressed as</i>		<i>mean±S. D; n=3</i>				

Table 2: Minimum inhibitory concentration obtained for each extract against four microorganisms.

Micro organism	MIC (mcg/ml)					
	PEE	CHE	EAE	ALC	AQE	STD
<i>E. coli</i>	>500	275.8	185.3	141.6	94.19	14.10
<i>S. aureus</i>	>500	≤400	97.94	142.3	223.7	5.22
<i>P. aeruginosa</i>	>500	295.3	92.3	136.87	≤ 400	8.69
<i>S. Pneumonia</i>	>500	≤500	136.6	289.26	≤ 400	19.85

All the extracts showed activity towards the tested microorganisms except petroleum ether extract. Ethyl acetate extract was more active towards *S. aureus* and *P. aeruginosa*. Alcohol and chloroform extract was active toward *E. coli* and *P. aeruginosa*. Aqueous extract was active against *E. coli* and *S. aureus*.

The ethyl acetate extract was most potent against the tested organisms with an MIC range of 90-190 mcg/ml. The lowest MIC (highest potency) exhibited was 92.3mcg/ml against *S. aureus*. Ethyl acetate and alcoholic

extracts showed good response on both gram-positive and gram negative bacteria. Although Gram negative bacteria tend to have higher intrinsic resistance to most antimicrobial agents, in spite of this, impressive activity was observed for ethyl acetate extract (*E. coli*-11.33 mm, *S. aureus*-16.0 mm) and alcoholic fractions (*E. coli*-15.33 mm and *P. aeruginosa*-10.66 mm) for concentration 500 mcg/ml.

The ethyl acetate extract was found to be active against *S. aureus* and *P. aeruginosa* which are the causative organisms for skin infections;

therefore this activity can be a scientific proof for the traditional use of the plant [8,9]. The plant extracts were also found to be active against *S. pneumonia* which cause respiratory tract infections, thus providing a scientific basis for the use of plant paste in cough and respiratory infections.



(1-Chloroform extract, 2- Ethyl acetate extract, 3- Alcoholic extract, 4- Aqueous extract. All extracts were taken in concentration 500 mcg/ml)

Fig. 1: Anti microbial activity obtained for different extracts against *E. coli* (left) and *S. aureus* (right)




Fig. 2: Anti microbial activity obtained for different extracts against *S. pneumonia* (left) and *P. aeruginosa* (right)

The increasing failure of chemotherapeutics and antibiotic resistance exhibited by pathogenic microbial infectious agents has led to the screening of several traditionally used medicinal plants for their potential antimicrobial activity as they will be safer, effective and cheap [3,10,11]. Thus, the result in the present experiment also ensures the traditional usage of *M. minuta* as antibacterial agent.

Further, the most active extracts can be subjected for isolation of the therapeutic antimicrobials and its formulation which will be relatively safer than synthetic medicines and offering profound therapeutic benefits at the more affordable price.

REFERENCES

1. Das K, Tiwari RKS, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agent: current methods and future trends. *J Medicinal Plant Res* 2010;4(2):104-11.
2. Dharamuir H. Bioactive Medicinal plants. Gene tech books: New Delhi; 2002;56:57-75.
3. Iwu MM, Duncan AR, Okunji CO. New Antimicrobials of Plant Origin. Perspectives on new crops and new uses. ASHS Press: Alexandria; 1999. p. 457-62.
4. Tiwari OP, Subhrata K, Bhattamisra, Pushpendra K, tripathi, Paras N. Anti aggressive activity of standardized extract of *Marsilea minuta* Linn. In rodent models of aggression. *Bio Sci Trends* 2010;4(4):190-4.
5. Pulok KM. Quality control of herbal drugs. 2nd ed. New Delhi: Business Horizons; 2007.
6. Mohana DC, Satish S, Raveesha KA. Antibacterial evaluation of some plant extracts against some human pathogenic bacteria. *Adv Biol Res* 2008;2:49-55.
7. Bonev B, Hooper J, Parisot J. Principles of assessing bacterial susceptibility to antibiotics using the agar diffusion method. *J Antimicrobial Chemotherapy* 2008;61(6):1295-301.
8. Spellberg B, Powers JH, brass EP, Miller LG, Edwards JR. trends in antimicrobial drug development: implications for the future. *Clin Infect Dis* 2004;38:1279-86.
9. Yuan Q, Wang J, Ruan J. Screening for bioactive compounds from *adiantum capillus- veneris* L. *J Chem Soc Pak* 2012;34(1):207-16.
10. Igbiosa OO, Igbiosa EO, Aiyegoro OA. Antimicrobial activity and phytochemical screening of stem bark extracts from *Jatropha curcas* (Linn). *Afr J Pharm Pharmacol* 2009;3(2):58-62.
11. Neeraj M, Behal KK. Antimicrobial activity of some species against selected microbes. *Int J Pharm Pharm Sci* 2010;2(3):187-96.