

Original Article

IDENTIFICATION, QUANTIFICATION AND VALIDATION OF STIGMASTEROL FROM *ALPINIA CALCARATA* USING HIGH-PERFORMANCE THIN LAYER CHROMATOGRAPHY METHOD

PRATHEEMA PHILOMINDOSS

Department of Chemistry, Holy Cross College, Tiruchirappalli 620002, India
Email: benprathee1021@gmail.com

Received: 15 May 2017 Revised and Accepted: 13 Jul 2017

ABSTRACT

Objective: The present study is designed to develop a new simple, precise, rapid and selective high-performance thin-layer chromatographic (HPTLC) method for the determination of stigmasterol in methanolic rhizomes extract of *Alpinia calcarata*.

Methods: As per International Conference on Harmonization (ICH) guidelines we have applied different concentrations of stigmasterol as standard on HPTLC plates for the quantification of stigmasterol from the *Alpinia calcarata* rhizomes. The concentration of standard stigmasterol is 1 mg/ml.

Results: The retention factor of stigmasterol was 0.58. Linearity was obtained in the range of 50 ng-250 ng for stigmasterol. The developed and validated HPTLC method was employed for stigmasterol in methanolic rhizomes extract of *Alpinia calcarata* for standardization of the content of the marker. The linear regression data for the calibration plots showed a good linear relationship with $r=0.99977$ for stigmasterol, respectively Satisfactory recoveries of 99.77 % were obtained for stigmasterol.

Conclusion: The results obtained in validation assays indicate the accuracy and reliability of the developed HPTLC method for the quantification of stigmasterol in methanolic rhizomes extract of *Alpinia calcarata*

Keywords: *Alpinia calcarata* Roscoe, Stigmasterol, HPTLC

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)
DOI: <http://dx.doi.org/10.22159/ijpps.2017v9i9.19988>

INTRODUCTION

Alpinia calcarata Roscoe belonging to the family Zingiberaceae is a rhizomatous perennial herb, which is commonly used in the traditional medicinal systems in Bangladesh, India, Indonesia, Thailand, Malaysia, Sri Lanka, Taiwan, Cambodia, and Vietnam [1]. *Alpinia calcarata* is used in the oriental part of the world as a food additive, spice and in indigenous system of medicine. It grows in dense forests at high altitudes and is considered as a native of India [2]. *Alpinia calcarata* rhizomes are branched and dense with a light to dark brown colour and known to possess a broad spectrum of medicinal properties. This is a very good source of pinocembrin 6 which induces mitochondrial apoptosis in colon cancer cells [3].

Experimentally, rhizomes are shown to possess antibacterial [4], antifungal [5], anthelmintic, antinociceptive [6], antioxidant [7], aphrodisiac [8], antidiabetic activities [9], rheumatism, fever and anticancer activity. It is also widely used to relieve colds and reducing swellings [10, 11]. High blood pressure, diuretic, stomach problems, analgesic [12], anticandidal, antiplatelet, antispasmodic [13] antiulcerous hypotensive [14, 15] insecticidal, muscle relaxant and Uterine stimulant.

Nowadays, HPTLC has become a routine analytical technique due to its advantages of reliability in quantitation of analytes at the micro and even in nanogram levels and cost effectiveness [16]. It has proved a very useful technique because of its low operating cost, high sample throughput and the need for minimum sample clean-up. The major advantage of HPTLC is in reducing analysis time and cost per analysis [17]. However, no single method was found in the literature to our knowledge to detect stigmasterol in methanolic extract of rhizomes of *Alpinia calcarata*. Therefore, the present study aimed to develop an HPTLC method for analysis of stigmasterol in *Alpinia calcarata* rhizomes which have not yet been reported.

The developed method was optimized and validated in accordance with International Conference on Harmonization (ICH) guidelines. Here, we developed a simple high-performance thin-layer chromatographic method for the rapid analysis of stigmasterol

compounds in *Alpinia calcarata* rhizomes. The method was found suitable for rapid screening of plant materials for their genotypic assessment and can be performed without any special sample pretreatment.

MATERIALS AND METHODS

Materials

Standard and chemicals

Stigmasterol (purity 99%), was purchased from Sigma-Aldrich, New Delhi. Ethyl acetate, toluene, glacial acetic acid and methanol used in the present research work were of HPLC grade and were procured from E. Merck Mumbai, India.

Collection of plant material

The Rhizomes of *Alpinia calcarata* were collected in the Month of July from the Kolli Hill-Chinna Kovilur, Tamil Nadu, India. The plant was identified and rhizomes of *Alpinia calcarata* were authenticated and confirmed from Dr. S. John Britto, Director, Rapinat herbarium, St. Joseph College, Tiruchirappalli, and Tamil Nadu for identifying the plants. The voucher specimen number PP001 (14.07.2016).

Preparation of methanol extracts

The rhizomes of *Alpinia Calcarata* were washed in running water, cut into small pieces and then shade dried for a week at 35-40 °C, after which it was grinded to a uniform powder of 40 mesh size. The methanol extracts were prepared by soaking 100 g each of the dried powder plant materials in 1 L of methanol using a soxhlet extractor continuously for 10 hr. The extracts were filtered through whatmann filter paper No. 42 (125 mm) to remove all unextractable matter, including cellular materials and other constitutions that are insoluble in the extraction solvent. The entire extracts were concentrated to dryness using a rotary evaporator under reduced pressure. The final dried samples were stored in labeled sterile bottles and kept at -20 °C. The filtrate obtained was used as sample solution for the further HPTLC analysis [17].

Identification and quantification of stigmaterol from *Alpinia calcarata* by HPTLC

Sample preparation

All the chemicals, including solvents, were of analytical grade from E. Merck, India. The HPTLC plates Si 60F₂₅₄ (20 cmX10 cm) were purchased from E. Merck (India). Standards of stigmaterol (99% purity) were purchased from Sigma (New Delhi, India). 100 mg/ml of methanolic extracts of rhizomes of *Alpinia calcarata* was taken for analysis. The extracts were filtered and vacuum dried at 45 °C. The dried extracts were separately redissolved in 1 ml of methanol and sample of varying concentration (5-30 µl) for stigmaterol were spotted for quantification. 1 mg of standard stigmaterol were prepared in 1 ml of chloroform, and different amounts of (50-250 ng) stigmaterol were loaded onto a TLC plate to get the calibration curve [18-21].

Thin layer chromatography

A Camag HPTLC system equipped with an automatic TLC sampler ATS₄, TLC scanner 3 and integrated software Win CATS version 3 was used for the analysis. Samples were washed on a precoated silica gel HPTLC plates Si 60F₂₅₄ (20 cm x 10 cm) plate of 200 µm-layer thickness, for quantification of stigmaterol in rhizomes of *Alpinia calcarata*. The samples and standards were applied on the plate as 8 mm wide bands with a constant application rate of 150Nl s⁻¹, with an automatic TLC sampler (ATS₄) under a flow of N₂ gas, 15 mm from the bottom, 15 mm from the side, and the space between two spots was 6 mm in the plate.

Detection and estimation of stigmaterol

The linear ascending development was carried out in a camag twin through chamber (20 cm x 10 cm), which was pre-saturated with a 25 ml mobile phase, toluene: ethyl acetate (8.0: 2.0 v/v) for stigmaterol for 30 min, at room temperature (25 °C±2 °C) and 50±5% relative humidity. The length of the chromatogram run was up to 90 mm. Subsequent to the development; the TLC plate was dried in a current of air, with the help of air dryer, in a wooden chamber with adequate ventilation.

The dried plate was dipped into freshly prepared libermann burchard reagent. Quantitative estimation of the plate was performed in the absorption-reflection mode at 538 nm, using a slit width 6.00 x 0.45 mm, with data resolution 100 µm/step and scanning speed 20 mm/sec. The source of radiation utilized was a tungsten lamp emitting continuous visible spectra of 366 nm. Determination of stigmaterol in methanolic extracts of rhizomes of *Alpinia calcarata* was performed by the external standard method, using pure standards. Each was carried out in triplicate [22, 23].

Method validation

This method was validated as per the ICH guidelines (International Conference on Harmonization in 1994, 1996 and 2005), the method validation parameters checked were linearity, precision, accuracy and recovery, limit of detection, limit of quantification, specificity, robustness and ruggedness. All measurements were performed in triplicates [24-26].

Calibration curve and linearity

The calibration was performed by analysis of working standard solutions of stigmaterol (50 to 250 ng for *Alpinia calcarata*) were spotted on precoated TLC plate, using semi automatic spotter under a nitrogen stream. The TLC plates were developed, dried by hot air and photometrically analyzed as described earlier. The calibration curves were prepared by plotting peak area versus concentration (ng/spot) corresponding to each spot.

Recovery

To determine the recovery, known concentrations of standards were added to a pre-analyzed sample of *Alpinia calcarata* rhizomes. The spiked samples were then analyzed by the proposed HPTLC method and the analysis was carried out in triplicate.

Precision

A stock solution containing stigmaterol compound was prepared in chloroform and six 10 µl (1000 ng/spot) bands were applied and analyzed by the developed method to determine instrument precision. Six different volumes of same concentration were spotted on a plate and analyzed by the developed method to determine variation arising from the method itself. To evaluate intra-day precision, six samples at three different concentrations (1000, 2000 and 3000 ng/spot) for stigmaterol were analyzed on the same day. The inter-day precision was studied by comparing assays performed on three different days.

Limit of detection and limit of quantification

The detection limit (LOD) of an individual analytical procedure is the lowest amount of analyte in a sample which can be detected but not necessarily quantitated as an exact value. LOD was calculated using the following formula

$$\text{LOD} = \frac{3.3 \times \text{standard deviation of the } y - \text{intercept}}{\text{slope of calibration curve}}$$

The quantification limit (LOQ) of an individual analytical procedure is the lowest amount of analyte in a sample which can be quantitatively determined with suitable precision and accuracy. LOQ was calculated using the following formula,

$$\text{LOQ} = \frac{10 \times \text{standard deviation of the } y - \text{intercept}}{\text{slope of calibration curve}}$$

Specificity

The specificity of the method was ascertained by analyzing standard compound stigmaterol and the compound stigmaterol is present in the rhizomes of *Alpinia calcarata*.

Method specifications

Silica gel 60 F254 precoated plates (20x 10 cm) were used with toluene: ethyl acetate (8.0:2.0 v/v) for stigmaterol as solvent system. The sample was spotted on precoated TLC plates by using Linomat 5 applicator. Ascending mode was used for development of thin layer chromatography. TLC plates were developing up to 80 mm and scanned in fluorescence mode at 366 nm. The contents of stigmaterol in the *Alpinia calcarata* were determined by comparing the area of the chromatogram of standard stigmaterol with a calibration curve of the marker compound of *Alpinia calcarata*, considering the isolated compound to be 100% pure.

RESULTS

Optimization of HPTLC chromatographic conditions

HPTLC fingerprint patterns have been therefore evolved for methanolic extracts of rhizomes of *Alpinia calcarata*. Stigmaterol standard was quantitated accurately using silica gel F₂₅₄ HPTLC pre-coated plates with the mobile phase toluene: ethyl acetate (8.0: 2.0 v/v), the R_f value was about 0.58. The chromatographs of stigmaterol and methanolic extract of rhizomes of *Alpinia calcarata* are shown in (fig. 1). The R_f value of stigmaterol was matched with the R_f value of extract was about 0.58 was shown in peak (fig. 2 (a) and (b)). A steroid compound stigmaterol was identified and isolated by HPTLC techniques.

Validation of HPTLC method

Calibration curve and linearity

The calibration curve was prepared by plotting peak area versus concentration (ng/spot) corresponding to each spot (fig. 3). The regression equation and correlation curves for stigmaterol in *Alpinia calcarata* were regression via height $y = 6.234 + 0.101X$ and $r = 0.99315$ $sdv = 6.52$. Fig. 3 (a) and regression via area $y = 132.296 + 2.085 X$ and $r = 0.99977$ $sdv = 1.17$. fig. 3(b).

Accuracy and recovery

The results showed that the percentage recoveries after sample processing and application were in the range of 99.77 % to 100.11 % (stigmaterol) (table 1). The percentage of stigmaterol in *Alpinia calcarata* rhizome (table 2).




Fig. 1: Quantitative estimation of stigmasterol in *Alpinia calcarata* rhizomes




Fig. 2: (a) HPTLC chromatogram of standard stigmasterol; (b) HPTLC chromatogram of stigmasterol in *Alpinia calcarata* rhizomes




Fig. 3 Linear graph for Stigmasterol in *Alpinia calcarata* in all tracks (concentration vs. area)

Precision

The developed method was found to be precise as indicated by percent RSD (Relative Standard Deviation) not more than 1.5 (tables 3 and 4).

Specificity

It was observed that the other herbal constituents present in the formulations did not interfere with the peak of stigmasterol.

Therefore the method was specific. The spectrum of standard compound stigmasterol and the corresponding spot present in *Alpinia calcarata* matched exactly, indicating no interference by the other plant constituents and excipients. The peak purity of stigmasterol was assessed by comparing the spectra at three different levels like peak start (S), peak apex (M) and peak end (E) positions of the spot. Good correlation $r = 0.99977$ and $SD = 1.17$ for stigmasterol were obtained between the standard and sample overlain spectra of stigmasterol (fig. 4).




Fig. 4: Spectral comparison of standard Stigmasterol (green colour) and Stigmasterol quantified from *Alpinia calcarata* rhizomes (pink colour)

Table 1: Recovery study of stigmasterol by HPTLC (n=3)

Compound	Amount of compound present in the plant material (mean, mg/100 mg)	Amount of standard added (mg)	Amount of standard found in the mixture (mg)	Recovery (%)
Stigmasterol	1912	1912	3830.31	100.11±1.14
		3824	5725.33	99.77±0.93

n is number of determination

Table 2: Amount of Stigmasterol in *Alpinia calcarata* rhizomes (n=3)

Compound	Quantity (mean) (mg/100 mg)	mean±SE	CV (%)
Stigmasterol	1.912	1.912±0.005	0.26

n is number of determination, SE is standard error, CV is cumulative value

Table 3: Intra-day and inter-day precision of the method (n = 6)

Compound	Amount (ng/spot)	Intra-day precision			Inter-day precision		
		Mean area	SD	%RSD	Mean area	SD	% RSD
Stigmasterol	1000	1580.81	1.53	0.096	1540.50	1.67	0.108
	2000	3165.32	1.71	0.054	3215.35	1.98	0.061
	3000	4790.20	1.48	0.030	4681.80	1.84	0.039

n is number of determination, SD is standard deviation, RSD is relative standard deviation

Table 4: Summary of validation parameter

Parameters	Stigmasterol
Linearity	
(i) Range	50-250 ng
(ii) Correlation coefficient	
(a) Height	0.99315
(b) Area	0.99977
(iii) Rf value	0.60
Precision (%RSD)	
(i) Instrument precision (CV%, n=6)	1.62
(ii) Method precision (CV%, n=6)	2.91
LOD (ng/spot)	50
LOQ (ng/spot)	214
Specificity	Specific
Robustness	Robust
Ruggedness (%RSD)	0.8110

n is a number of determination, RSD is a relative standard deviation, CV is cumulative value, LOD is Limit of detection, LOQ is Limit of quantification, Rf is retention factor.

Limit of detection and limit of quantification

The limit of detection was found to be 80 ng/spot for stigmaterol while the limit of quantification was found to be 212 ng/spot for stigmaterol (table 4).

Robustness

Robustness tests examine the effect of the operational parameters on the analysis results. By introducing small changes in mobile phase composition, the results indicated that the method was robust (table 5).

Table 5: Robustness of the method (n=6)

Compound	Amount (ng/spot)	Mobile phase	%RSD
Stigmaterol	1000	Toluene: Ethyl acetate (7: 3v/v)	0.96
		Toluene: Ethyl acetate (8:2 v/v)	1.45

n is number of determination, RSD is relative standard deviation

Ruggedness of the method

It expresses the precision within laboratories variations like different days, different analyst, and different equipment. The ruggedness of the method was assessed by spiking the standard 6 times in two different days with the different analyst (table 4).

DISCUSSION

HPTLC is a simple, rapid and accurate method for analyzing plant material. HPTLC fingerprint has better resolution and estimation of active constituents is done with reasonable accuracy in a shorter time. The HPTLC method can be used for phytochemical profiling of plants and quantification of compounds present in plants. With increasing demand for herbal products as medicines and cosmetics, there is an urgent need for standardization of plant products [16]. HPTLC finger print analysis has become the most of its simplicity and reliability. It can serve as a tool for identification, authentication, qualitative, quantitative analysis and quality control of herbal drug [27]. Thin-layer chromatography (TLC) and HPTLC methods are commonly applied for the identification, assay and testing for purity, stability, dissolution or content uniformity of raw materials (herbal and animal extracts, fermentation mixtures, drugs and excipients) and formulated products (pharmaceuticals, cosmetics, nutriments) [28]. Stigmaterol is reported to show antioxidant [29], anti-inflammatory [30], analgesic and anthelmintic [30], hepatoprotective and anticancer activity [31-35].

Previous study has reported that quantitative analysis of stigmaterol and *dl*- α -tocopherol acetate, two marker compounds in *Leptadenia reticulata* by high-performance thin-layer chromatographic methods [20]. Quantitative determination of stigmaterol and lupeol from *Ficus religiosa* by HPTLC [17]. HPTLC method which has been developed and validated for quantitation of stigmaterol from petroleum ether extract of dried leaf, stem and flower of *Ageratum conyzoides* [36]. Qualitative and quantitative estimation of galangin from *Alpinia galangal* and *Alpinia officinarum* by HPTLC method [37].

Determination of stigmaterol in dietary supplements by gas chromatography was performed [38], and HPTLC determination of stigmaterol was done [39]. Four new labdane-type diterpenoids, calcaratarins A-D (1-4), along with six known labdane-type diterpenoids, a known element-type sesquiterpenoid, and a known coumarin were isolated from the rhizomes of *Alpinia calcarata* [40]. The major components of essential oils isolated from *Alpinia calcarata* rhizomes were camphene (3.86%), beta-myrcene (4.39%), eucalyptol (14.05%), linalol (2.48%), pyrazine (1.72%), L-camphor (7.90%) and berneol (5.67%) might possess significant anti-inflammatory activity and analgesic effect and could be a potential source for treatment of different inflammatory diseases [41]. Phytochemical analysis of the *Alpinia calcarata* rhizomes extracted using soxhlet apparatus with different solvents revealed the presence of polyphenols, tannins, flavonoids, steroid, glycosides and alkaloids [1]. Beta alanine, isovaleric acid, catechol, citral protocatechuic acid, iso menthone, nerol umbelliferone, chrysin, pigenindi-methyl ether, vanillic acid, myristic acid, acetoxycha-vicol acetate, nerolidol, himachalol, dihydro resveratrol, confertin, valerenic acid, alpha-linolenic acid isolated from rhizomes of methanolic extract of *Alpinia calcarata* by LCMS/MS analysis [2].

Isolation of stigmaterol and β -sitosterol from the Rhizomes of *Alpinia allughas* Ros and revealed to the presence of steroids, terpenoids, alkaloids and phenolics [42].

The presented study clearly gave evidence of the bioactive quantitative of stigmaterol in methanolic extracts of *Alpinia calcarata* rhizomes. The developed HPTLC method for the quantification of above stigmaterol compounds is simple, precise, specific, sensitive, and accurate. Further, this method can be effectively used for routine quality control of herbal materials as well as formulations containing any or both of these compounds.

CONCLUSION

In conclusion, an HPTLC method has been developed with some modifications and it can be used for the quantitative determination of stigmaterol in methanolic extract of rhizomes of *Alpinia calcarata*; its main advantages are its simplicity, accuracy and selectivity. The average recovery values of stigmaterol were found to be about 99.94%, which showed the reliability and suitability of the method. This method could also be used for the estimation of these compounds in other herbal preparations and might be useful for standardization purposes.

ACKNOWLEDGEMENT

P. P acknowledges Dr. S. John Britto, Director, rapinat herbarium, St. Joseph College, Tiruchirapalli, Tamil Nadu for identifying the plants. P. P acknowledges Assistant Professor, DR. L. Cathrine of Holy Cross College, Tiruchirapalli, Tamil Nadu for constant support for this research.

AUTHOR CONTRIBUTION

All authors contributed equally to this manuscript.

CONFLICTS OF INTERESTS

The authors declare that they have no conflict of interest. It has not been published elsewhere. That it has not been simultaneously submitted for publication elsewhere. All authors agree to the submission to the journal.

REFERENCES

- Mohanasundari L, Suja S. Qualitative phytochemical screening of rhizomes on *Alpinia calcarata* and *Alpinia speciosa*. J Pharmacogn Phytochem 2015;4 Suppl 2:53-6.
- Silvy Mathew, John Britto S. LCMS/MS analysis of methanolic rhizome extracts of *Alpinia calcarata* roscow (*Zingiberaceae*)-a multipotent medicinal plant. Int J Adv Res 2016;4 Suppl 5:1817-22.
- Kumar MAS, Nair M, Hema PS, Mohan J, Santhoshkumar TR. Pinocebrin triggers Bax-dependent mitochondrial apoptosis in colon cancer cells. Mol Carcinog 2007;46 Suppl 3:231-41.
- George M, Pandalai KM. Investigations on plant antibiotics. Indian J Med Res 1949;37:169-81.
- Pushpangadan P, Atal CK. Ethno-medico-botanical investigations in Kerala. J Ethnopharmacol 1984;111:59-77.
- Arambewela LSR, Arawawala LDAM, Ratnasooriya WD. Antinociceptive activities of aqueous and ethanolic extracts of *Alpinia calcarata* rhizomes in rats. J Ethnopharmacol 2004;95:311-6.

7. Arambewela LSR, Arawwawala LDAM. Antioxidant activities of ethanolic and hot aqueous extracts of *Alpinia calcarata* rhizomes, Aust J Med Herbalism 2005;17:91-4.
8. Ratnasooriya WD, Jayakody JR. Effects of aqueous extract of *Alpinia calcarata* rhizomes on reproductive competence of male rats. Acta Biol Hung 2006;57:23-35.
9. Arambewela LSR, Arawwawala LDAM, Ratnasooriya WD. Hypoglycemic and antihyperglycemic activities of the aqueous and the ethanolic extracts of *Alpinia calcarata* rhizomes in rats. Pharmacogn Mag 2009;5:412-8.
10. Ahmed AD, Medine G, Meryen S, Hatice O, Fikerenttin S, Isa K. Anti-microbial effects of *Ocimum basilicum* (Labiatae) extract. Turkey J Biol 2005;29:155-60.
11. Perveen R, Islam F, Khanum J, Yeasmin T. Preventive effect of ethanolic extract of *Alpinia Calcarata* Rosc on ehrlich's ascitic carcinoma cell induced malignant ascites in mice. Asian Pac J Trop Med 2012;5:121-5.
12. Leal-Cardoso JH. Effects of essential oil of *Alpinia Zerumbet* on the compound action potential of the ratsciatic nerve. Phytomedicine 2004;11 Suppl 6:549-53.
13. Bezerra MA. Myorelaxant and antispasmodic effects of the essential oil of *Alpinia speciosa* rat ileum. Phytother Res 2000;14 Suppl 7:549-51.
14. Lahlou S. Antihypertensive effects of the essential oil of *Alpinia Zerumbet* and its main constituent, terpinen-4-ol, in DOCA-salt hypertensive conscio rats. Fundam Clin Pharmacol 2003;17 Suppl 3:323-30.
15. De Moura RS. Antihypertensive and endothelium dependent vasodilator effects of *Alpinia Zerumbet*, a medicinal plant. J Cardiovasc Pharmacol 2005;46 Suppl 3:288-94.
16. Rathee D, Thanki M, Agarwal R, Anandjiwala S. Simultaneous quantification of bergenin, catechin, gallicin, gallic acid and quantification of β -Sitosterol using HPTLC from *Bergenia ciliate* (Haw). Pharm Anal Acta 2010;1:104.
17. Deepti Rathee, Sushila Rathee, Permender Rathee, Aakash Deep, Sheetal Anandjiwala, Dharmender Rathee. HPTLC densitometric quantification of stigmasterol and lupeol from *Ficus religiosa*. Arab J Chem 2015;8:366-71.
18. Suthar AC, Banavaliker MM, Biyani MK, Priyadarsini Indira K, Sudarsan V, Mohan H. A high performance thin layer chromatography method for quantitative estimation of lupeol in *Crataeva nurvala*. Indian Drugs 2001;38 Suppl 9:474-8.
19. Badami S, Gupta MK, Ramaswamy S, Rai SR, Nanjiam M, Bendell DJ, et al. Determination of betulin in *Grewia itiaefolia* by HPTLC. J Separ Sci 2004;27:129-31.
20. Purnima D, Hamrapurkar PK. HPTLC Determination of stigmasterol and tocopherol acetate in *Leptadenia reticulata* and its formulation. J Plan Chromatogr 2007;20 Suppl 3:183-7.
21. Geneva. ICH, Q2A (R1), Validation of Analytical Procedure: Text and Methodology, International Conference on Harmonization Specification; 2005. p. 5-10.
22. Ravindra CS, Sanjay BK, Kalaichelvan VK. Phytochemical profile studies on the steroids of methanolic leaf extract of medicinally important plant *Holoptelea integrifolia* (Roxb.) planch using high performance thin layer chromatography. Asian J Pharm Clin Res 2014;7:4.
23. Lavanya MS, Gnanamani A, Ilavarasan R. Physico-chemical, phytochemical and high performance thin layer chromatography analysis of the whole plant of *Orthosiphon thymiflorus* (Roth.) sleesen. Asian J Pharm Clin Res 2015;8:1.
24. Geneva. ICH, Q2B, Guideline on Validation of Analytical Procedure: Methodology, International Conference on Harmonization; 1996. p. 1-8.
25. Geneva. ICH, Q2A, Text on Validation of Analytical Procedure, International Conference on Harmonization; 1994. p. 1-5.
26. Sarfaraj Hussain, Sheeba Fareed, Mohammad Ali, Sarfaraz Alam, Akhlakquer Rahman, Srivastava AK. Phytochemical investigation and simultaneous estimation of bioactive lupeol and stigmasterol in *Abutilon indicum* by validated HPTLC method. J Coastal Life Med 2014;2 Suppl 5:394-401.
27. Kpoviessi DSS, Gbaguidia F, Gbénou J, Accrombessia G, Moudachiroua M, Rozet E. Validation of a method for the determination of sterols and triterpenes in the aerial part of *Justicia anselliana* (Nees) T. Anders by capillary gas chromatography. J Pharm Biomed Anal 2008;48:1127-35.
28. Delporte C, Backhouse N, Erazo S, Negrete R, Vidal P, Silva X. Analgesic-anti-inflammatory properties of *Proustia pyrifolia*. J Ethnopharmacol 2005;99 Suppl 11:119-24.
29. McAnuff MA, Harding WW, Omoruyi FO, Jacobs H, Morrison EY, Asemota HN. Hypoglycemic effects of steroidal saponin isolated from *Jamaican bitteryam*, *Dioscorea polygonoides*. Food Chem Toxicol 2005;43 Suppl 11:1667-72.
30. Sunitha S, Nagaraj M, Varalakshmi P. Hepatoprotective effect of lupeol and lupeol linoleate on tissue antioxidant defence system in cadmium-induced hepatotoxicity in rats. Fitoterapia 2001;72:516-23.
31. Chaturvedi PK, Bhui K, Shukla Y. Lupeol: connotations for chemoprevention. Cancer Lett 2008;263:1-13.
32. Saleem M, Alam A, Arifin S, Shah MS, Ahmed B, Sultana S. Lupeol, a triterpene, inhibits early responses of tumor promotion induced by benzoil peroxide in murine skin. Pharmacol Res 2001;43:127-34.
33. Nigam N, Prasad S, George J, Shukla Y. Lupeol induces p53 and cyclin-B-mediated G2/M arrest and targets apoptosis through activation of caspase in mouse skin. Biochem Biophys Res Commun 2009;381:253-8.
34. Saleem M. Lupeol, A novel anti-inflammatory and anticancer dietary triterpene. Cancer Lett 2009;285:109-15.
35. Biringanine G, Chiarelli MT, Faes M, Duez P. A validation protocol for the HPTLC standardization of herbal products: application to the determination of acteoside in leaves of *Plantago palmate* Hook. Talanta 2006;69:418-24.
36. Kamboj A, Saluja AK. Isolation of stigmasterol and β -sitosterol from petroleum ether extract of aerial parts of *Ageratum conyzoides* (Asteraceae). Int J Pharm Pharm Sci 2011;3 Suppl 1:94-6.
37. Vijaykumar M Kal, Ajay G Namdeo. HPTLC densitometric evaluation by simultaneous estimation of galangin in *Alpinia galanga* and *Alpinia officinarum* Pharm Lett 2015;7 Suppl 7:158-64.
38. Sorenson WR, Sullivan D. Determination of campesterol, stigmasterol, and beta-sitosterol in saw palmetto raw materials and dietary supplements by gas chromatography: single-laboratory validation. J AOAC Int 2006;89:22-34.
39. Hamrapurkar PD, Karishma P. HPTLC determination of stigmasterol and tocopherol acetate in *Leptadenia reticulata* and in its formulation. J Planar Chromatogr Mod TLC 2007;20 Suppl 3:183-7.
40. Ling-Yi K, Min-Jian Q, Masatake N. Diterpenoids from the rhizomes of *Alpinia calcarata*. J Nat Prod 2000;63 Suppl 7:939-42.
41. Mizanur R, Atiar R, Hashem MA, Ullah M, Shakila A, Vidyanath C. Anti-inflammatory, analgesic and GC-MS analysis of essential oil of *Alpinia calcarata* rhizome. Int J Pharma Bio Sci 2012;3 Suppl 4:55-63.
42. Sonali Sethi, Om Prakash, Pant AK. Phytochemical analysis and isolation of stigmasterol and β -Sitosterol from the rhizomes of *Alpinia allughas* Rosc. Int J Curr Res Biosci Plant Biol 2015;2 Suppl 5:124-9.

How to cite this article

- Pratheema Philomindoss. Identification, quantification and validation of stigmasterol from alpinia calcarata using high performance thin layer chromatography method. Int J Pharm Pharm Sci 2017;9(9):126-131.