

Original Article

**CALLUS INDUCTION, PHYTOCHEMICAL STUDIES AND ANTIBACTERIAL ACTIVITY OF
DECALEPIS ARAYALPATHRA (JOSEPH AND CHANDRAS) VENTER**

H. R. RAVEESHA, K. S. ASHALATHA

Department of Botany, Bangalore University, Jnanabharathi Campus, Bengaluru 560056 Karnataka
Email: hrraveesh74@gmail.com

Received: 16 Jun 2017 Revised and Accepted: 02 Nov 2017

ABSTRACT

Objective: The aim of the present study was designed for the induction of callus from leaf explants of *Decalepis arayalpathra* (*D. arayalpathra*) and to analyse their phytochemical constituents and antibacterial activity.

Methods: The explants were cultured on Murashige and Skoog (MS) medium supplemented with different concentration of 2, 4-dichlorophenoxyacetic acid (2, 4-D) and later subcultured to the combination of 6-benzylaminopurine (BAP) and 1-naphthaleneacetic acid (NAA). The phytochemical constituents were analysed in the different solvent extracts using standard methods. Antibacterial activity of the different solvent extracts was carried out using agar well diffusion method against reference standards.

Results: Callus induction was observed on MS medium supplemented with different concentration and combination of auxins and cytokinins. Maximum callus induction was noticed on media supplemented with 2, 4-D (2 mg/l) and BAP (1 mg/l)+NAA (0.5 mg/l) respectively. The phytochemical screening revealed the presence of alkaloids, flavonoids, phenols, tannins, steroids and terpenoids, glycosides, coumarins and quinone etc. All the solvent extracts showed varying degree of antibacterial activities against the bacterial strains (*Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescence* and *Staphylococcus aureus*). The maximum antibacterial activity of leaf was observed in aqueous (*Klebsiella pneumoniae*, 19.00±1.00) and methanolic extract (*Klebsiella pneumoniae*, 18.33±1.15). Whereas petroleum ether extract of the callus showed maximum inhibition (*Bacillus subtilis*, 17.00±1.00) compared to other extracts.

Conclusion: The study revealed the presence of secondary metabolites in the leaf and callus extracts of *D. arayalpathra*. The methanolic extracts possess higher antibacterial activity compared to other solvent extracts. However, further studies have to be carried out for the isolation and identification of antimicrobial compounds against pathogens.

Keywords: Antibacterial, Auxins, Callus, *Decalepis arayalpathra*, Phytochemicals

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)
DOI: <http://dx.doi.org/10.22159/ijpps.2017v9i12.20743>

INTRODUCTION

Plants are an exemplary source of traditional medicine and pharmaceutical drugs for humankind since time immemorial. In the worldwide, it has been estimated that about 20,000 plants species are used as drugs. In India, herbal medicines have been used in the Indian traditional system of medicine (Ayurveda, Unani and Siddha) for the treatment of various diseases *SD arayalpathra* is one of the important medicinal plants, belonging to the family Periplocaceae, found in the southern region of the Western Ghats of India [1]. Tuberous roots of *D. arayalpathra* are consumed as pickles and as a popular cool drink known as Nannari [2-4]. The root extract was used by the kani tribes for the treatment of peptic ulcer and has a rejuvenating tonic [5]. Recent pharmacological studies have also revealed that root extract of the plant had immune-modulatory and antitumor activity [6] and as well as blood purifier [7-8]. Conventional propagation of *D. arayalpathra* is at stake with several factors like poor fruit set, seed germination and rooting on stem cutting [9-10]. Due to its high demand, the destructive harvesting of the roots leads to extinction of the species which necessitates another alternative method for propagation and conservation. Therefore, plant tissue culture techniques offer a powerful tool for mass multiplication of many plant species.

The development of drug resistance in human pathogen against commonly used antibiotics has opened a new avenue for the search of new antibacterial compounds from natural sources. Many of the plant species have been evaluated for antimicrobial properties, but the majority of them have not been systematically evaluated and a lot of attention is being derived to evaluate the plant extracts as an antibacterial agent against resistant plant pathogens. Recently, plant tissue culture techniques are used to enhance the secondary metabolite production in plants [11]. Studies on phytochemical and

antibacterial activities of the *in vitro* regenerated callus are limited. Therefore the present study was investigated to develop an effective protocol for callus induction and to analyse their phytochemical and antibacterial activities.

MATERIALS AND METHODS

Plant material

The plants were collected from the Western Ghats, Njaraneeli, Kerala and authenticated in the Department of Botany, Bangalore University. A voucher specimen is deposited in the herbarium (BUB, No. 2268). The plants were maintained in greenhouse at Department of Botany, Bangalore University, Bengaluru.

Callus induction

Leaf explants of *D. arayalpathra* were washed under running tap water (30 min) to remove soil particles then followed by different sterilants (bavastin and teepol). Explants were rinsed with distilled water for 2-3 times after each treatment. Finally, the explants were sterilized with 0.1% mercuric chloride for 2 min followed by washing with double distilled water for four times. Explants were then inoculated on MS media supplemented with different concentration of auxins (0.5, 1, 2, 3 and 4 mg/l of 2, 4-D), combination of auxins (0.5 mg/l of NAA) and cytokinins (0.5, 1, 2, 3 and 4 mg/l of BAP). Cultures were maintained at 25±2 °C under white fluorescent light for 16 hr photoperiod.

Preparation for plant extracts

10 gms of leaf and callus were grinded using mortar and pestle in different solvents (ethanol, methanol, petroleum ether, chloroform, hexane, acetone, butanol and distilled water). The extracts were filtered through Whatman No. 1 filter paper. The procedure was

repeated for another two cycles to ensure complete extraction of phytochemical compounds [12]. The filtrates were lyophilized and stored at 4 °C until further analysis.

Preliminary phytochemical analysis

The preliminary phytochemical analysis was carried out by standard methods [13-14]. Briefly, the lyophilized extracts were dissolved in respective solvents and screened for the qualitative analysis for the presence of alkaloids, flavonoids, proteins, phenols, tannins, steroids and terpenoids, phytosterols, glycosides, coumarins, carbohydrates, betacyanin, resins, phlobatannins, starch, volatile oils, emodols [15-16].

Antibacterial assay

The solvent extracts were assayed against the following organisms *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas fluorescense* and *Staphylococcus aureus*. Bacterial strains were obtained from Department of Microbiology, Bangalore University, Bengaluru. *In vitro* antibacterial activity was performed by agar well diffusion method according to the protocol of Johnson *et al.* [17]. Wells were bored into nutrient agar using a sterile 8 mm diameter cork borer. Different solvent extracts (150 µl) were added into the wells using sterilized pipettes and allowed to diffuse at room temperature for 2 h. The plates were incubated at 37 °C for 24 h.

After the incubation, the diameter of the zone of inhibition was recorded in millimetre and compare with standard antibiotics (tetracycline and streptomycin). The experiments were repeated thrice in triplicates.

Statistical analysis

The results were expressed as mean± standard deviation (SD). Data were analyzed statistically by one-way analysis of variance followed by Duncan's multiple range tests using SPSS software. Probability values $p < 0.05$ were considered significant.

RESULTS

In vitro callus induction for leaf explants

The induction of callus was observed from leaf explants within two weeks of inoculation on MS media supplemented with various plant growth regulators. Maximum callus induction was observed after eight weeks of cultures on MS media supplemented with 2 mg/l of 2, 4-D and combinations of BAP (1 and 2 mg/l)+NAA (0.5 mg/l) (table. 1, fig. 1 A-F). Further increase in growth hormone concentration, decrease in the percent of callus induction. The morphology and growth of the callus were affected by varying concentration of 2, 4-D (creamish callus) and the combination of BAP+NAA (green friable callus). There was no callus induction on basal medium.

Table 1: Effect of plant growth hormones on callus induction from leaf explants of *D. arayalpathra*

Growth regulators	Concentration	% response±SD	Nature of callus
2, 4-D	0.5 mg/l	80.00±5.00 ^{cd}	Creamish callus
	1.0 mg/l	90.00±5.00 ^{ef}	Creamish green callus
	2.0 mg/l	96.67±2.89 ^f	Creamish yellow callus
	3.0 mg/l	85.00±5.00 ^{de}	Creamish yellow callus
	4.0 mg/l	55.00±5.00 ^a	Creamish yellow callus
BAP+NAA	0.5 mg/l+0.5 mg/l	90.00±5.00 ^{ef}	Friable green callus
	1.0 mg/l+0.5 mg/l	96.67±2.89 ^f	Greenish callus
	2.0 mg/l+0.5 mg/l	95.00±5.00 ^f	Friable green callus
	3.0 mg/l+0.5 mg/l	75.00±5.00 ^{bc}	Creamish green callus
	4.0 mg/l+0.5 mg/l	70.00±5.00 ^b	Creamish green callus

Values represent the mean±SD. Means with the different letters in columns indicate significant differences at 5% level.




Fig. 1: A-F. *In vitro* callusing from leaf explants of *D. arayalpathra*. A-C: MS+2, 4-D (0.5 mg/l), (1 mg/l), (2 mg/l) respectively. D: MS+BAP (0.5 mg/l)+NAA (0.5 mg/l), E: MS+BAP (1 mg/l)+NAA (0.5 mg/l), F: MS+BAP (2 mg/l)+NAA (0.5 mg/l)

Phytochemical analysis

The secondary metabolites produced by the plants play an important role in plant defence against prey, microorganisms, stress as well as interspecies protections. These secondary metabolites have been used as a drug from the time immemorial, hence screening of phytochemicals serve as the initial steps in predicting

the potential active compounds in the plant extracts [18]. Phytochemical studies revealed the presence of alkaloids, flavonoids, proteins, phenols, tannins, steroids and terpenoids, phytosterols, glycosides, coumarins, carbohydrates, betacyanin, resins, phlobatannins, volatile oils. Ethanol, methanol and acetone were proven to be better solvents for the extraction of major phytochemicals compare to other solvents (table 2).

Table 2: Preliminary phytochemical analysis on fresh leaf and callus of *D. arayalpathra*

Tests	Fresh leaf								Fresh callus							
	I	II	III	IV	V	VI	VII	VIII	I	II	III	IV	V	VI	VII	VIII
Alkaloids	+	+	+	+	-	+	-	-	-	+	-	-	-	+	-	+
Flavonoids	+	+	-	-	-	+	+	+	+	+	-	-	-	+	-	+
Proteins	+	+	+	+	+	+	+	+	+	+	-	-	+	+	+	+
Phenols	+	+	-	-	-	+	+	+	+	+	-	-	-	-	-	+
Tannins	+	+	+	+	-	+	+	+	-	+	-	-	-	-	-	+
Steroids and Terpenoids	+	+	+	+	-	-	+	-	-	-	-	-	-	-	-	+
Phytosterols	+	+	+	+	-	-	+	-	-	-	-	-	-	-	-	+
Glycosides	+	+	-	-	-	+	+	+	+	+	-	-	-	-	-	-
Coumarins	+	+	+	+	-	+	-	+	-	+	+	+	-	+	-	-
Carbohydrates	+	+	-	+	-	-	+	+	+	+	-	+	+	-	-	+
Betacyanin	+	+	+	+	+	+	-	+	+	+	-	-	-	+	-	+
Resins	+	+	-	-	-	-	-	-	-	+	+	+	+	-	-	+
Phlobatannins	+	+	-	-	-	-	-	+	-	-	-	-	-	+	-	-
Starch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Volatile oils	+	+	-	-	-	+	-	+	-	-	-	-	-	-	-	+
Emodols	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

I-Ethanol, II-Methanol, III-Petroleum ether, IV-Chloroform, V-Hexane, VI-Acetone, VII-Butanol, VIII-Aqueous

Antibacterial activity

The present study revealed the antibacterial activities of the leaf and callus extracts (150 µg/ml) against bacterial strains (*Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Klebsiella pneumoniae* and

Pseudomonas fluorescence) by agar well diffusion method. Aqueous and methanol leaf extracts showed maximum inhibition against *Klebsiella pneumoniae* compared to other extracts (fig. 2). Whereas in callus, ethanol and petroleum ether extracts showed higher inhibitory activity against *Escherichia coli* and *Bacillus subtilis* (fig. 3).




Fig. 2: Zone of inhibition on different solvents of *D. arayalpathra* fresh leaf, data are presented as the mean±SE of the mean of triplicates

DISCUSSION

Plants are important sources of potentially useful substances for the development of new therapeutic agents [19]. The aromatic roots of *D. arayalpathra* have been overexploited for therapeutic use leading to drastic reduction in the population of these species in the wild habitat. The plant biotechnology provides new tools for collection, multiplication and conservation of plant biodiversity by using *in vitro* culture techniques [20]. There are limited reports on

micropropagation of *D. arayalpathra* [10, 21]. Therefore, the present study was investigated to find out the effective protocol for the micropropagation of *D. arayalpathra* using leaf explants.

The efficiency of callus induction depends on the type of growth regulators, explants source and culture medium. In our study, the green friable callus induction was observed on MS media supplemented with a lower concentration of BAP and NAA. Formation of green friable callus may be due to their role in DNA

synthesis and mitosis [22]. Our results are in agreement with the studies of Umesh [23] and Sudha et al. [21] in *Decalepis hamiltonii* using nodal explants. Similar results were reported in other plant

species (*Amaranthus* spp., and *Ruta graveolens*) wherein, callus induction was observed from leaf explants inoculated on MS medium supplemented with BA and NAA [24-25].




Fig. 3: Zone of inhibition on different solvents of *D. aryalpathra* leaf callus, data are presented as the mean \pm SE of the mean of triplicates

Phytochemical constituents present in plants such as alkaloids, flavonoids, and tannins etc., serves as a defense mechanism against predation by many microorganisms. Chandra and Kaneria [26] have reported that the accumulation of secondary metabolites may vary in different plant parts and leaf is one such plant parts which have highest accumulation of metabolites [27]. Our qualitative analysis revealed the presence of major phytochemicals in the leaf compared to callus. Previous studies reported lesser accumulation of phytochemicals in the callus compared to other plant parts [28, 29]. Aqueous, methanol and acetone extracts are the best solvents to isolate major phytochemicals compared to other solvents. Johnson et al. [17] reported that major phytochemicals were extracted in ethanol and methanol leaf and callus extracts of *Baliospermum montanum*.

Plant extracts can be a good source of antibiotics against various bacterial pathogens. In the present study, antibacterial activities of leaf and callus extracts of *D. aryalpathra* were calculated by measuring the diameter of the zone around the well. The maximum inhibition was observed in leaf extracts against major pathogens compared to callus due to their lesser accumulation of secondary metabolites. Various workers have reported that root, leaf and callus extracts can inhibit both gram positive and gram negative bacteria [30-32]. The mode of action of antimicrobial agents may be due to interference with cell wall synthesis, inhibition of protein synthesis, interference with nucleic acid synthesis, and inhibition of a metabolic pathway [33].

CONCLUSION

In the present study, an efficient protocol was developed for callus induction from leaf explants of *D. aryalpathra*. The phytochemical analysis of the leaf and callus revealed the presence of bioactive compounds with antibacterial activity against many pathogens. Further studies have to be carried to characterize these antimicrobial compounds and large-scale production through *in vitro* propagation.

CONFLICT OF INTERESTS

No conflict of interest

REFERENCES

1. Sharma S, Shahzad A. An overview on *decalepis*; a genus of woody medicinal climbers. Open Sci Publications 2014;1:1-13.

- Vedavathy S. *Decalepis hamiltonii* Wight and Arn—an endangered source of indigenous health drink. Nat Prod Radiance 2004;3:22-3.
- Vijayakumar V, Pullaiah T. An ethno-medico-botanical study of Prakasam district, Andhra Pradesh, India. Fitoterapia 1998; 69:483-9.
- Harish R, Srivastava A, Divakar S, Shivanandappa T. Isolation of antioxidant compounds from the methanolic extract of the roots of *Decalepis hamiltonii* (Wight and Arn.). J Agric Food Chem 2005;53:7709-14.
- Pushpangadan P, Rajasekharan A, Ratheeshkumar PK, Jawahar CR, Radhakrishnan K. Amrithapala (*Janakia aryalpathra* Joseph and Chandrasekharan), a new drug from the Kani tribe of Kerala. Anc Sci Life 1990;9:212-4.
- Subramaniam A, Rajasekharan S, Latha PG, Evans DA, Pushpangadan P. Immuno-modulatory and antitumor activities of *Janakia aryalpathra*. Fitoterapia 1996;57:140-4.
- Chopra RN, Nayar SI, Chopra LC, Asolkar LV, Kakkar KK. Glossary of Indian Medicinal Plants; Council of Scientific and Industrial Research, New Delhi; 1956.
- Jacob KC. An unrecorded economic product *Decalepis hamiltonii* W. and Arn, family asclepidaceae. Madras Agric J 1937;25:176.
- Sudha CG, Seeni S. Establishment and analysis of fast-growing normal root culture of *Decalepis aryalpathra*, a rare endemic medicinal plant. Curr Sci 2001;81:371-4.
- Gangaprasad A, Decruse SW, Seeni S, Nair GM. Micropropagation and ecorestoration of *Decalepis aryalpathra* (Joseph and Chandra.) venter—an endemic and endangered ethnomedicinal plant of Western Ghats. Indian J Biotechnol 2005;4:265-70.
- Lee Y, Lee DE, Lee HS, Kim SK, Lee WS, Kim SH, et al. Influence of auxins, cytokinins and nitrogen on the production of rutin from callus and adventitious roots of the white mulberry tree (*Morus alba* L.). Plant Cell Tiss Org Cult 2011;105:9-19.
- Dhanapal ACTN, Ming TW, Aung HP, Hao SJ. Preliminary screening of *Artemisia argyi* for antioxidant potentials. Int J Pharmacogn Phytochem Res 2016;8:347-55.
- Harbone JB. Phytochemical Methods. Chapman and Hall Ltd, London; 1973. p. 33-80.

14. Borah U, Dash B, Dash S, Deka J, Kalita J. Preliminary phytochemical screening and *in vitro* antimicrobial activity of ethanolic extract of whole aerial part of the herb *Leucas plukenetii* spreng (Family-Lamiaceae). Int J Curr Pharm Res 2017;9:87-90.
15. Mohan B, Nayak JB, Sunil Kumar R, Shiva Kumari LP, Mohan CH, Rajani B. Phytochemical screening, GC-MS analysis of *Decalepis hamiltonii* Wight and Arn. An endangered medicinal plant. J Pharmacogn Phytochem 2016;5:10-6.
16. Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction; a review. Int Pharm Sci 2011;1:98-106.
17. Johnson M, Wesely EG, Zahir Hussain MI, Selvan N. *In vivo* and *in vitro* phytochemical and antibacterial efficacy of *Baliospermum montanum* (Willd.) Muell. Arg Asian Pacific J Trop Med 2010;3:894-7.
18. Chew YL, Chan EWL, Tan PL, Lim YY, Stanslas J, Goh JK. Assessment of phytochemical content, polyphenolic composition, antioxidant and antibacterial activities of Leguminosae medicinal plants in Peninsular Malaysia. BMC Complem Altern Med 2011;11:12.
19. Rajeswari A. Evaluation of phytochemical constituents, quantitative analysis and antimicrobial efficacy of potential herbs against selected microbes. Asian J Pharm Clin Res 2015;8:232-7.
20. Samydurai P, Saradha M, Ramakrishnan R, Santhosh Kumar S, Thangapandian V. Micropropagation prospective of cotyledonary explants of *Decalepis hamiltonii* Wight and Arn.- an endangered edible species. Indian J Biotechnol 2016; 15: 256-60.
21. Sudha CG, Krishnan PN, Pushpangadan P, Seeni S. *In vitro* propagation of *Decalepis arayalpathra*, a critically endangered ethnomedicinal plant *in vitro* cell. Dev Biol-Plant 2005;41:648-54.
22. Skoog F, Miller CO. Chemical regulation of growth and organ formation in plant tissue cultures *in vitro*. Symp Soc Exp Biol 1957;11:118-31.
23. Umesh TG. *In vitro* callus induction and antioxidant potential of *Decalepis hamiltonii* (Wight and Arn). Int J Pharm Pharm Sci 2014;6:452-6.
24. Pannu J, Thalwal S, Gupta A. Comparison of antimicrobial activity and phytochemical constituents of *in vivo* and *in vitro* grown *Amaranthus spinosus* plants. Int J Pharm Pharm Sci 2013;5:703-7.
25. Ahmad N, Faisal M, Anis M, Aref LM. *In vitro* callus induction and plant regeneration from leaf explants of *Ruta graveolens* L. South Afr J Bot 2010;76:597-600.
26. Chandra S, Kaneria M. Indian nutraceutical plant leaves as a potential source of natural antimicrobial agents. In: A Mendez Vilas. Ed. Science against microbial Pathogens: Communicating Current Research and Technological Advances, Fomatex Research Center Publication, Spain; 2011. p. 1251-9.
27. Pires De Abreu LR, Ortiz RM, Castro de SC. HPLC determination of amoxicillin comparative bioavailability in healthy volunteers after a single dose administration. J Pharm Pharm Sci 2003;6:223-30.
28. Arumugam T, Ayyanar M, Koli Pillai YJ, Sekar T. Phytochemical screening and antibacterial activity of leaf and callus extracts of *Centella asiatica*. Bangladesh J Pharmacol 2011;6:55-60.
29. Thenmozhi M, Sivaraj R. *In vitro* evaluation of the antibacterial activity of *Petunia* leaf and callus extracts. J Agric Tech 2011;7:321-30.
30. Murugan VK, Arumugam T, Tamilvannan V, Umarani K, Dhmocharan R, Ravikumar S. Phytochemical analysis and *in vitro* antibacterial properties of selected medicinal plants. Eur J Pharm Med Res 2015;2:189-94.
31. Prakash P, Thiyagarajan G, Manivasagaperumal R. Phytochemical screening and antibacterial activity of root extracts of *Decalepis hamiltonii* Wight and Arn. Int J Pharm Res Rev 2014;3:33-8.
32. Rajani B, Mohan B, Uma Devi M, Shiva Kumarich LP. Phytochemical studies and antibacterial activity of *Decalepis hamiltonii* wight and arn, an endangered medicinal plant. J Med Plant Studies 2016;4:88-91.
33. Laxmi A, Siddhartha S, Archana M. Antimicrobial screening of methanol and aqueous extracts of *Swertia chirata*. Int J Pharm Pharm Sci 2011;3:142-6.