

## IN VIVO ANTIOXIDANT ACTIVITY OF *LIMNOPHILA HETEROPHYLLA* AND *MICHELIA CHAMPACA*

RAJA S., RAVINDRANADH K.

GITAM Institute of Pharmacy, GITAM University, Visakhapatnam Andhra Pradesh, India 530045

Email: srja61@gmail.com

Received: 14 Aug 2017 Revised and Accepted: 02 Nov 2017

### ABSTRACT

**Objective:** The present study was aimed at investigating the *in-vivo* antioxidant activity of the methanol extracts of *Limnophila heterophylla* and *Michelia champaca* leaves.

**Methods:** Methanol extract of both plants were administered to rats separately at three different doses of 125, 250 and 500 mg/kg for 21 d to evaluate oxidative stress parameters such as ferric reducing ability of plasma (FRAP), thiobarbituric acid reactive substance (TBARS) and reduced glutathione (GSH) and to evaluate antioxidant enzyme levels of catalase (CAT) and superoxide dismutase (SOD).

**Results:** The methanol extracts of both the plants significantly ( $p < 0.05$ ) elevated the ferric reducing ability of plasma (FRAP) on days 7, 14 and 21 of treatment. Significant ( $p < 0.05$ ) decrease of thiobarbituric acid reactive substance (TBARS) levels along with an increase in the superoxide dismutase (SOD) enzyme level in the liver and kidney at three different doses both the plants was observed. Treatment at a dose of 500 mg/kg b. w of both plants caused a significant increase only in the level of CAT in the liver and kidney. However, there was no significant effect of a thiobarbituric acid reactive substance (TBARS), superoxide dismutase (SOD) and catalase (CAT) in the heart and reduced glutathione (GSH) level in liver, heart and kidney at three different doses both the plants.

**Conclusion:** These outcomes recommend that the leaves of *Limnophila heterophylla* and *Michelia champaca* have a potent antioxidant activity which may be responsible for some of its reported pharmacological actions.

**Keywords:** *Limnophila heterophylla*, *Michelia champaca*, FRAP, Catalase, TBARS, GSH

© 2017 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open-access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ijpps.2017v9i12.22013>

### INTRODUCTION

In the development of diseases, free radicals have emerged because the major entity inflicting harm to cells. These cytotoxic metabolites are generated by aerobic metabolism within the cell that successively considerably will increase pathological conditions, resulting in free radical mediate denaturation of protein, enzymatic deactivation, base hydroxylation of nucleic acids, cross-linking or strand cutting, mutation or maybe death [1]. However, the physiological system has a series of defense mechanism including antioxidant enzymes-superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX), Thiobarbituric acid reactive substances (TBARS), reduced glutathione (GSH), and other free radical scavengers,  $\beta$ -carotene, vitamin C, vitamin E,  $\alpha$ -lipoic acid, curcumin, rutin, BHT and glutathione to protect the cell against cytotoxic ROS (reactive oxygen species) [2]. The antioxidants in a biological system can be either enzymatic or non-enzymatic. The enzymatic antioxidants include catalase (CAT), superoxide dismutase (SOD), and glutathione which catalyse neutralization of many types of free radicals [3], while the nonenzymatic antioxidants include vitamin C, selenium, vitamin E, carotenoids, and polyphenols. The enzymatic antioxidants catalyse neutralization of many types of free radicals [3]. However, when the balance between the oxygen species and antioxidants is altered, a state of oxidative stress results, possibly leading to everlasting cellular damage. There is evidence that antioxidants may be useful in preventing the deleterious consequences of oxidative stress, and there is an increasing interest in the protective biochemical function of natural antioxidants contained in vegetables, fruits and medicinal herbs [4]. Generally, plants, herbs, and spice, rich in phenolic compounds like flavonoids, have been demonstrated to have anti-inflammatory, antiallergenic, antiviral, antiaging and anticarcinogenic activities which can be attributed to their antioxidant properties [5]. In this respect, flavonoids and polyphenolic compounds have received the greatest attention [6-8].

*Limnophila heterophylla* is an aquatic herb, mainly submerged, but with shoots that often emerge above the water surface, rooting at nodes. The plant finds lot of applications in the traditional system of medicine to treat wounds [9]. Different parts of *Limnophila heterophylla* possess varied pharmacological activities like COX inhibitor [10], antimicrobial [11] and wound healing [12]. The plant encloses terpene, flavanoids, terpenoids and oils [13]. *Michelia*, known by the scientific name *Michelia champaca*, is a very tall tree that grows up to 30m tall. *Michelia champaca* is used ethnomedicinally for the handling of astringent, constipation, dyspepsia, dysmenorrhea, fever, febrifuge, nausea, stomachic, skin disease, tonic, ulcers and wounds [14]. Earlier pharmacological reports of *Michelia champaca* had demonstrated its cytotoxic activity [15], anti-inflammatory [16], antihyperglycemic [17], leishmanicidal [18], antibacterial [19], wound healing [20], diuretic [21], antiulcer [22], antifertility [23], antihelminthic [24] and cardioprotective [25] activities. Several phytoconstituents like alkaloids, flavonoids, triterpenoids, saponins, tannins, sterols and steroids have been isolated from different parts *Michelia champaca*.

Based on these reports, this study was designed to investigate the *in vivo* antioxidant activity of methanol extracts of leaves of *Limnophila heterophylla* and *Michelia champaca*.

### MATERIALS AND METHODS

#### Plant materials

The plants were collected from Tirupati (Andhra Pradesh), India and further, plants were distinguished, affirmed and validated by Dr. Madavchetty, Professor, Botany office, Sri Venkateswara University, Tirupati. A voucher specimen of these plants (*Limnophila heterophylla*-GIP006/2013-2014 and *Michelia champaca*-GIP005/2013-2014) have been kept in the GITAM Institute of Pharmacy, GITAM University, Visakhapatnam, Andhra Pradesh, India.

## Chemicals and reagents

All chemicals used in the study were of analytical grade and they were procured from Coastal Enterprises Pvt. Ltd., Visakhapatnam, Andhra Pradesh and India.

## Extraction

Leaves of *Limnophila heterophylla* and *Michelia champaca* were dried under shade and then powdered with a mechanical grinder to obtain a coarse powder. The powder was passed through 40 mesh sieve and extracted with methanol separately in Soxhlet apparatus at 60 °C. The solvent was completely removed by rotary vacuum evaporator and concentrated. The extracts were freeze-dried and stored in a vacuum desiccator for further *in vivo* antioxidant studies.

## Test animals and groups

Wistar albino rats (200–250 gm) of either sex were maintained under standard environmental conditions and had free access to feed and water ad libitum. Experiments on animals were performed based on animal ethics guidelines of Institutional Animal Ethics Committee (IAEC). Rats were divided into seven groups of six animals each. Group, I served as control and was given the vehicle alone (normal saline). Group II, III and IV received methanol extract of *Limnophila heterophylla* orally at 125, 250 and 500 mg/kg body weight, individually. Group V, VI and VII received methanol extract of *Michelia champaca* orally at 125, 250 and 500 mg/kg body weight, separately. The treatments were given for 21 d and all the animals were sacrificed by decapitation on 22<sup>nd</sup> day of the experiment. Blood was collected through the direct cardiac puncture and it was used for *in vivo* antioxidant activity. The heart, liver and kidney were removed, washed in cold saline and stored in liquid nitrogen for further biochemical studies. This *in vivo* antioxidant activity was analyzed by the method described by Rajlakshmi *et al.* [26].

## Serum preparation

Blood was allowed to clot for 30 min, then centrifuged at 2500 rpm for 15 min and serum was harvested. The serum was prepared using standard method as described by Yesufu *et al.*, [27] and it was used for the estimation of superoxide dismutase (SOD), catalase (CAT) and malonyldialdehyde (MDA).

## Preparation of rat heart, liver and kidney homogenate

Tissue homogenate was prepared in a ratio of 1 gm of wet tissue to 10 times (w/v) 0.05 mol/l ice-cold phosphate buffer (pH 7.4) and homogenised using a homogenizer (Tissue homogeniser). A 0.2 ml sample of homogenate was used for assessment of thiobarbituric acid reactive substance (TBARS). The leftover part of the homogenate was separated into two parts. One part was mixed with 10% trichloroacetic acid (1:1), centrifuged at 5000g (4 °C, for 10 min) and the supernatant was used for reduced glutathione (GSH) estimation. The second part of the homogenate was centrifuged at 15 000g at 4 °C for 60 min and the supernatant was used for superoxide dismutase (SOD) and catalase (CAT) estimation.

## Ferric reducing ability of plasma (FRAP) assay

Total plasma antioxidant capacity was measured according to the ferric reducing ability of plasma (FRAP) method [28]. The blood samples were collected from the rat retro-orbital venous plexus into heparinised glass tubes at 0, 7 14 and 21 d of treatment. Briefly, 3 ml of freshly prepared and warm (37 °C) FRAP reagent (1 ml of 10 mmol/l TPTZ [2,4,6 tripyridyl-s-triazine] solution in 40 mmol/l HCl, 1 ml 20 mmol/l FeCl<sub>2</sub>·6H<sub>2</sub>O, 10 ml of 0.3 mmol/l acetate buffer [pH 3.6]) was mixed with 0.375 ml distilled water and 0.025 ml of test samples. The absorbance of developed colour in the organic layer was measured at 593 nm. The temperature was maintained at 37 °C. The readings at 180 sec were selected for the calculation of FRAP values. Ferrous sulphate (FeSO<sub>4</sub>·7H<sub>2</sub>O) was used as a standard for calibration and the data expressed as nmol Fe<sup>2+</sup>/l.

## Superoxide dismutase assay

Superoxide dismutase (SOD) activity was analysed by the method described by Rai *et al.*, [29]. Assay mixture contain 0.1 ml of supernatant/serum, 1.2 ml of sodium pyrophosphate buffer (pH 8.3;

0.052M), 0.1 ml of phenazine methosulfate (186 mmol), 0.3 ml of nitroblue tetrazolium (300 mmol), and 0.2 ml of NADH (750 mmol). The reaction was started by the addition of NADH. After incubation at 30 °C for 90s, the reaction was stopped by the addition of 0.1 ml of glacial acetic acid. The reaction mixture was stirred vigorously with four ml of n-butanol. Color intensity of the chromogen in the butanol was measured spectrophotometrically at 560 nm and the concentration of superoxide dismutase (SOD) was expressed as units/mg of protein.

## Catalase assay

Catalase activity (CAT) was measured by the method of Aebi [30]. A 0.1 ml of supernatant/serum was added to a cuvette containing 1.9 ml of 50 mmol phosphate buffer (pH 7.0). The reaction was started by the addition of 1.0 ml of freshly prepared 30 mmol H<sub>2</sub>O<sub>2</sub>. The rate of the decomposition of H<sub>2</sub>O<sub>2</sub> was measured spectrophotometrically at 240 nm. The activity of catalase (CAT) was expressed as units/mg of protein.

## Estimation of thiobarbituric acid reactive substance (TBARS)

Lipid peroxidation (LPO) was measured by the method of Liu *et al.*, [31]. Acetic acid 1.5 ml (20%; pH 3.5), 1.5 of TBA (0.8%), and 0.2 ml of sodium dodecyl sulfate (8.1%) was added to 0.1 ml of supernatant/serum and heated at 100 °C for 60 min. The mixture was cooled to room temperature. To this, 5 ml of n-butanol: pyridine mixture and 1 ml of distilled water were added and vortexed vigorously. After centrifugation at 1200g for 10 min, the organic layer was separated and the absorbance was measured at 532 nm using a spectrophotometer. Malonyldialdehyde (MDA) was an end product of LPO, which reacts with TBA to form pink chromogen-TBA reactive substance. It was calculated using a molar extinction coefficient of 1.56 X 10<sup>5</sup>M<sup>-1</sup> cm<sup>-1</sup> and it was expressed as nM/g wet wt.

## Estimation of reduced glutathione (GSH)

Glutathione was measured according to the method of Ellman [32]. An equal quantity of homogenate was mixed with 10% trichloroacetic acid and it was centrifuged to separate the proteins. To 0.01 ml of this supernatant, 2 ml of phosphate buffer (pH 8.4), 0.5 ml of 5'-dithio, bis (2-nitrobenzoic acid) and 0.4 ml double distilled water was added. The mixture was vortexed and the absorbance was read at 412 nm within 15 min. The concentration of reduced glutathione was expressed as µg/g tissue.

## Statistical analysis

All experimental data were expressed as mean±standard error of the mean (SEM). This Statistical analysis was carried out using one-way analysis of variance (ANOVA) followed by Dunnett-t-test with the SPSS statistical software for comparison to the control group. p<0.05 was considered as statistically significant.

## RESULTS

### Serum enzymatic levels of MDA, SOD and CAT

The results of the serum enzymatic levels of malonyldialdehyde (MDA), superoxide dismutase (SOD) and catalase (CAT) of individual methanol extract of *Limnophila heterophylla* and *Michelia champaca* were mentioned in table 1. At a dose of 250 mg/kg both the extracts exhibited moderate significant (p<0.05, p<0.01 and p<0.001) decrease in the serum level of malonyldialdehyde (MDA) in the test animals (group II to VII) when compared to control. However, only the higher dose (500 mg/kg) of both the plant methanol extracts produced a significant (p<0.001) decrease in the serum levels of malonyldialdehyde (MDA) when compared with control. The extracts of treated rats also evoked a dose-related significant (p<0.05, p<0.01 and p<0.001) increase in the serum levels of catalase (CAT) and superoxide dismutase (SOD) when compared to control. Methanol extracts of both the plants at a dose of 250 mg/kg showed moderate significant (p<0.01) elevation in serum levels of catalase (CAT) and superoxide dismutase (SOD). However, only the higher dose (500 mg/kg) of both the plant methanol extracts caused a significant (p<0.001) increase in the serum levels of catalase (CAT) and superoxide dismutase (SOD) when compared with control. However, there was mild significant (p<0.05) changes of malonyldialdehyde (MDA), superoxide dismutase (SOD) and catalase (CAT) levels were observed in 125 mg/kg of both the plant methanol extracts.

Table 1: Serum enzymatic levels of MDA, SOD and CAT

Group	MDA (nM of TBARS mg <sup>-1</sup> of protein)	SOD U/mg protein	CAT U/mg protein
Group-I	0.46±0.32	46.26±1.28	26.22±0.01
Group-II	0.39±0.01*	48.14±2.62*	26.82±0.18*
Group-III	0.33±0.12**	53.6±1.37**	28.44±1.12**
Group-IV	0.29±1.32***	54.8±1.09***	40.06±1.87***
Group-V	0.44±0.02*	45.23±1.23*	22.12±0.08*
Group-VI	0.36±0.32**	49.7±1.08**	26.32±1.23**
Group-VII	0.32±0.42***	51.7±1.22***	37.25±1.82***

[p<0.05\*, p<0.01\*\* and p<0.001\*\*\*]

#### Ferric reducing ability of plasma (FRAP) assay

The ferric reducing ability of plasma (FRAP) level of rats after administration of methanol extracts of *Limnophila heterophylla* and *Michelia champaca* over a period of 21 d was presented in fig. 1. In the control group, there was no significant change in ferric reducing ability of plasma (FRAP) value on days 7 (870 nM Fe<sup>2+</sup>/l) 14 (869 nM Fe<sup>2+</sup>/l) and 21 (868 nM Fe<sup>2+</sup>/l) compared with day 0 (864 nM Fe<sup>2+</sup>/l). But, in group II, III and IV on days 7 (947, 1088 and 1101 nM Fe<sup>2+</sup>/l, respectively), 14 (1104, 1202 and 1287 nM Fe<sup>2+</sup>/l, respectively) and 21 (1122, 1309 and 1344 nM Fe<sup>2+</sup>/l, respectively) there was a significant (p<0.05, p<0.01 and p<0.001)

increase in ferric reducing ability of plasma (FRAP) value compared with day 0 (869, 870 and 871 nM Fe<sup>2+</sup>/l, respectively). Similarly, in group V, VI and VII on days 7 (941, 1075 and 1095 nM Fe<sup>2+</sup>/l, respectively), 14 (1097, 1189 and 1265 nM Fe<sup>2+</sup>/l, respectively) and 21 (1103, 1297 and 1321 nM Fe<sup>2+</sup>/l, respectively) there was a significant (p<0.05, p<0.01 and p<0.001) increase in ferric reducing ability of plasma (FRAP) value compared with day 0 (868, 870 and 871 nM Fe<sup>2+</sup>/l, respectively). Maximum enrichment of ferric reducing ability of plasma (FRAP) level was found in group IV and VII, which correspond to animals administered with 500 mg/kg body weight of the methanol extract of *Limnophila heterophylla* and *Michelia champaca*, individually.




Fig. 1: Changes in rat total antioxidant capacity of the ferric reducing ability of plasma (FRAP) measured by Fe<sup>2+</sup> equivalent after administration of methanol extract of *Limnophila heterophylla* and *Michelia champaca*. Values were mean±s. d (n=6). Group II, III, IV, V, VI and VII (Plant Extracts Treated Rats) Compared to Group I (control rats). p<0.05\* and p<0.01\*\*

#### Estimation of superoxide dismutase (SOD)

The treatment of methanol extract of both plants caused minor significant (p<0.05, p<0.01 and p<0.001) decrease at 125 mg/kg (510 and 498 units/mg of protein), 250 mg/kg (528 and 524 units/mg of protein) and 500 mg/kg (536 and 532 units/mg of protein) bodyweight in the level of superoxide dismutase (SOD) in the heart compared with the control (522 units/mg of protein). However, the level of superoxide dismutase (SOD) in the kidney and liver of the both plants treated rats was not dose related and was found to be significantly increased (p<0.05, p<0.01 and p<0.001) at the three different doses of 125 mg/kg (206, 198 and 428, 419 units/mg of protein, respectively), 250 mg/kg (218, 209 and 462, 455 units/mg of protein, respectively) and 500 mg/kg (220, 211 and 478, 469 units/mg of protein, respectively) compared with the control (group-I) (204 and 424 units/mg of protein, respectively). The results were shown in fig. 2, 3 and 4.

#### Estimation of catalase (CAT)

The administration of methanol extract of both plants to normal rats for 21 d induced a dose-dependent increase in the level of catalase (CAT) in liver (fig. 2) and kidney (fig. 4), but a decrease in the heart (fig. 3). Methanol extract of both plants were significantly increased at 500 mg/kg body weight dose of the treatment (p<0.001) for liver (344 and 339 units/mg of protein) and kidney (402 and 394 units/mg of protein) compared with the control group of liver and kidney (312 and 385 units/mg of protein, respectively). However, there was a little change in the endogenous antioxidant levels in heart tissue. Similarly, for both the plants at a dose of 125 and 250 mg/kg a significant increase (p<0.05 and p<0.01) in the catalase (CAT) levels were observed for liver (308 and 298 units/mg of protein) and kidney (387 and 374 units/mg of protein) compared with the control group.





Fig. 2: Changes in rat liver thiobarbituric acid reactive substance (TBARS; nmol/g wet weight [wt]), superoxide dismutase (SOD; units/mg protein), catalase (CAT; units/mg protein) and reduced glutathione (GSH; µg/g wet wt) following oral administration of methanol extracts of *Limnophila heterophylla* and *Michelia champaca*. Values were mean±SD (n = 6). p<0.05\* and p<0.01\*\*, compared with control

### Estimation of thiobarbituric acid reactive substance (TBARS)

The effect of different doses of individual methanol extract of *Limnophila heterophylla* and *Michelia champaca* on the lipid peroxidation and endogenous antioxidants of liver, heart and kidney of rats was shown in fig. 2, 3 and 4 respectively. For groups II, III, and IV, a significant ( $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ ) decrease in thiobarbituric acid reactive substance (TBARS) concentration in liver (339, 287 and 284 nmol/g wet weight [wt] tissue, respectively) and kidney (182, 176 and 168 nmol/g wet wt tissue, respectively) was observed and

compared with the control group (341 and 184 nmol/g wet wt tissue in liver and kidney, respectively). Similarly, for groups V, VI and VII a significant ( $p < 0.05$ ,  $p < 0.01$  and  $p < 0.001$ ) decrease in thiobarbituric acid reactive substance (TBARS) concentration in liver (329, 292 and 288 nmol/g wet weight [wt] tissue, respectively) and kidney (176, 179 and 168 nmol/g wet wt tissue, respectively) was observed and compared with the control group (341 and 184 nmol/g wet wt tissue in liver and kidney, respectively). Also, the experiment study indicated that a negligible change in the level of thiobarbituric acid reactive substance (TBARS) in the heart was observed for all treated groups.




**Fig. 3: Changes in rat heart thiobarbituric acid reactive substance (TBARS; nmol/g wet weight [wt]), superoxide dismutase (SOD; units/mg protein), Catalase (CAT; units/mg protein) and Reduced Glutathione (GSH;  $\mu\text{g/g}$  wet wt) following oral administration of methanol extracts of *Limnophila heterophylla* and *Michelia champaca*, values were mean  $\pm$  SD (n = 6).  $p < 0.05^*$ , compared with control**

### Estimation of reduced glutathione (GSH)

For groups II, III and IV, there was no significant change of reduced glutathione (GSH) level in liver (207, 205 and 206 mg/g wet wt tissue, respectively), heart (285, 287 and 284  $\mu\text{g/g}$  wet wt tissue, respectively) and kidney (91, 87 and 89  $\mu\text{g/g}$  wet wt tissue,

respectively) compared with the control group of liver (fig. 2), heart (fig. 3) and kidney (fig. 4) (206, 286 and 88  $\mu\text{g/g}$  wet wt tissue, respectively). Similarly, the methanol extract of *Michelia champaca* showed no significant change in the level of reduced glutathione (GSH) in liver, heart and kidney at a dose of 125, 250 and 500 mg/kg.



**Fig. 4: Changes in rat kidney thiobarbituric acid reactive substance (TBARS; nmol/g wet weight [wt]), superoxide dismutase (SOD; units/mg protein), catalase (CAT; units/mg protein) and reduced glutathione (GSH;  $\mu\text{g/g}$  wet wt) following oral administration of methanol extracts of *Limnophila heterophylla* and *Michelia champaca*. values were mean  $\pm$  SD (n = 6).  $p < 0.05^*$ , compared with control**

## DISCUSSION

Excessive production of reactive oxygen species (ROS) plays a very important role within the pathologic process and progression of many diseases together with completely different organs [33]. The principle for the utilization of antioxidants is well established in interference and treatment of chronic diseases wherever aerobic stress plays a serious aetiopathological role. varied population studies support that consumption of natural sources of fruits and vegetables, rich in antioxidant compounds, are related to a lower incidence of aerobic stress evoked diseases [34].

In the present investigation, the ferric reducing ability of plasma (FRAP) test measured total antioxidant capacity determined by non-enzymatic antioxidants. Numerous approaches have been established to assess the total antioxidant capacity of plasma or

serum because of the difficulty in measuring each antioxidant component separately in the serum or plasma [35]. One of these is the ferric reducing ability of plasma (FRAP), which measures the reduction of  $\text{Fe}^{3+}$  to  $\text{Fe}^{2+}$  in the presence of water-soluble exogenous antioxidants [28]. The significant increase in ferric reducing ability of plasma (FRAP) level after oral administration of methanol extract of both plants (*Limnophila heterophylla* and *Michelia champaca*) indicates the presence of bio-available antioxidants in these plants. As mentioned above, the maximum enhancement was obtained in group IV and VII after 21 d of handling. Also, ROS react with all biological substance; however, the most susceptible ones are polyunsaturated fatty acids. Reactions with these cell membrane constituents lead to lipid peroxidation (LPO) [36]. Increased LPO impairs membrane function by decreasing membrane fluidity and changing the activity of membrane-bound enzymes and receptor

[37]. Thiobarbituric acid reactive substance (TBARS) levels were measured as a marker of LPO and malondialdehyde (MDA) production. Malondialdehyde is an endogenous genotoxic product of enzymatic and ROS-induced LPO whose adducts are known to exist in DNA isolated from healthy human being [38]. In our study, the level of thiobarbituric acid reactive substance (TBARS) in the extracts treated groups decreased in a dose-dependent manner when compared to control. The present study showed the depletion in the lipid peroxidation as observed by significant decrease in the thiobarbituric acid reactive substance (TBARS) level of the liver and kidney in the plant extracts treated groups, but there was no change of thiobarbituric acid reactive substance (TBARS) level in heart as compared to control.

The superoxide dismutase (SOD) catalyzes the dismutation of superoxide to hydrogen peroxide and oxygen, thereby reducing the likelihood of superoxide anion reacting with nitric oxide to form reactive peroxynitrite [39]. It is an effective defence of the cell against the endogenous and exogenous generation of superoxide [40]. Catalase (CAT) is a ubiquitous enzyme that catalyzes the decomposition of hydrogen peroxide, a reactive oxygen species, which is a toxic product of both normal aerobic metabolism and pathogenic ROS production [41]. The ROS scavenging activity of superoxide dismutase (SOD) is effective only when it is followed by the action of catalase (CAT) and glutathione peroxidase (GPX), because of the dismutase activity of superoxide dismutase (SOD) generates hydrogen peroxide from the superoxide ion, which is more toxic than oxygen-derived free radicals and required to be scavenged further by catalase (CAT) and glutathione peroxidase (GPX) [42]. The administration of methanol extract of both plants at 500 mg/kg body weight significantly increased the level of superoxide dismutase (SOD) and catalase in liver and kidney.

This shows the antioxidant nature of both the extracts. Generally, results for the kidney have shown fewer changes in antioxidant activity compared to the liver [43]. However, decrease in the level of superoxide dismutase (SOD) and catalase (CAT) was observed in the heart, which could explain the present observation. Reduced glutathione is a protective molecule against chemical-induced cytotoxicity [44]. Glutathione is involved in many imperative cellular functions, ranging [43] from the control of physicochemical belongings of cellular proteins and peptides to the detoxification of free radicals [45]. However, long-term administration of the methanol extracts of two plants did not show significant results in reduced glutathione (GSH) levels of liver, heart and kidney indicating a protective antioxidant effect.

## CONCLUSION

It can be concluded that the methanol extracts of *Limnophila heterophylla* and *Michelia champaca* had significant *in vivo* antioxidant activity. The antioxidant action of methanol extracts of both plants may be attributed to the presence of known bioactive compounds (flavonoids), which provides maximum conjugation with radical species, thus reducing the number of free radicals available as well as oxidative stress-related ailments (diabetes, Alzheimer's disease, atherosclerosis, arthritis, cancer) of major organs such as liver, kidney and heart.

## ACKNOWLEDGEMENT

The authors are grateful to the University Grant Commission for providing financial assistance in the form of UGC-MRP fellowship to GITAM Institute of pharmacy, GITAM University, Visakhapatnam, Andhra Pradesh, India.

## AUTHORS CONTRIBUTION

The complete research work was suggested and designed by Raja S. Extraction process and *in vivo* antioxidant activity was carried out by Ravindranadh K. The manuscript was drafted by Ravindranadh K. The manuscript was edited by Ravindranadh K. Authors read and approved the final manuscript.

## CONFLICT OF INTERESTS

Declared none

## REFERENCES

- Maxwell SR. Prospects for the use of antioxidant therapies. *Drugs* 1995;49:345-61.
- Kazim H, Stephen R, Hazelrig G. Oxidative injury due to chronic nitric oxide synthase inhibition in rat: effect of regular exercise on the heart. *Biochim Biophys Acta Mol Basis Dis* 2002;1587:75-82.
- Jacob RA. The integrated antioxidant system. *Nutr Res* 1995;15:755-66.
- Rice-Evans C, Halliwell B, Lunt GG. Free Radicals and Oxidative Stress: Environment, Drugs and Food additives. 2<sup>nd</sup> edition. Portland press: London; 1995.
- Aqil F, Ahmad I, Mehmood Z. Antioxidant and free radical scavenging properties of twelve traditionally used Indian medicinal plants. *Turkish J Biol* 2006;30:177-83.
- Costantino L, Albasini A, Rastelli G, Benvenuti S. Activity of polyphenolic crude extracts as scavengers of superoxide radicals and inhibitors of xanthine oxidase. *Planta Med* 1992;5:342-4.
- Mukherjee PK. Quality control of herbal drugs-an approach to the evaluation of botanicals, Business Horizons, New Delhi, India; 2002. p. 560-7.
- Merfort I, Heilmann J, Weis M, Pietta P, Gardana C. Radical scavenger activity of three flavonoid metabolites studied by inhibition of chemiluminescence in human PMNs. *Planta Med* 1996;62:289-92.
- Arul Manikandan PN. Folk herbal medicine: a survey on the paniya tribes of the mundakunnu village of the Nilgiri Hills, South India. *Anc Sci Life* 2005;25:21-7.
- Brahmachari G, Jash SK, Mandal LC, Mondal A, Roy R. Cyclooxygenase (COX)-Inhibitory Flavonoid from *Limnophila heterophylla*. *Rasayan J Chem* 2008;1:288-91.
- Padiya RH, Patel ED, Acharya RN. Evaluation of antimicrobial activity of *Limnophila heterophylla* (Roxb.) benth. (Scrophulariaceae) whole plant. *Int J Ayur Med* 2013;4:27-3.
- Reddy GBS, Melkhani AB, Kalyani GA, Rao JV, Shirwaikar A, Kotian M, et al. Chemical and pharmacological investigations of *Limnophila conferta* and *Limnophila heterophylla*. *Int J Pharmacog* 1991;29:145-53.
- Rastogi RP, Mehrotra BN. Compendium of Indian Medicinal Plants. Eds. CDRI and NISCOM, New Delhi, India; 1998;4:435.
- Gupta S, Mehla K, Chauhan D, Nair A. Anti-inflammatory activity of leaves of *Michelia champaca* investigated on acute inflammation induced rats. *Lett Am J Pharm* 2011;30:819-22.
- Hoffmann JJ, Torrance SJ, Wiedhopf RM, Cole JR. Cytotoxic agents from *Michelia champaca* and *Talauma ovata*: Parthenolide and Costunolide. *J Pharm Sci* 1977;66:883-4.
- Vimala R, Nagarajan S, Alam M, Susan T, Joy S. Anti-inflammatory and antipyretic activity of *Michelia champaca* Linn., (white variety), *Ixora brachiata* Roxb, and *Rhynchosia cana* (Willd.) DC flower extract. *Indian J Exp Biol* 1997;35:1310-4.
- Jarald EE, Joshi SB, Jain DC. Antidiabetic activity of flower buds of *Michelia champaca* Linn, *Indian J Pharmacol* 2008;40:256.
- Takahashi M, Fuchino H, Satake M, Agatsuma Y, Sekita S. *In vitro* screening of leishmanicidal activity in myanmar timber extracts. *Biol Pharm Bull* 2004;27:921-5.
- Parimi U, Kolli D. Antibacterial and free radical scavenging activity of *Michelia champaca* Linn. Flower extracts. *Free Radicals Antioxid* 2012;2:58-61.
- Shanbhag T, Kodidela S, Shenoy S, Amuthan A, Kurra S. Effect of *Michelia champaca* linn flowers on burn wound healing in wistar rats. *Int J Pharm Sci Rev Res* 2011;7:112-5.
- Ahamad H, Mishra A, Gupta R, Saraf SA. Determination of gallic acid in *Michelia champaca* Linn. (champa) leaves and stem bark by HPTLC. *Pharm Lett* 2011;3:307-17.
- Mullaicharam AR, Kumar MS. Effect of *Michelia champaca* linn on pylorous ligated rats. *J Appl Pharm Sci* 2011;1:60-4.
- Taprial S, Kashyap D, Mehta V, Kumar S, Kumar D. Antifertility effect of hydroalcoholic leaves extract of *Michelia champaca* L.: An ethnomedicine used by Bhatra women in the Chhattisgarh state of India. *J Ethnopharmacol* 2013;147:671-5.
- Dama G, Bidkar J, Deore S, Jori M, Joshi P. Helmintholytic activity of the methanolic and aqueous extracts of leaves of *Michelia champaca*. *Res J Pharmacol Pharmacodyn* 2011;3:25-6.

25. Rajshree S, Ranjana V, *Michelia champaca* L. (Swarna champa) a review. Int J Enhanc Res Sci Tech Eng 2016;5:78-82.
26. Rajlakshmi D, Banerjee SK, Sood S, Maulik SK. *In vitro* and *in vivo* antioxidant activity of different extracts of the leaves of *Clerodendron colebrookianum* Walp in the rat. J Pharm Pharmacol 2003;55:1681-6.
27. Yesufu HB, Bassi PU, Khaz IZ, Abdulrahman FI, Mohammed GT. Phytochemical screening and hepatoprotective properties of aqueous root bark extract of *Sarcocephalus latifolius* (smith) bruce (African peach). Arch Clin Microbiol 2010;1:1-5.
28. Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power' the FRAP assay. Anal Biochem 1996;23:70-6.
29. Rai S, Wahile A, Mukherjee K, Saha BP, Mukherjee PK. Antioxidant activity of *Nelumbo nucifera* (sacred lotus) seeds. J Ethnopharmacol 2006;104:322-7.
30. Aebi H. Catalase *in vitro*. Methods Enzymol 1984;10:121-6.
31. Liu J, Edamatsu R, Kabuto H, Mori A. Antioxidant action of guilingji in the brain of rats with FeCl<sub>3</sub>-induced epilepsy. Free Radical Biol Med 1990;9:451-4.
32. Ellman GL. Tissue sulfhydryl groups. Arch Biochem Biophys 1959;82:70-7.
33. Visioli F, Keaney JF, Halliwell B. Antioxidant and cardiovascular disease; panacea or tonics for tired sheep. Cardiovascular Res 2000;47:409.
34. Vayalil PK. Antioxidant and antimutagenic properties of aqueous extract of date fruit (*Phoenix dactylifera* L, Arecaceae). J Agric Food Chem 2002;50:610-7.
35. Cao G, Prior RL. Comparison of different analytical methods for assessing the total antioxidant capacity of human serum. Clin Chem 1998;4:1309-15.
36. Bakirel T, Bakirel U, Keles OU, Ulgen SG, Yardibi H. *In vivo* assessment of antidiabetic and antioxidant activities of rosemary (*Rosmarinus officinalis*) in alloxan-diabetic rabbits. J Ethnopharmacol 2008;116:64-73.
37. Arulselvan P, Subramanian SP. Beneficial effects of *Murraya koenigii* leaves on antioxidant defense system and ultrastructural changes of pancreatic  $\beta$ -cells in experimental diabetes in rats. Chem Biol Interact 2007;165:155-64.
38. Niedernhofer LJ, Daniels JS, Rouzer CA, Greene RE, Marnett LJ. Malondialdehyde, a product of lipid peroxidation is mutagenic in human cells. J Biol Chem 2003;278:31426-33.
39. Maritim AC, Sanders RA, Watkins JB. Diabetes, oxidative stress and antioxidants: a review. J Biochem Mol Toxicol 2003;17:24-38.
40. Brawn K, Fridovich I. Superoxide radical and superoxide dismutase threat and defense. Acta Physiol Scand Suppl 1980;49:9-18.
41. Kohen R, Nyska A. Oxidation of biological systems: oxidative stress phenomena, antioxidants, redox reactions, and methods for their quantification. Toxicol Pathol 2002;30:620-50.
42. Blake DR, Allen RE, Lunee J. Free radicals in biological systems: a review oriented to the inflammatory process. Br Med Bull 1987;4:371-85.
43. Jadwiga JL, Marek M, Elzbeita B. Effect of sesquiterpene lactones on antioxidant enzymes and some drug metabolizing enzymes in rat liver and kidney. Planta Med 2000;66:199-205.
44. Orrenius S, Moldeus P. The multiple roles of glutathione in drug metabolism. Trends Pharmacol Sci 1984;5:432.
45. Meister A, Anderson ME. Glutathione. Annu Rev Biochem 1983;52:711-60.